

Cornu Spirals

From Euler to Ferrari

Max Mussavian

Arcana, May 2022

- Definitions: Curves, Lengths and Curvature
- Relating the Curve Length to its Curvature
- The Euler Spiral
- Other Fun Curves
- “Curvature determines the Curve”
- Application 1: Designing Roads and Railways
- Application 2: The Racing Line



What is a Curve?

Before we begin note that everything we look at here is in 2 dimensions \mathbb{R}^2

Definition (A Parametric Curve)

A parametric curve is a twice differentiable function that has the form $x = g(t)$ and $y = h(t)$ defined on an open interval (a, b) . The set of points traced out by the curve is called the trace.



TONBRIDGE
SCHOOL

How Long is a Curve?

Definition (Arc Length)

The length of a curve s is given by $s = \int_{t_1}^{t_2} \sqrt{x'^2 + y'^2} dt$

Intuition: The velocity at time t is $\begin{pmatrix} x'(t) \\ y'(t) \end{pmatrix}$ and hence the speed at time t is $\sqrt{x'(t)^2 + y'(t)^2}$. Distance travelled (arc length) is the integral of speed with respect to time.

Alternatively approximate the trace by line segments. The length of line segments converges to the arc length as the segments get smaller and smaller.



TONBRIDGE
SCHOOL

Some Simple Examples

- Circle: We can define a circle with radius r as $x(t) = r \cos(t)$ and $y(t) = r \sin(t)$. The arc length is

$$s = \int_0^{2\pi} \sqrt{r^2 \sin^2 t + r^2 \cos^2 t} dt = [r]_0^{2\pi} = 2\pi r$$

- Parabola: We can define a parabola as $x(t) = t$ and $y(t) = t^2$. The Arc Length between 0 and 1 is

$$\begin{aligned} s &= \int_0^1 \sqrt{1 + 4t^2} dt \\ &= \left[\frac{1}{2} t \sqrt{1 + 4t^2} + \frac{1}{4} \ln \left(2t + \sqrt{1 + 4t^2} \right) \right]_0^1 \\ &\approx 1.48 \end{aligned}$$



TONBRIDGE
SCHOOL

How Curved is a Curve?

The curvature at a point on the curve is the reciprocal of the radius of the circle that approximates the curve.

Definition (Curvature)

The curvature of curve is κ given by $\kappa = \frac{x'y'' - y'x''}{(x'^2 + y'^2)^{3/2}}$

Note: Curvature is signed in two dimensions. Positive curvature corresponds “bending to the left” while negative curvature “bends to the right”.



TONBRIDGE
SCHOOL

Explaining Definition for Curvature

For any circle with radius r we have
 $s = r\theta$.

Therefore for “kissing” circle $\frac{ds}{dt} = r \frac{d\theta}{dt}$
 But $\tan \theta = \frac{y'(t)}{x'(t)}$ and taking derivatives
 w.r.t. t we have $\sec^2 \theta \frac{d\theta}{dt} = \frac{x'y'' - y'x''}{x'^2}$.


Further $\frac{1}{\cos^2 \theta} = \frac{x'^2 + y'^2}{x'^2}$ therefore
 $\frac{d\theta}{dt} = \frac{x'y'' - y'x''}{x'^2 + y'^2}$

From above $\frac{ds}{dt} = \sqrt{x'^2 + y'^2}$.

Combining gives us

$$r = \frac{ds/dt}{d\theta/dt} = \frac{(x'^2 + y'^2)^{3/2}}{x'y'' - y'x''}$$

Finally note $\kappa = \frac{1}{r}$



TONBRIDGE
SCHOOL

Some Simple Examples

- Circle: Recall $x(t) = r \cos(t)$ and $y(t) = r \sin(t)$

$$\kappa = \frac{r^2 \sin^2 t + r^2 \cos^2 t}{(r^2 \sin^2 t + r^2 \cos^2 t)^{\frac{3}{2}}} = \frac{1}{r}$$

- Parabola: Recall $x(t) = t$ and $y(t) = t^2$.

$$\kappa = \frac{2}{(1 + 4t^2)^{\frac{3}{2}}}$$



TONBRIDGE
SCHOOL

Relating Curve Length to Curvature

Let's try the following parametrisation for x and y

$$\begin{aligned}x(t) &= \int_0^t \cos f(u) du \\y(t) &= \int_0^t \sin f(u) du\end{aligned}$$

This gives us

$$\begin{aligned}x' &= x'(t) = \cos f(t) & \text{and} & & x'' &= -f'(t) \sin f(t) \\y' &= y'(t) = \sin f(t) & \text{and} & & y'' &= f'(t) \cos f(t)\end{aligned}$$



TONBRIDGE
SCHOOL

Relating Curve Length to Curvature

This gives us the following for slope, arc length and curvature

$$\frac{dy}{dx} = \frac{\sin f(t)}{\cos f(t)} = \tan f(t)$$

$$s = \int_0^t \sqrt{\cos^2 f(u) + \sin^2 f(u)} du = t$$

$$\kappa = \frac{f'(t) \cos^2 f(t) + f'(t) \sin^2 f(t)}{\cos^2 f(t) + \sin^2 f(t)} = f'(t)$$



TONBRIDGE
SCHOOL

Define the Curve by Curve Length and Curvature

We can get replace “time” variable t by the curve length s .
And the curvature at point t is $f'(t)$. Which means

$$f(t) = \int \kappa(t) dt$$

Thus the equations for the curve become

$$\begin{aligned}x = x(s) &= \int_0^s \cos \left(\int_0^u \kappa(t) dt \right) du \\y = y(s) &= \int_0^s \sin \left(\int_0^u \kappa(t) dt \right) du\end{aligned}$$

Hence the curve is defined by curve length and curvature alone.



TONBRIDGE
SCHOOL

A Very Simple Example

We can make the curvature κ constant and equal to 1. Then $\int_0^u \kappa(t) dt = u$ and

$$\begin{aligned}x = x(s) &= \int_0^s \cos u du = \sin s \\y = y(s) &= \int_0^s \sin u du = -\cos s + 1\end{aligned}$$

which is the parametric curve for a circle with centre $(0, 1)$ and radius 1.



TONBRIDGE
SCHOOL

The Euler Spiral

A more interesting example is the curvature equal to the arc length.

$$\kappa(s) = s$$

Then $\int_0^u \kappa(t) dt = \frac{u^2}{2}$ and

$$x = x(s) = \int_0^s \cos \frac{u^2}{2} du$$

$$y = y(s) = \int_0^s \sin \frac{u^2}{2} du$$

These integrals are called “Fresnel Integrals”.

They cannot be solved analytically. But what do they look like numerically?



TONBRIDGE
SCHOOL

Making a Plotter in Python

```
def x_func(upper, lower, func):  
    def integrand(t):  
        return np.cos(func(t))  
    result, _ = integrate.quad(integrand, upper, lower)  
    return result
```

```
def y_func(upper, lower, func):  
    def integrand(t):  
        return np.sin(func(t))  
    result, _ = integrate.quad(integrand, upper, lower)  
    return result
```

Euler Spiral

```
[7] k = s  
    spy.integrate(k, s)  
  
     $\frac{s^2}{2}$ 
```

```
[8] def func_euler(x):  
    return 0.5 * np.power(x, 2)
```

```
[9] arc_length, x_coord, y_coord = x_y_coordinates(x_func, y_func, func_euler, s_steps)
```


First write functions to calculate
x and y coordinates More words
here



TONBRIDGE
SCHOOL

The Euler Spiral


Figure: The Euler Spiral aka Cornu Curve



TONBRIDGE
SCHOOL

The Euler Spiral - x y Coordinates

Figure: Fresnel Integrals with arguments $\frac{u^2}{2}$




These converge to $\pm \frac{\sqrt{\pi}}{2} \approx 0.8862$.



TONBRIDGE
SCHOOL

Other Fun Curves: Even Powers of s


Figure: $\kappa(s) = s^2$



TONBRIDGE
SCHOOL

More Fun Curves: Mix in a Bit of a Circle


Figure: $\kappa(s) = s^2 - 2.19$



TONBRIDGE
SCHOOL

More Fun Curves: Polynomials


Figure: $\kappa(s) = 5s^4 - 18s^2 + 5$



TONBRIDGE
SCHOOL

More Fun Curves: Trigonometric Functions


Figure: $\kappa(s) = \cos(s) - s \sin(s)$



TONBRIDGE
SCHOOL

More Fun Curves: Hyperbolic Functions

Figure: $\kappa(s) = \sinh(s) - 5.19$



TONBRIDGE
SCHOOL

"Curvature determines the Curve"

Say something about the fundamental theorem of plane curves...



TONBRIDGE
SCHOOL

Designing Roads and Railways

- 1 Transition curves are used to link straight sections of motorways or railways.
- 2 They are designed to give passengers a smooth ride.
- 3 In particular so sudden changes in acceleration.

Figure: Cloverleaf Motorway Interchange



Why Transition Curves are Euler Spirals

- The acceleration along the transition curve is given by

$$a = s''(t)\vec{T} + \kappa s'(t)^2\vec{N}$$

Where \vec{T} is the unit tangent vector and \vec{N} is the unit normal vector.

- If the car/train is going round the curve at constant speed $s'(t) = \text{constant}$ and $s''(t) = 0$.
- The acceleration at constant speed only depends on the curvature κ and speed $s'(t)$ in the direction of the normal vector.



TONBRIDGE
SCHOOL

Case 1: Semicircular Curves

A closed track is made up of four segments.

- 1 A straight track of length 1km
- 2 A semicircular track of length 1km. This has radius $1,000/\pi m \approx 318.31m$
- 3 A straight track of length 1km
- 4 A semicircular track of length 1km.


Let the vehicle go around the track at a constant speed of $60km/h = 16\frac{2}{3}m/s$.



TONBRIDGE
SCHOOL

Case 1: Position and Acceleration

Figure: Position and Acceleration



Note: The acceleration is a step function.

As a passenger you would feel the full centrifugal force pushing you outward the moment you entered the curve.



TONBRIDGE
SCHOOL

Case 2: Euler Spiral Transition Curves

- A closed track with same width and height.
- Replace semicircles with two parts of an Euler Spiral.
- Recall that the curvature is proportional to arc length. $\kappa = \alpha s$ for some α .
- The width and height of an Euler Spiral that turns through $\pi/2$ is given by

$$\begin{aligned} \text{width} &= \sqrt{\pi/\alpha} C(1) \\ \text{height} &= \sqrt{\pi/\alpha} S(1) \end{aligned}$$

where $S(z) = \int_0^z \sin(\pi t^2/2) dt$ and $C(z) = \int_0^z \cos(\pi t^2/2) dt$ are the standard Fresnel integrals.


$$\alpha = \frac{\pi S(1)^2}{r^2} \approx 5.95 \times 10^{-6}$$



TONBRIDGE
SCHOOL

Case 2: Euler Spiral Transition Curves

Figure: Position and Acceleration



Note: The acceleration is a increases linearly as we move through the curve.


The maximum acceleration however at the apex is greater than with the semicircular track.



TONBRIDGE
SCHOOL

Case 1 vs Case 2

Figure: Curves Compared



TONBRIDGE
SCHOOL