



# Workshop 11

COMP90051 Machine Learning


Semester 2, 2020

# Learning Outcomes

By the end of this workshop you should be able to:

1. explain why variable elimination order affects the efficiency of inference on directed PGMs
2. specify a PGM based on a natural language description
3. (extension) perform approximate inference on a PGM using PyStan

# Context for Worksheet 11a



- Pacbaby's parents are trying to teach her to discriminate between Pacmen ( $Y = 1$ ) and ghosts ( $Y = -1$ )
- She will use visual features such as presence of bowtie, hat, moustache etc., denoted by  $X_1, X_2, \dots, X_6$
- The features are *not independent*, so Pacbaby's parents decide to use a tree-augmented Naïve Bayes (TANB) model

## Q1a: TANB model


Assume all features  $\mathbf{X} = (X_1, \dots, X_6)$  are observed. What is the classification rule? Your answer should be in terms of the conditional distributions.

- Classification rule is the class that maximises the posterior probability

$$y^* = \arg \max_y p(Y = y | \mathbf{X} = \mathbf{x})$$

- Applying Bayes' rule and exploiting conditional dependence structure we have


$$p(Y = y | \mathbf{X} = \mathbf{x}) \propto p(y) p(x_1 | y) p(x_2 | x_1, y) \\ p(x_3 | x_2, y) p(x_4 | x_2, y) p(x_5 | x_4, y) p(x_6 | x_4, y)$$



## Q1b: Efficient variable elimination

Specify an efficient elimination order for the query  $p(Y|X_5 = x_5)$ . How many variables are in the biggest factor induced by variable elimination? Which variables are they?

- Recall each step of elimination:
  - \* Removes a node
  - \* Connects node's remaining neighbours
- Time complexity is **exponential** in the largest clique of the induced graph
- Different elimination orderings produce different cliques



# Q1b: Efficient variable elimination

Try eliminating in the order  $X_6 \rightarrow X_3 \rightarrow X_4 \rightarrow X_2 \rightarrow X_1$

$$\begin{aligned}
 p(Y|x_5) &\propto \sum_{X_1, X_2, X_3, X_4, X_6} p(Y)p(X_1|Y)p(X_2|X_1, Y)p(X_3|X_2, Y) \\
 &\quad p(X_4|X_2, Y)p(x_5|X_4, Y)p(X_6|X_4, Y) \\
 &= p(Y) \sum_{X_1} p(X_1|Y) \sum_{X_2} p(X_2|X_1, Y) \sum_{X_4} \underbrace{p(X_4|X_2, Y)p(x_5|X_4, Y)p(X_6|X_4, Y)}_{\phi^1(X_2, X_4, Y)} \\
 &= p(Y) \sum_{X_1} p(X_1|Y) \sum_{X_2} \underbrace{p(X_2|X_1, Y)\phi^1(X_2, X_4, Y)}_{\phi^2(X_1, X_2, Y)} \\
 &= p(Y) \sum_{X_1} \underbrace{p(X_1|Y)\phi^2(X_1, X_2, Y)}_{\phi^3(X_1, Y)} \\
 &= \phi^3(Y)
 \end{aligned}$$


# Q1b: Efficient variable elimination

Try eliminating in the order  $X_6 \rightarrow X_3 \rightarrow X_2 \rightarrow X_4 \rightarrow X_1$

$$\begin{aligned}
 p(Y|x_5) &\propto \sum_{X_1, X_2, X_3, X_4, X_6} p(Y)p(X_1|Y)p(X_2|X_1, Y)p(X_3|X_2, Y) \\
 &\quad p(X_4|X_2, Y)p(x_5|X_4, Y)p(X_6|X_4, Y) \\
 &= p(Y) \sum_{X_1} p(X_1|Y) \sum_{X_4} p(x_5|X_4, Y) \sum_{X_2} \underbrace{p(X_2|X_1, Y)p(X_4|X_2, Y)}_{\phi^1(X_1, X_2, X_4, Y)} \\
 &= p(Y) \sum_{X_1} p(X_1|Y) \sum_{X_4} \underbrace{p(x_5|X_4, Y)\phi^2(X_1, X_4, Y)}_{\phi^3(X_1, X_4, Y)} \\
 &= p(Y) \sum_{X_1} \underbrace{p(X_1|Y)\phi^4(X_1, Y)}_{\phi^5(X_1, Y)} \\
 &= \phi^6(Y)
 \end{aligned}$$

## Q1b: Efficient variable elimination






- Let's try a graphical approach.
- Re-arrange graph and moralise—add an edge between any nodes that share a child





# Q1b: Efficient variable elimination







Try eliminating in the order  $X_6 \rightarrow X_3 \rightarrow X_4 \rightarrow X_2 \rightarrow X_1$




Induced graph is same as top left. Largest clique size is 3.

# Q1b: Efficient variable elimination

Try eliminating in the order  $X_6 \rightarrow X_3 \rightarrow X_2 \rightarrow X_4 \rightarrow X_1$




Induced graph has an additional edge between  $X_1$  and  $X_4$ . Largest clique size is 4.



## Q1c: Efficient variable elimination






Specify an efficient elimination order for the query  $p(X_3 | X_5 = x_5)$ . How many variables are in the biggest factor induced by variable elimination? Which variables are they?

We'll use the graphical approach.



# Q1c: Efficient variable elimination






Try eliminating in the order  $X_6 \rightarrow X_4 \rightarrow X_1 \rightarrow Y \rightarrow X_2$




Induced graph is same as top left. Largest clique size is 3.

# Q1c: Efficient variable elimination

Try eliminating in the order  $X_6 \rightarrow X_2 \rightarrow Y \rightarrow X_4 \rightarrow X_1$





Induced graph has an additional edge between  $X_1$  and  $X_4$ . Largest clique size is 5.



## Q2a: CPTs

Use the following facts to fill out the conditional probability tables for the NB and TANB models:

- Pacbaby observes  $Y = 1$  or  $Y = -1$  50% of the time
- Given  $Y = 1$ , Pacbaby observes  $M = 1$  (moustache) 50% of the time and  $S = 1$  (sunglasses) 50% of the time
- When Pacbaby observes  $Y = -1$ , the frequency of observations are identical (equal probabilities of  $M = 1, -1$  and  $S = 1, -1$ )
- When Pacbaby observes  $Y = 1$ , anyone with a moustache wears sunglasses and anyone without a moustache does not wear sunglasses
- If  $Y = -1$  the presence/absence of a moustache has no influence on sunglasses



# Q2a: CPTs

## NB model

| $P(Y = y)$ |          |
|------------|----------|
| $y = 1$    | $y = -1$ |
|            |          |

|     | $P(M = m Y = y)$ |          |
|-----|------------------|----------|
| $y$ | $m = 1$          | $m = -1$ |
| 1   |                  |          |
| -1  |                  |          |

|     | $P(S = s Y = y)$ |          |
|-----|------------------|----------|
| $y$ | $s = 1$          | $s = -1$ |
| 1   |                  |          |
| -1  |                  |          |

## TANB model

| $P(Y = y)$ |          |
|------------|----------|
| $y = 1$    | $y = -1$ |
|            |          |

|     | $P(M = m Y = y)$ |          |
|-----|------------------|----------|
| $y$ | $m = 1$          | $m = -1$ |
| 1   |                  |          |
| -1  |                  |          |

|     |     | $P(S = s Y = y, M = m)$ |          |
|-----|-----|-------------------------|----------|
| $y$ | $m$ | $s = 1$                 | $s = -1$ |
| 1   | 1   |                         |          |
| -1  | 1   |                         |          |
| 1   | -1  |                         |          |
| -1  | -1  |                         |          |

# Q2a: CPTs

## NB model

| $P(Y = y)$ |          |
|------------|----------|
| $y = 1$    | $y = -1$ |
| 0.5        | 0.5      |

| $P(M = m Y = y)$ |         |          |
|------------------|---------|----------|
| $y$              | $m = 1$ | $m = -1$ |
| 1                | 0.5     | 0.5      |
| -1               | 0.5     | 0.5      |

| $P(S = s Y = y)$ |         |          |
|------------------|---------|----------|
| $y$              | $s = 1$ | $s = -1$ |
| 1                | 0.5     | 0.5      |
| -1               | 0.5     | 0.5      |

## TANB model

| $P(Y = y)$ |          |
|------------|----------|
| $y = 1$    | $y = -1$ |
| 0.5        | 0.5      |

| $P(M = m Y = y)$ |         |          |
|------------------|---------|----------|
| $y$              | $m = 1$ | $m = -1$ |
| 1                | 0.5     | 0.5      |
| -1               | 0.5     | 0.5      |

| $P(S = s Y = y, M = m)$ |     |         |          |
|-------------------------|-----|---------|----------|
| $y$                     | $m$ | $s = 1$ | $s = -1$ |
| 1                       | 1   | 1       | 0        |
| -1                      | 1   | 0.5     | 0.5      |
| 1                       | -1  | 0       | 1        |
| -1                      | -1  | 0.5     | 0.5      |



## Q2b: Query

Pacbaby sees someone with a moustache wearing a pair of sunglasses.

What prediction does the NB model make? What probability does it assign to its prediction?

Under the NB model

$$\begin{aligned} p(Y|M=1, S=1) &\propto p(Y)p(M=1|Y)p(S=1|Y) \\ &= \begin{cases} \left(\frac{1}{2}\right)^3, Y=1 \\ \left(\frac{1}{2}\right)^3, Y=-1 \end{cases} \end{aligned}$$

So there is a tie between the two classes.

## Q2b: Query

Pacbaby sees someone with a moustache wearing a pair of sunglasses.

What prediction does the TANB model make? What probability does it assign to its prediction?

Under the TANB model

$$\begin{aligned} p(Y|M=1, S=1) &\propto p(Y)p(M=1|Y)p(S=1|M=1, Y) \\ &= \begin{cases} \left(\frac{1}{2}\right)^2, & Y=1 \\ \left(\frac{1}{2}\right)^3, & Y=-1 \end{cases} \end{aligned}$$

Normalising we have  $p(Y=1|M=1, S=1) = \frac{2}{3}$ . So the model predicts that a Pacman was observed.

# Worksheet 11b