
TAGES ANZEIGER MATH PROBLEMS

The Zürich daily *Tages Anzeiger* has a weekly math column. Here are some problems (translated from German to English) and my solutions.

2024-05-02: Overlapping squares

Three squares with the given side lenghts overlap as follows:



The gray surface is twice as large as the black surface. What is the area of the white surface?

SOLUTION

The area of gray + white is $11^2 + 7^2 = 170$. The area of black + white is $9^2 + 5^2 = 106$

Let w be the white area. Then 170 - w = 2(106 - w) and it follows that w = 42.

2024-05-02: Interesting boxes

The width and height of a box (a rectangular parallelepiped) are two consecutive positive integers. Its depth is equal to the product of those two integers. Prove that the diagonal of such a box is always an integer.

SOLUTION

Denote the bottom face of the box by the vertices ABCD and let the corresponding vertices of the top face be WXYZ. The front face is given by ABXW.

Then, given the problem description, |AB| = x, |BX| = x + 1, |BC| = x(x + 1).

Define a diagonal AC across the bottom face, and let $d_1 = |AC|$. By the Pythagorean theorem,

$$d_1^2 = x^2 + x^2(x+1)^2 = x^4 + 2x^3 + 2x^2.$$

A diagonal of the parallelepiped is AY. Let $d_2 = |AY|$. By the Pythagorean theorem,

$$d_2^2 = d_1^2 + (x+1)^2$$

$$= x^4 + 2x^3 + 3x^2 + 2x + 1$$

$$= (x^2 + x + 1)^2$$

Since x is an integer, $d_2 = (x^2 + x + 1)$ is an integer also.

Let Q(n) be the sum of digits of a natural number n. For all $n, Q(2n) \leq 2Q(n)$.

Find a number n such that $Q(n) = 2024 \cdot Q(3n)$.

SOLUTION

Restating the problem, we know that for all n:

$$\frac{Q(2n)}{Q(n)} \le 2$$

and we want some n such that:

$$\frac{Q(3n)}{Q(n)} = \frac{1}{2024} \le 2$$

Let's use the notation $x\{k\}y$ to denote a natural number in base 10 where the digit x is repeated k times and followed by the digit y.

Playing around a bit to build intuition, for all positive k, $2(9\{k\}9) = 19\{k\}8$ and

$$\frac{Q(19\{k\}8)}{Q(9\{k\}9)} = \frac{1+9k+8}{9k+9} = 1$$

Similarly, $2(1\{k\}) = 2\{k\}$ and

$$\frac{Q(2\{k\})}{Q(1\{k\})} = 2$$

Intuitively, we want to find some increasing sequence of numbers n_i that contain a repeating pattern of digits such that $Q(n_i)$ grows much more quickly than $Q(3n_i)$ as the pattern of digits increases in length. Can we find a 3n that consists mostly of 0s while n does not?

The simplest pattern for which this is true is numbers of the form $n = 3\{k\}y$ with $y \in \{4, 5, 6\}$. Then $3n = 10\{k\}z$, where $z \in \{2, 5, 8\}$, respectively.

We may be onto something! Let's build a table:

n	$3\{k\}4$	$3\{k\}5$	$3\{k\}6$
3n	$10\{k\}2$	$10\{k\}5$	$10\{k\}8$
Q(n)	3k+4	3k + 5	3k+6
Q(3n)	3	6	9

 $\frac{3k+4}{3} = 2024$ has no integer solution (for any $m, 3m-4 \mod 3 = 2$).

 $\frac{3k+5}{6} = 2024$ has no integer solution (for any $m, 6m-5 \mod 3 = 1$).

But $\frac{3k+6}{9} = 2024$ does have an integer solution $(9m-6 \mod 3 = 0)$. k = 6070, so $n = 3\{6070\}6$ (and $3n = 10\{6070\}8$).