

# Problem Set 5

Released: March 20, 2023  
Due: March 31, 2023, 11:59pm

Submit your solutions to this homework on Canvas in a group of 2 or 3. Your solutions must be submitted in pdf produced using LaTeX.

**Definition 1.** Let  $\mathcal{C}$  be a cartesian category. A natural numbers object (NNO) in  $\mathcal{C}$  consists of

- An object  $N \in \mathcal{C}$
- Morphisms  $zero : 1 \rightarrow N$  and  $succ : N \rightarrow N$
- such that for any  $z : 1 \rightarrow V$  and  $s : V \rightarrow V$  there exists a unique morphism  $rec(z, s) : N \rightarrow V$  that satisfies
  - $rec(z, s) \circ zero = z$
  - $rec(z, s) \circ succ = s \circ rec(z, s)$

Diagrammatically,

$$\begin{array}{ccccc}
 1 & \xrightarrow{zero} & N & \xrightarrow{succ} & N \\
 & \searrow z & \downarrow rec(z,s) & & \downarrow rec(z,s) \\
 & & V & \xrightarrow{s} & V
 \end{array}$$

## Problem 1 Programming with Peano

Let  $\mathcal{C}$  be a bicartesian closed category with a natural numbers object  $(N, zero, succ)$ .

- Define a morphism  $add : N \times N \rightarrow N$  that when  $\mathcal{C}$  is the category of sets is the usual addition operation on natural numbers.
- Prove that zero is a left and right unit of add. That is

$$add \circ (zero \circ !, id_N) = id_N : N \rightarrow N$$

and

$$add \circ (id_N, zero \circ !) = id_N : N \rightarrow N$$

HINT: depending on how you define add, one of these two will be easy and one will require the uniqueness property of an NNO.

- Prove that addition is commutative:

$$\text{add} \circ (\pi_2, \pi_1) = \text{add}$$

.....

**Definition 2.** Let  $\mathcal{C}$  be a cartesian category and  $X \in \mathcal{C}_0$  an object of  $\mathcal{C}$ . An  $X$ -list object consists of

- An object  $L_X \in \mathcal{C}_0$
- Morphisms  $\text{nil} : X \rightarrow L_X$  and  $\text{cons} : X \times L_X \rightarrow L_X$
- such that for any  $n : X \rightarrow V$  and  $c : X \times V \rightarrow V$  there exists a unique  $\text{fold}(n, c) : L_X \rightarrow V$  satisfying

$$\text{fold}(n, c) \circ \text{nil} = n : 1 \rightarrow V$$

and

$$\text{fold}(n, c) \circ \text{cons} = c \circ (\pi_1, \text{fold}(n, c)) : X \times L_X \rightarrow V$$

## Problem 2 Functoriality of Lists

Assume that  $\mathcal{C}$  is a cartesian category, and for each  $X \in \mathcal{C}_0$  we have an  $X$ -list object  $(L_X, \text{nil}_X, \text{cons}_X)$ .

- Extend the list operation to a functor  $L_- : \mathcal{C} \rightarrow \mathcal{C}$ .
- Show that if  $\mathcal{C}$  is a cartesian *closed* category, the action of the functor can be *internalized* as a family of morphisms

$$\text{map} : (X \Rightarrow Y) \Rightarrow L_X \Rightarrow L_Y$$

where  $A \Rightarrow B$  is the exponential  $B^A$ .

.....