

Proof of the main theorem

Theorem

Let C be a Clifford circuit measuring commuting Pauli operators S_1, \dots, S_r . Then, for any subset of qubits L , we have

$$\text{depth}(C) \geq \frac{n_{\text{cut}}}{64|\partial L|}.$$

Corollary

For families of local-expander quantum LDPC codes of length n , a syndrome extraction circuit C implemented as a local Clifford circuit on a $\sqrt{N} \times \sqrt{N}$ grid of qubits satisfies

$$\text{depth}(C) \geq \Omega\left(\frac{n}{\sqrt{N}}\right).$$

Strategy

- Partition the circuit's qubits into two subsets L and R .

Strategy

- Partition the circuit's qubits into two subsets L and R .
- Lower bound the amount of correlation required between L and R to measure the Pauli operators.


Strategy

- Partition the circuit's qubits into two subsets L and R .
- Lower bound the amount of correlation required between L and R to measure the Pauli operators.
- Upper bound the amount of correlation introduced per operation.


Strategy

- Partition the circuit's qubits into two subsets L and R .
- Lower bound the amount of correlation required between L and R to measure the Pauli operators.
- Upper bound the amount of correlation introduced per operation.
- Combine both arguments to derive a lower bound for the depth of the circuit.

Measuring correlations




Measuring correlations



Mutual information

$$I(b_1; b_2) = 0$$

Measuring correlations




Mutual information

$$I(b_1; b_2) = 0$$

$$I(b'_1; b'_2) = 1$$


Measuring correlations




Mutual information

$$I(b'_1, b'_2, E_1; E_2) = 1$$

Measuring correlations



Measuring correlations



Classical operations can artificially boost mutual information.

Measuring correlations

- Build a circuit C' with the same action and similar depth as C by pushing all measurements and classical operations at the end.

Measuring correlations



- Build a circuit C' with the same action and similar depth as C by pushing all measurements and classical operations at the end.
- Consider the circuit $C' \circ E \circ C'$.

Measuring correlations

- ❖ Build a circuit C' with the same action and similar depth as C by pushing all measurements and classical operations at the end.
- ❖ Consider the circuit $C' \circ E \circ C'$.
- ❖ Compute the mutual information



$$I(O_L^{(2)}, E_L; O_R^{(2)}, E_R | O^{(1)}).$$

Circuit transformations





$$\text{depth}(C') \leq 4 \cdot \text{depth}(C) + 2$$

Circuit transformations





Circuit transformations







Both ancillas are the same node in the connectivity graph and in the same partition.






Circuit transformations




Circuit transformations




Circuit transformations



Circuit transformations




Circuit transformations



$$\text{depth}(U) \leq 4 \cdot \text{depth}(C)$$


Circuit transformations



$$\text{depth}(U) \leq 4 \cdot \text{depth}(C)$$

$$|\partial L| = |\partial L'|$$

The double measurement circuit



$$I(O_{\bar{L}}^{(2)}, E_{\bar{L}}; O_{\bar{R}}^{(2)}, E_{\bar{R}} | O^{(1)}) \text{ with } \bar{L} = L' \cup A_E$$

Bounds on the mutual information

Lower bound

For the double measurement circuit, we have

$$I(O_L^{(2)}, E_L; O_R^{(2)}, E_R | O^{(1)}) \geq \frac{n_{\text{cut}}}{2}.$$

Bounds on the mutual information

Lower bound

For the double measurement circuit, we have

$$I(O_L^{(2)}, E_L; O_R^{(2)}, E_R | O^{(1)}) \geq \frac{n_{\text{cut}}}{2}.$$

Upper bound

For the double measurement circuit, we have

$$I(O_L^{(2)}, E_L; O_R^{(2)}, E_R | O^{(1)}) \leq 32|\partial L|\text{depth}(C).$$

Proof of the lower bound

Note $m_i^{(t)} = \bigoplus_{o \in O_i^{(t)}} o$, the outcome for the measurement of S_i in circuit t .

Proof of the lower bound

Note $m_i^{(t)} = \bigoplus_{o \in O_i^{(t)}} o$, the outcome for the measurement of S_i in circuit t .

Note $M_{\bar{L}}^{(t)} = \{m_{i,\bar{L}}^{(t)} = \bigoplus_{o \in O_i^{(t)} \cap \bar{L}} o\}$ and similarly for $M_{\bar{R}}^{(t)}$.

Proof of the lower bound

Note $m_i^{(t)} = \bigoplus_{o \in O_i^{(t)}} o$, the outcome for the measurement of S_i in circuit t .

Note $M_L^{(t)} = \{m_{i,\bar{L}}^{(t)} = \bigoplus_{o \in O_i^{(t)} \cap \bar{L}} o\}$ and similarly for $M_{\bar{R}}^{(t)}$.

From the data processing inequality

$$I(O_L^{(2)}, E_{\bar{L}}; O_{\bar{R}}^{(2)}, E_{\bar{R}} | O^{(1)}) \geq I(M_L^{(2)}, E_{\bar{L}}; M_{\bar{R}}^{(2)}, E_{\bar{R}} | O^{(1)})$$

Proof of the lower bound

Note $m_i^{(t)} = \bigoplus_{o \in O_i^{(t)}} o$, the outcome for the measurement of S_i in circuit t .

Note $M_L^{(t)} = \{m_{i,L}^{(t)} = \bigoplus_{o \in O_i^{(t)} \cap L} o\}$ and similarly for $M_R^{(t)}$.

From the data processing inequality

$$\begin{aligned} I(O_L^{(2)}, E_L; O_R^{(2)}, E_R | O^{(1)}) &\geq I(M_L^{(2)}, E_L; M_R^{(2)}, E_R | O^{(1)}) \\ &= H(M_L^{(2)}, E_L | O^{(1)}) - H(M_L^{(2)}, E_L | M_R^{(2)}, E_R, O^{(1)}) \end{aligned}$$

Proof of the lower bound

Note $m_i^{(t)} = \bigoplus_{o \in O_i^{(t)}} o$, the outcome for the measurement of S_i in circuit t .

Note $M_L^{(t)} = \{m_{i,L}^{(t)} = \bigoplus_{o \in O_i^{(t)} \cap \bar{L}} o\}$ and similarly for $M_R^{(t)}$.

From the data processing inequality

$$\begin{aligned} I(O_L^{(2)}, E_L; O_R^{(2)}, E_R | O^{(1)}) &\geq I(M_L^{(2)}, E_L; M_R^{(2)}, E_R | O^{(1)}) \\ &= H(M_L^{(2)}, E_L | O^{(1)}) - H(M_L^{(2)}, E_L | M_R^{(2)}, E_R, O^{(1)}) \\ &= H(M_L^{(2)}, E_L, O^{(1)}) - H(O^{(1)}) - H(E_L) \end{aligned}$$

Proof of the lower bound

Note $m_i^{(t)} = \bigoplus_{o \in O_i^{(t)}} o$, the outcome for the measurement of S_i in circuit t .

Note $M_L^{(t)} = \{m_{i,L}^{(t)} = \bigoplus_{o \in O_i^{(t)} \cap \bar{L}} o\}$ and similarly for $M_R^{(t)}$.

From the data processing inequality

$$\begin{aligned} I(O_L^{(2)}, E_L; O_R^{(2)}, E_R | O^{(1)}) &\geq I(M_L^{(2)}, E_L; M_R^{(2)}, E_R | O^{(1)}) \\ &= H(M_L^{(2)}, E_L | O^{(1)}) - H(M_L^{(2)}, E_L | M_R^{(2)}, E_R, O^{(1)}) \\ &= H(M_L^{(2)}, E_L, O^{(1)}) - H(O^{(1)}) - H(E_L) \\ &= H(M_L^{(2)} | E_L, O^{(1)}). \end{aligned}$$

Proof of the lower bound

Note S_{cut} the operators with support on both L and R .

Proof of the lower bound

Note S_{cut} the operators with support on both L and R .

Note $S_{\text{cut}, \bar{L}}$ the operators $S_j \in S_{\text{cut}}$ for which m_j depends on at least one outcome in O_L .

Proof of the lower bound

Note S_{cut} the operators with support on both L and R .

Note $S_{\text{cut}, \bar{L}}$ the operators $S_i \in S_{\text{cut}}$ for which m_i depends on at least one outcome in O_L .

Note $M_{\bar{L}, \text{cut}}^{(t)}$ the outcome of $M_{\bar{L}}^{(t)}$ corresponding to $S_{\text{cut}, \bar{L}}$.

Proof of the lower bound

Note S_{cut} the operators with support on both L and R .

Note $S_{\text{cut}, \bar{L}}$ the operators $S_i \in S_{\text{cut}}$ for which m_i depends on at least one outcome in O_L .

Note $M_{\bar{L}, \text{cut}}^{(t)}$ the outcome of $M_{\bar{L}}^{(t)}$ corresponding to $S_{\text{cut}, \bar{L}}$.

From the data processing inequality

$$H(M_{\bar{L}}^{(2)} | E_{\bar{L}}, O^{(1)}) \geq H(M_{\bar{L}, \text{cut}}^{(2)} | E_{\bar{L}}, O^{(1)})$$

Proof of the lower bound

Note S_{cut} the operators with support on both L and R .

Note $S_{\text{cut}, \bar{L}}$ the operators $S_i \in S_{\text{cut}}$ for which m_i depends on at least one outcome in O_L .

Note $M_{\bar{L}, \text{cut}}^{(t)}$ the outcome of $M_{\bar{L}}^{(t)}$ corresponding to $S_{\text{cut}, \bar{L}}$.

From the data processing inequality

$$\begin{aligned} H(M_{\bar{L}}^{(2)} | E_{\bar{L}}, O^{(1)}) &\geq H(M_{\bar{L}, \text{cut}}^{(2)} | E_{\bar{L}}, O^{(1)}) \\ &\geq H(M_{\bar{L}, \text{cut}}^{(2)} | E_{\bar{L}}, O^{(1)}, M_{\bar{R}}^{(2)}) \end{aligned}$$

Proof of the lower bound

Note S_{cut} the operators with support on both L and R .

Note $S_{\text{cut},\bar{L}}$ the operators $S_i \in S_{\text{cut}}$ for which m_i depends on at least one outcome in O_L .

Note $M_{\bar{L},\text{cut}}^{(t)}$ the outcome of $M_{\bar{L}}^{(t)}$ corresponding to $S_{\text{cut},\bar{L}}$.

From the data processing inequality

$$\begin{aligned} H(M_{\bar{L}}^{(2)} | E_{\bar{L}}, O^{(1)}) &\geq H(M_{\bar{L},\text{cut}}^{(2)} | E_{\bar{L}}, O^{(1)}) \\ &\geq H(M_{\bar{L},\text{cut}}^{(2)} | E_{\bar{L}}, O^{(1)}, M_{\bar{R}}^{(2)}) \\ &= H(m_i(E_{\bar{R}}) : S_i \in S_{\text{cut},\bar{L}} | E_{\bar{L}}, O^{(1)}, M_{\bar{R}}^{(2)}) \end{aligned}$$

Proof of the lower bound

Note S_{cut} the operators with support on both L and R .

Note $S_{\text{cut},\bar{L}}$ the operators $S_i \in S_{\text{cut}}$ for which m_i depends on at least one outcome in O_L .

Note $M_{\bar{L},\text{cut}}^{(t)}$ the outcome of $M_{\bar{L}}^{(t)}$ corresponding to $S_{\text{cut},\bar{L}}$.

From the data processing inequality

$$\begin{aligned} H(M_{\bar{L}}^{(2)} | E_{\bar{L}}, O^{(1)}) &\geq H(M_{\bar{L},\text{cut}}^{(2)} | E_{\bar{L}}, O^{(1)}) \\ &\geq H(M_{\bar{L},\text{cut}}^{(2)} | E_{\bar{L}}, O^{(1)}, M_{\bar{R}}^{(2)}) \\ &= H(m_i(E_{\bar{R}}) : S_i \in S_{\text{cut},\bar{L}} | E_{\bar{L}}, O^{(1)}, M_{\bar{R}}^{(2)}) \\ &= H(m_i(E_{\bar{R}}) : S_i \in S_{\text{cut},\bar{L}}) \end{aligned}$$

Proof of the lower bound

Note S_{cut} the operators with support on both L and R .

Note $S_{\text{cut},\bar{L}}$ the operators $S_i \in S_{\text{cut}}$ for which m_i depends on at least one outcome in O_L .

Note $M_{\bar{L},\text{cut}}^{(t)}$ the outcome of $M_{\bar{L}}^{(t)}$ corresponding to $S_{\text{cut},\bar{L}}$.

From the data processing inequality


$$\begin{aligned} H(M_{\bar{L}}^{(2)} | E_{\bar{L}}, O^{(1)}) &\geq H(M_{\bar{L},\text{cut}}^{(2)} | E_{\bar{L}}, O^{(1)}) \\ &\geq H(M_{\bar{L},\text{cut}}^{(2)} | E_{\bar{L}}, O^{(1)}, M_{\bar{R}}^{(2)}) \\ &= H(m_i(E_{\bar{R}}) : S_i \in S_{\text{cut},\bar{L}} | E_{\bar{L}}, O^{(1)}, M_{\bar{R}}^{(2)}) \\ &= H(m_i(E_{\bar{R}}) : S_i \in S_{\text{cut},\bar{L}}) \\ &= |S_{\text{cut},\bar{L}}|. \end{aligned}$$

Proof of the lower bound

By symmetry

$$I(O_{\bar{L}}^{(2)}, E_{\bar{L}}; O_{\bar{R}}^{(2)}, E_{\bar{R}} | O^{(1)}) \geq \max\{|S_{\text{cut}, \bar{L}}|, |S_{\text{cut}, \bar{R}}|\} \geq \frac{n_{\text{cut}}}{2}.$$

Proof of the upper bound



$$S_{A'_2, A_E}(\rho_L(t_5); \rho_R(t_5)) = I(O_L^{(2)}, E_L; O_R^{(2)}, E_R) \geq I(O_L^{(2)}, E_L; O_R^{(2)}, E_R | O^{(1)})$$

Proof of the upper bound

- Given a set of qubits and a partition into subsets A, B . Let ρ be a density matrix on $A \cup B$ and G be a two-qubit unitary gate acting qubit of A and a qubit of B . Note $\rho' = G\rho G^\dagger$, then

$$S(\rho'_A; \rho'_B) \leq S(\rho_A, \rho_B) + 4.$$

- Single qubit gates and measurements are CPTP maps. Thus, they can't increase the mutual entropy.

Proof of the upper bound


- Given a set of qubits and a partition into subsets A, B . Let ρ be a density matrix on $A \cup B$ and G be a two-qubit unitary gate acting qubit of A and a qubit of B . Note $\rho' = G\rho G^\dagger$, then

$$S(\rho'_A; \rho'_B) \leq S(\rho_A, \rho_B) + 4.$$

- Single qubit gates and measurements are CPTP maps. Thus, they can't increase the mutual entropy.
- Discarding a subsystem can't increase the mutual entropy.


Proof of the upper bound

Proof of the upper bound




$$S(\rho_L(t_0); \rho_R(t_0)) = 0$$

Proof of the upper bound




$$S(\rho_L(t_1); \rho_R(t_1)) \leq 4\text{depth}(U)|\partial L|$$

Proof of the upper bound




$$S(\rho_L(t_2); \rho_R(t_2)) \leq 4\text{depth}(U)|\partial L|$$

Proof of the upper bound




$$S(\rho_L(t_3); \rho_R(t_3)) \leq 8\text{depth}(U)|\partial L|$$

Proof of the upper bound




$$S(\rho_L(t_4); \rho_R(t_4)) \leq 8\text{depth}(U)|\partial L|$$

Proof of the upper bound



$$S_{A'_2, A_E}(\rho_L(t_4); \rho_R(t_4)) \leq 8 \text{depth}(U) |\partial L|$$

Proof of the upper bound



$$S_{A'_2, A_E}(\rho_L(t_5); \rho_R(t_5)) \leq 8 \text{depth}(U) |\partial L|$$