

Proof of the main theorem

Theorem

Let C be a Clifford circuit measuring commuting Pauli operators S_1, \dots, S_r . Then, for any subset of qubits L , we have

$$\text{depth}(C) \geq \frac{n_{\text{cut}}}{64|\partial L|}.$$

Corollary

For families of local-expander quantum LDPC codes of length n , a syndrome extraction circuit C implemented as a local Clifford circuit on a $\sqrt{N} \times \sqrt{N}$ grid of qubits satisfies

$$\text{depth}(C) \geq \Omega\left(\frac{n}{\sqrt{N}}\right).$$

Strategy

- Partition the circuit's qubits into two subsets L and R .

Strategy

- Partition the circuit's qubits into two subsets L and R .
- Lower bound the amount of correlation required between L and R to measure the Pauli operators.


Strategy

- Partition the circuit's qubits into two subsets L and R .
- Lower bound the amount of correlation required between L and R to measure the Pauli operators.
- Upper bound the amount of correlation introduced per operation.


Strategy

- Partition the circuit's qubits into two subsets L and R .
- Lower bound the amount of correlation required between L and R to measure the Pauli operators.
- Upper bound the amount of correlation introduced per operation.
- Combine both arguments to derive a lower bound for the depth of the circuit.

Measuring correlations




Measuring correlations



Mutual information

$$I(b_1; b_2) = 0$$

Measuring correlations



Mutual information

$$I(b_1; b_2) = 0$$

$$I(b'_1; b'_2 | b_1, b_2) = 1$$