

Algebra for N00bs

By Max

Table of Contents

1. First Order Logic // you already know this
2. Words and Languages // aka easy algebra
3. Finite Deterministic Automata
4. Regular Expressions & Languages
5. Monoids
6. Metric Spaces // the secret ingredient to ML

// Plus a short note at the end about how Pin denotes functions ...

1st Order Logic

Variables: x_0, x_1, x_2, \dots

max-is-smart, max-is-dumb

1st Order Logic

Variables: x_0, x_1, x_2, \dots

max-is-smart, max-is-dumb

There are countably
many of these.

1st Order Logic

Variables: x_0, x_1, x_2, \dots

max-is-smart, max-is-dumb

There are countably
many of these.

Operators: $\sim, ^, \vee, =$

max-is-dumb \vee max-is-smart

Input: Properties
Output: Properties

1st Order Logic

Variables: x_0, x_1, x_2, \dots

There are countably many of these.

max-is-smart, max-is-dumb

Operators: $\sim, ^, \vee, =$

Input: Properties
Output: Properties

max-is-dumb \vee max-is-smart

Parentheses: (...)

1st Order Logic

Variables: x_0, x_1, x_2, \dots

There are countably many of these.

max-is-smart, max-is-dumb

Operators: $\sim, ^, \vee, =$

Input: Properties
Output: Properties

max-is-dumb \vee max-is-smart

Parentheses: (...)

Predicates: $P_i(y_0, \dots, y_n)$

Input: Variables
Output: Properties

are-siblings(Max, Sam)

1st Order Logic

Variables: x_0, x_1, x_2, \dots

There are countably many of these.

max-is-smart, max-is-dumb

Operators: $\sim, \wedge, \vee, =$

Input: Properties
Output: Properties

max-is-dumb \vee max-is-smart

Parentheses: (...)

Quantifier: \exists

$\forall x P(x)$ means $\sim \exists x \sim P(x)$

$\exists x$ jacob-thinks(x)

Predicates: $P_i(y_0, \dots, y_n)$

Input: Variables
Output: Properties

are-siblings(Max, Sam)

1st Order Logic

Variables: x_0, x_1, x_2, \dots

max-is-smart, max-is-dumb

There are countably many of these.

Operators: $\sim, \wedge, \vee, =$

max-is-dumb \vee max-is-smart

Input: Properties
Output: Properties

Parentheses: (...)

Quantifier: \exists

$\forall x P(x)$ means $\sim \exists x \sim P(x)$

$\exists x$ jacob-thinks(x)

Predicates: $P_i(y_0, \dots, y_n)$

are-siblings(Max, Sam)

Input: Variables
Output: Properties

Equality: "="

This ingredient is different in HoTT

1st Order Logic

Variables: x_0, x_1, x_2, \dots

max-is-smart, max-is-dumb

There are countably many of these.

Operators: $\sim, \wedge, \vee, =$

max-is-dumb \vee max-is-smart

Input: Properties
Output: Properties

Parentheses: (...)

Quantifier: \exists $\forall x P(x)$ means $\sim \exists x \sim P(x)$

$\exists x$ jacob-thinks(x)

Predicates: $P_i(y_0, \dots, y_n)$

are-siblings(Max, Sam)

Input: Variables
Output: Properties

Functions: $f_i(y_0, \dots, y_n) \mapsto y_k$

father-of(Max) = Frank

Input: Variables, Output: Variables

Equality: "="

This ingredient is different in HoTT

1st Order Logic

Variables: x_0, x_1, x_2, \dots

max-is-smart, max-is-dumb

There are countably many of these.

Operators: $\sim, \wedge, \vee, =$

max-is-dumb \vee max-is-smart

Input: Properties
Output: Properties

Parentheses: (...)

Quantifier: \exists $\forall x P(x)$ means $\sim \exists x \sim P(x)$

$\exists x$ jacob-thinks(x)


Predicates: $P_i(y_0, \dots, y_n)$

are-siblings(Max, Sam)

Input: Variables
Output: Properties

Theory: a finite set Γ of variables, assumed to be *true*. \vdash is entailment.

... more on this in a moment.



Functions: $f_i(y_0, \dots, y_n) \mapsto y_k$

father-of(Max) = Frank

Input: Variables, Output: Variables

Equality: "="

This ingredient is different in HoTT

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

// For example, Peano Arithmetic:

```
/* { nat(0),  
     $\forall x(x=x)$ ,  
     $\forall x,y(x=y \Rightarrow y=x)$ ,  
     $\forall x,y,z (x=y \wedge y=z) \Rightarrow x=z$ ,  
     $\forall x,y (\text{nat}(x) \wedge x=y) \Rightarrow \text{nat}(y)$ ,  
     $\forall x \text{ nat}(x) \Rightarrow \text{nat}(S(x))$ ,  
     $\forall x,y (x = y) \Leftrightarrow (S(x) = S(y))$ ,  
     $\forall x \sim(S(x)=0)$  } */
```

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

// Does there exist a proof, which assumes only Γ , and concludes φ ?

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$ // Is it an axiom?

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$ // Is it an axiom?
 - If x is in Γ it's called a non-logical axiom.

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$ // Is it an axiom?
 - If x is in Γ it's called a non-logical axiom.
 - If $\{ \} \vdash x$ then x is a "tautology", or a logical axiom.

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$ // Is it an axiom?
 - If x is in Γ it's called a non-logical axiom.
 - If $\{ \} \vdash x$ then x is a "tautology", or a logical axiom.
 - I'm not getting into logical axioms right now because I don't want to talk about modus ponens ...

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$
- If $\Gamma \vdash x$ and $\Gamma \vdash y$ then $\Gamma \vdash x \wedge y$ // Can we prove this and that?

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$
- If $\Gamma \vdash x$ and $\Gamma \vdash y$ then $\Gamma \vdash x \wedge y$
- If $\Gamma \vdash x$ or $\Gamma \vdash y$ then $\Gamma \vdash x \vee y$ // Can we prove this or that?

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$
- If $\Gamma \vdash x$ and $\Gamma \vdash y$ then $\Gamma \vdash x \wedge y$
- If $\Gamma \vdash x$ or $\Gamma \vdash y$ then $\Gamma \vdash x \vee y$
- If $\Gamma, y \vdash z$ and $\Gamma, y \vdash \sim z$ then $\Gamma \vdash \sim y$ // Proof by way of contradiction

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$
- If $\Gamma \vdash x$ and $\Gamma \vdash y$ then $\Gamma \vdash x \wedge y$
- If $\Gamma \vdash x$ or $\Gamma \vdash y$ then $\Gamma \vdash x \vee y$
- If $\Gamma, y \vdash z$ and $\Gamma, y \vdash \sim z$ then $\Gamma \vdash \sim y$

Semantic Entailment: $\Gamma \models \varphi$ if, for every theory Π containing Γ , $\Pi \not\models \sim \varphi$.

// The property is compatible with all similar theories.

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$
- If $\Gamma \vdash x$ and $\Gamma \vdash y$ then $\Gamma \vdash x \wedge y$
- If $\Gamma \vdash x$ or $\Gamma \vdash y$ then $\Gamma \vdash x \vee y$
- If $\Gamma, y \vdash z$ and $\Gamma, y \vdash \sim z$ then $\Gamma \vdash \sim y$

Semantic Entailment: $\Gamma \models \varphi$ if, for every theory Π containing Γ , $\Pi \not\models \sim \varphi$.

Consistency: $\Gamma \vdash \varphi$ if and only if $\Gamma \not\models \sim \varphi$. // Don't prove a contradiction.

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$
- If $\Gamma \vdash x$ and $\Gamma \vdash y$ then $\Gamma \vdash x \wedge y$
- If $\Gamma \vdash x$ or $\Gamma \vdash y$ then $\Gamma \vdash x \vee y$
- If $\Gamma, y \vdash z$ and $\Gamma, y \vdash \sim z$ then $\Gamma \vdash \sim y$

Semantic Entailment: $\Gamma \models \varphi$ if, for every theory Π containing Γ , $\Pi \not\models \sim \varphi$.

Consistency: $\Gamma \vdash \varphi$ if and only if $\Gamma \not\models \sim \varphi$.

Soundness: If $\Gamma \vdash \varphi$ then $\Gamma \models \varphi$. // Don't prove anything that can be disproven by a compatible theory.

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$
- If $\Gamma \vdash x$ and $\Gamma \vdash y$ then $\Gamma \vdash x \wedge y$
- If $\Gamma \vdash x$ or $\Gamma \vdash y$ then $\Gamma \vdash x \vee y$
- If $\Gamma, y \vdash z$ and $\Gamma, y \vdash \sim z$ then $\Gamma \vdash \sim y$

Semantic Entailment: $\Gamma \models \varphi$ if, for every theory Π containing Γ , $\Pi \not\models \sim \varphi$.

Consistency: $\Gamma \vdash \varphi$ if and only if $\Gamma \not\models \sim \varphi$.

Soundness: If $\Gamma \vdash \varphi$ then $\Gamma \models \varphi$.

Completeness: If $\Gamma \models \varphi$ then $\Gamma \vdash \varphi$. // All theories w/ these axioms allow φ

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$
- If $\Gamma \vdash x$ and $\Gamma \vdash y$ then $\Gamma \vdash x \wedge y$
- If $\Gamma \vdash x$ or $\Gamma \vdash y$ then $\Gamma \vdash x \vee y$
- If $\Gamma, y \vdash z$ and $\Gamma, y \vdash \sim z$ then $\Gamma \vdash \sim y$

Semantic Entailment: $\Gamma \models \varphi$ if, for every theory Π containing Γ , $\Pi \not\models \sim \varphi$.

Consistency: $\Gamma \vdash \varphi$ if and only if $\Gamma \not\models \sim \varphi$.

Soundness: If $\Gamma \vdash \varphi$ then $\Gamma \models \varphi$.

Completeness: If $\Gamma \models \varphi$ then $\Gamma \vdash \varphi$.

Thm 1: You can't have both consistency & completeness.

1st Order Logic <<Surprise Gödel>>

Theory: a finite set Γ of predicates, assumed to be *true*.

Syntactic Entailment: $\Gamma \vdash \varphi$.

- If x in Γ then $\Gamma \vdash x$
- If $\Gamma \vdash x$ and $\Gamma \vdash y$ then $\Gamma \vdash x \wedge y$
- If $\Gamma \vdash x$ or $\Gamma \vdash y$ then $\Gamma \vdash x \vee y$
- If $\Gamma, y \vdash z$ and $\Gamma, y \vdash \sim z$ then $\Gamma \vdash \sim y$

Semantic Entailment: $\Gamma \models \varphi$ if, for every theory Π containing Γ , $\Pi \not\models \sim \varphi$.

Consistency: $\Gamma \vdash \varphi$ if and only if $\Gamma \not\models \sim \varphi$.

Soundness: If $\Gamma \vdash \varphi$ then $\Gamma \models \varphi$.

Completeness: If $\Gamma \models \varphi$ then $\Gamma \vdash \varphi$.

Thm 1: You can't have both consistency & completeness.

Thm 2: Γ can't prove its own consistency.

Table of Contents

- ~~1. First Order Logic // you already know this~~
2. Words and Languages // aka easy algebra
3. Finite Deterministic Automata
4. Regular Expressions & Languages
5. Monoids
6. Metric Spaces // the secret ingredient to ML

// Plus a short note at the end about how Pin denotes functions ...

Words and Languages

Alphabet: a non-empty
finite set.

$A = \{ 0, 1, 2, \dots, 9 \}$

Words and Languages

Alphabet: a non-empty
finite set.

$A = \{ 0, 1, 2, \dots, 9 \}$

Word: a finite sequence
from the alphabet A .

69420

Words and Languages

Alphabet: a non-empty finite set.

$A = \{ 0, 1, 2, \dots, 9 \}$

Free Monoid: The set of all words over A .

$A^* = \mathbb{N}$ (with leading 0s)

Word: a finite sequence from the alphabet A .

69420

Words and Languages

Alphabet: a non-empty finite set.

$A = \{ 0, 1, 2, \dots, 9 \}$

Free Monoid: The set of all words over A .

$A^* = \mathbb{N}$ (with leading 0s)

Word: a finite sequence from the alphabet A .

69420

Alternatively, any element of A^* .

1 denotes "".

Words and Languages

Alphabet: a non-empty finite set.

$A = \{ 0, 1, 2, \dots, 9 \}$

Free Monoid: The set of all words over A .

$A^* = \mathbb{N}$ (with leading 0s)

Word: a finite sequence from the alphabet A .

69420

Alternatively, any element of A^* .

1 denotes "".

Language over A : any subset of A^*

Words and Languages

Alphabet: a non-empty finite set.

$A = \{ 0, 1, 2, \dots, 9 \}$

Free Monoid: The set of all words over A .

$A^* = \mathbb{N}$ (with leading 0s)

Word: a finite sequence from the alphabet A .

69420


Alternatively, any element of A^* .

1 denotes "".

Language over A : any subset of A^*

\mathbb{N} , positives, even numbers, odd numbers, binary, binary programs that terminate, binary programs that were written by Microsoft engineers, prime numbers, real-world RSA moduli, ternary encodings of Max's passwords ...

Words and Languages



69420

Initial

The diagram consists of a light green rectangular box containing the number 69420. The first digit, 6, is colored red, while the remaining digits (9, 4, 2, 0) are black. A thin black arrow points vertically upwards from the word 'Initial' to the red digit 6.

Words and Languages

69420

Ending



Words and Languages



69420

(Proper) prefix or left factor

Words and Languages



69420

(Proper) postfix or right factor

Words and Languages

69420

(Proper) subword

Words and Languages

$$|69420| = 5$$

Words and Languages

$$|69420| = 5$$

$$|69420|_6 = 1$$

Words and Languages

$$|69420| = 5$$

$$|69420|_6 = 1$$

$$|69666|_6 = 4$$

Words and Languages

$$|69420| = 5$$

$$|69420|_6 = 1$$

$$|69666|_6 = 4$$

W is called **multilinear** if $|w|_x \leq 1$ for all x in A

Words and Languages

$$69 * 420 = 69420$$

* is associative, non-commutative

*'s identity is 1

$$420^3 = 420420420$$

Words and Languages

$$\{69\} * \{666, 314, 420, 000\} = \{69666, 69314, 69420, 69000\}$$

* is associative, non-commutative

*'s identity is $\mathbb{1} = \{1\} = \{""\}$

$$\{22, 33\}^2 = \{2222, 2233, 3322, 3333\}$$

Words and Languages

$$\{6, 9\} \setminus \{6, 300\} = \{9\}$$

Words and Languages

$$\{6, 9\} \setminus \{6, 300\} = \{9\}$$

$$\{6, 9\} \cup \{6, 300\} = \{6, 9, 300\}$$

Words and Languages

$$\{6, 9\} \setminus \{6, 300\} = \{9\}$$

$$\{6, 9\} \cup \{6, 300\} = \{6, 9, 300\}$$


$$\{6, 9\}^c = \text{Everything in } A^* \text{ except for 6 and 9}$$

Table of Contents


- ~~1. First Order Logic // you already know this~~
- ~~2. Words and Languages // aka easy algebra~~
3. Finite Deterministic Automata
4. Regular Expressions & Languages
5. Monoids
6. Metric Spaces // the secret ingredient to ML

// Plus a short note at the end about how Pin denotes functions ...


Finite Deterministic Automata



Finite Deterministic Automata




Finite Deterministic Automata



Set.

... how Pin defines things




... how CS people usually
define things

Finite Deterministic Automata




... how Pin defines things

Set.
State Space.



... how CS people usually
define things


Finite Deterministic Automata




... how Pin defines things

... how CS people usually
define things


Finite Deterministic Automata



Finite Deterministic Automata



Finite Deterministic Automata




... how Pin defines things

Finite if Q (aka Σ) is finite.

... how CS people usually define things

Finite Deterministic Automata




... how Pin defines things

Finite if Q (aka Σ) is finite.

Deterministic if $*$ (aka δ) is a function.

... how CS people usually define things

Finite Deterministic Automata



$Q = \{ s0, s1, s2 \}$ // aka S


$A = \{ x, y, z \}$ // aka Σ

$s0 * x = s1$ // $\delta(s0, x) = s1$

$s1 * z = s0$ // $\delta(s1, z) = s0$


$s2 * y = s1$ // $\delta(s2, y) = s1$

Finite Deterministic Automata



We can extend our definition to include one or more **initial states** and **final states**.


Finite Deterministic Automata



We can extend our definition to include one or more **initial states** and **final states**.

A **path** = a sequence of consecutive transitions.


Finite Deterministic Automata



We can extend our definition to include one or more **initial states** and **final states**.

A **path** = a sequence of consecutive transitions.

Finite Deterministic Automata




We can extend our definition to include one or more **initial states** and **final states**.

A **path** = a sequence of consecutive transitions.

- “Final” if ends in a **final state**.

Finite Deterministic Automata




We can extend our definition to include one or more **initial states** and **final states**.

A **path** = a sequence of consecutive transitions.

- “Final” if ends in a **final state**.
- “Initial” if starts in an **initial state**.

Finite Deterministic Automata




We can extend our definition to include one or more **initial states** and **final states**.

A **path** = a sequence of consecutive transitions.


- “Final” if ends in a **final state**.
- “Initial” if starts in an **initial state**.
- “Successful” if both **Final** and **Initial**.

Finite Deterministic Automata



We can extend our definition to include one or more **initial states** and **final states**.

Finite Deterministic Automata



We can extend our definition to include one or more **initial states** and **final states**.

\mathcal{A} accepts the language L iff there exists initial states I and final states F s.t. L equals all of the successful paths from I to F .

$\{ y, yzx, yzxzx, yzxzxzx, \dots \}$

Table of Contents

~~1. First Order Logic // you already know this~~

~~2. Words and Languages // aka easy algebra~~

~~3. Finite Deterministic Automata~~

4. Regular Expressions & Languages

5. Monoids

6. Metric Spaces // the secret ingredient to ML

// Plus a short note at the end about how Pin denotes functions ...

Regular Expressions & Languages

Let A be a finite alphabet. $\text{Rat}(A^*)$, is the smallest subset of A^* satisfying all the following:

...

Regular Expressions & Languages

Let A be a finite alphabet. $\text{Rat}(A^*)$, is the smallest subset of A^* satisfying all the following:

- $\text{Rat}(A^*)$ contains $\{ \}$

Regular Expressions & Languages

Let A be a finite alphabet. $\text{Rat}(A^*)$, is the smallest subset of A^* satisfying all the following:

- $\text{Rat}(A^*)$ contains $\{ \}$
- If a is in A then $\{ a \}$ is in $\text{Rat}(A^*)$

Regular Expressions & Languages

Let A be a finite alphabet. $\text{Rat}(A^*)$, is the smallest subset of A^* satisfying all the following:

- $\text{Rat}(A^*)$ contains $\{ \}$
- If a is in A then $\{ a \}$ is in $\text{Rat}(A^*)$
- If X and Y are in $\text{Rat}(A^*)$ then so are XY , $X \cup Y$, and X^* .

Regular Expressions & Languages

Let A be a finite alphabet. $\text{Rat}(A^*)$, is the smallest subset of A^* satisfying all the following:

- $\text{Rat}(A^*)$ contains $\{ \}$
- If a is in A then $\{ a \}$ is in $\text{Rat}(A^*)$
- If X and Y are in $\text{Rat}(A^*)$ then so are XY , $X \cup Y$, and X^* .

Aka "Rational".

Regular Expressions & Languages

Let A be a finite alphabet. $\text{Rat}(A^*)$, is the smallest subset of A^* satisfying all the following:

- $\text{Rat}(A^*)$ contains $\{ \}$
- If a is in A then $\{ a \}$ is in $\text{Rat}(A^*)$
- If X and Y are in $\text{Rat}(A^*)$ then so are XY , $X \cup Y$, and X^* .

Aka "Rational".

Theorem: Every finite language is regular.

Regular Expressions & Languages

Let A be a finite alphabet. $\text{Rat}(A^*)$, is the smallest subset of A^* satisfying all the following:

- $\text{Rat}(A^*)$ contains $\{ \}$
- If a is in A then $\{ a \}$ is in $\text{Rat}(A^*)$
- If X and Y are in $\text{Rat}(A^*)$ then so are XY , $X \cup Y$, and X^* .

Aka "Rational".

Theorem: Every finite language is regular.

$\{ a , ab, ba \}^*$, A^*abaA^* , abA^* , set of words of even length

Regular Expressions & Languages

Let A be a finite alphabet. $\text{Rat}(A^*)$, is the smallest subset of A^* satisfying all the following:

- $\text{Rat}(A^*)$ contains $\{ \}$
- If a is in A then $\{ a \}$ is in $\text{Rat}(A^*)$
- If X and Y are in $\text{Rat}(A^*)$ then so are XY , $X \cup Y$, and X^* .

Aka "Rational".

Theorem: Every finite language is regular.

$\{ a, ab, ba \}^*$, A^*abaA^* , abA^* , set of words of even length

A^*abaA^* is the "regular expression" defining the "regular language" $\{ a_0 \dots a_k aba a_{k+1} \dots a_{k+t} \mid a_i \text{ in } A \text{ and } t, k \text{ naturals} \}$

Table of Contents

~~1. First Order Logic~~ // you already know this

~~2. Words and Languages~~ // aka easy algebra

~~3. Finite Deterministic Automata~~

~~4. Regular Expressions & Languages~~

5. Monoids

6. Metric Spaces // the secret ingredient to ML

// Plus a short note at the end about how Pin denotes functions ...


Monoids

M


$*$

1


Monoids




Monoids




Monoids



Monoids



Monoids



A^* is the “free monoid” over A , with $*$ = concatenation and $1 = ""$.

Monoids

Suppose that φ is a function ...



... s.t. $\varphi(1)=e$, and $\varphi(mk)=\varphi(m)\varphi(k)$.


Then we say φ is a **morphism**.

Table of Contents

- ~~1. First Order Logic // you already know this~~
- ~~2. Words and Languages // aka easy algebra~~
- ~~3. Finite Deterministic Automata~~
- ~~4. Regular Expressions & Languages~~
- ~~5. Monoids~~
6. Metric Spaces // the secret ingredient to ML


// Plus a short note at the end about how Pin denotes functions ...

Metric Spaces



X is a Set.


Metric Spaces



X is a Set.

$d : X \times X \rightarrow \mathbb{R}$ is a function,
Satisfying:

Metric Spaces




X is a Set.

$d : X \times X \rightarrow \mathbb{R}$ is a function,
Satisfying:

- $d(x, y) = 0$ iff $x = y$

Metric Spaces




X is a Set.

$d : X \times X \rightarrow \mathbb{R}$ is a function,
Satisfying:

- $d(x, y) = 0$ iff $x = y$
- $d(x, y) = d(y, x)$

Metric Spaces




X is a Set.

$d : X \times X \rightarrow \mathbb{R}$ is a function,
Satisfying:

- $d(x, y) = 0$ iff $x = y$
- $d(x, y) = d(y, x)$
- $d(x, y) + d(y, z) \geq d(x, z)$

Metric Spaces




X is a Set.

$d : X \times X \rightarrow \mathbb{R}$ is a function,
Satisfying:

- $d(x, y) = 0$ iff $x = y$
- $d(x, y) = d(y, x)$
- $d(x, y) + d(y, z) \geq d(x, z)$


Then d is a “metric” on X , and
 (X, d) is a “metric space”.

Metric Spaces



Consider a sequence through the space.


Metric Spaces



Consider a sequence through the space.

We can use our metric to measure the distance of every segment in the sequence.

Metric Spaces




Consider a sequence through the space.

We can use our metric to measure the distance of every segment in the sequence.

Maybe the sequence is infinite.

Metric Spaces



Consider a sequence through the space.

We can use our metric to measure the distance of every segment in the sequence.

Maybe the sequence is infinite.

If the sequence is infinite, and the distances converge to zero, then the sequence is “cauchy”.

Metric Spaces

$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \dots$

$(10, 10, 10), (10, 1, 1), (10, 0.1, 0.1),$
 $(10, 0.001, 0.001), \dots$

$3, 3.1, 3.14, 3.141, 3.1415, 3.14159, \dots$

Consider a sequence through the space.

We can use our metric to measure the distance of every segment in the sequence.


Maybe the sequence is infinite.

If the sequence is infinite, and the distances converge to zero, then the sequence is “cauchy”.

Metric Spaces

$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \dots$

$(10, 10, 10), (10, 1, 1), (10, 0.1, 0.1),$
 $(10, 0.001, 0.001), \dots$

$3, 3.1, 3.14, 3.141, 3.1415, 3.14159, \dots$  π

OBSERVE: A Cauchy sequence might *not* converge in the space.

Metric Spaces

$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \dots$

$(10, 10, 10), (10, 1, 1), (10, 0.1, 0.1),$
 $(10, 0.001, 0.001), \dots$

$3, 3.1, 3.14, 3.141, 3.1415, 3.14159, \dots$

OBSERVE: A Cauchy sequence might *not* converge in the space.

If every Cauchy sequence converges inside the space, then we say the metric space is “Complete”.

Metric Spaces

$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \dots$

$(10, 10, 10), (10, 1, 1), (10, 0.1, 0.1),$
 $(10, 0.001, 0.001), \dots$

$3, 3.1, 3.14, 3.141, 3.1415, 3.14159, \dots$

OBSERVE: A Cauchy sequence might *not* converge in the space.

If every Cauchy sequence converges inside the space, then we say the metric space is “Complete”.

Suppose (X, d) is a metric space, X is a subset of Z .

Metric Spaces

$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \dots$

$(10, 10, 10), (10, 1, 1), (10, 0.1, 0.1),$
 $(10, 0.001, 0.001), \dots$

$3, 3.1, 3.14, 3.141, 3.1415, 3.14159, \dots$

OBSERVE: A Cauchy sequence might *not* converge in the space.

If every Cauchy sequence converges inside the space, then we say the metric space is “Complete”.

Suppose (X, d) is a metric space, X is a subset of Z .

We say X is *dense* in Z if for all z in Z , either z is in X , or there exists a Cauchy sequence in X which converges to z .

Metric Spaces

1, $\frac{1}{2}$, $\frac{1}{3}$, $\frac{1}{4}$, $\frac{1}{5}$, $\frac{1}{6}$, ...

(10, 10, 1)
(10, 0.001)

3, 3.1, 3.14, ...

OBSERVE: A sequence of points in a metric space converges to a point in the space if and only if the sequence is Cauchy.

If every Cauchy sequence converges inside the space, the space is complete.

QUESTION If (X, d) is not complete, can I “fill in the holes” to make it complete?

metric space Z .
if for every $\epsilon > 0$ there exists $\delta > 0$ such that for any sequence in X , if the sequence is Cauchy, then it converges to z .

Metric Spaces

Let (X, d) be a metric space.

Metric Spaces

Let (X, d) be a metric space.

For each pair of Cauchy sequences \mathbf{z}, \mathbf{v} in the space, let

$$D(\mathbf{z}, \mathbf{v}) = \lim_{n \rightarrow \infty} d(\mathbf{z}[n], \mathbf{v}[n])$$

Metric Spaces

Let (X, d) be a metric space.

For each pair of Cauchy sequences \mathbf{z}, \mathbf{v} in the space, let

$$D(\mathbf{z}, \mathbf{v}) = \lim_{n \rightarrow \infty} d(\mathbf{z}[n], \mathbf{v}[n])$$

Let $C(X)$ = the Cauchy sequences in (X, d) , such as \mathbf{z} and \mathbf{v} .

Metric Spaces

Let (X, d) be a metric space.

For each pair of Cauchy sequences \mathbf{z}, \mathbf{v} in the space, let

$$D(\mathbf{z}, \mathbf{v}) = \lim_{n \rightarrow \infty} d(\mathbf{z}[n], \mathbf{v}[n])$$

Let $C(X)$ = the Cauchy sequences in (X, d) , such as \mathbf{z} and \mathbf{v} .

Let $\mathbf{z} \sim \mathbf{v}$ iff $D(\mathbf{z}, \mathbf{v}) = 0$.

Metric Spaces

Let (X, d) be a metric space.

For each pair of Cauchy sequences \mathbf{z}, \mathbf{v} in the space, let

$$D(\mathbf{z}, \mathbf{v}) = \lim_{n \rightarrow \infty} d(\mathbf{z}[n], \mathbf{v}[n])$$

Let $C(X)$ = the Cauchy sequences in (X, d) , such as \mathbf{z} and \mathbf{v} .

Let $\mathbf{z} \sim \mathbf{v}$ iff $D(\mathbf{z}, \mathbf{v}) = 0$. Let $[\mathbf{z}] = [\mathbf{v}] = \{ \mathbf{j} \text{ in } C(x) \mid \mathbf{j} \sim \mathbf{z} \}$.

Metric Spaces

Let (X, d) be a metric space.

For each pair of Cauchy sequences \mathbf{z}, \mathbf{v} in the space, let

$$D(\mathbf{z}, \mathbf{v}) = \lim_{n \rightarrow \infty} d(\mathbf{z}[n], \mathbf{v}[n])$$

Let $C(X)$ = the Cauchy sequences in (X, d) , such as \mathbf{z} and \mathbf{v} .

Let $\mathbf{z} \sim \mathbf{v}$ iff $D(\mathbf{z}, \mathbf{v}) = 0$. Let $[\mathbf{z}] = [\mathbf{v}] = \{ \mathbf{j} \text{ in } C(x) \mid \mathbf{j} \sim \mathbf{z} \}$.

Let $[C(X)] = \{ [\mathbf{x}] \mid \mathbf{x} \text{ in } C(x) \}$.

Metric Spaces

Let (X, d) be a metric space.

For each pair of Cauchy sequences \mathbf{z}, \mathbf{v} in the space, let

$$D(\mathbf{z}, \mathbf{v}) = \lim_{n \rightarrow \infty} d(\mathbf{z}[n], \mathbf{v}[n])$$

Let $C(X)$ = the Cauchy sequences in (X, d) , such as \mathbf{z} and \mathbf{v} .

Let $\mathbf{z} \sim \mathbf{v}$ iff $D(\mathbf{z}, \mathbf{v}) = 0$. Let $[\mathbf{z}] = [\mathbf{v}] = \{ \mathbf{j} \text{ in } C(X) \mid \mathbf{j} \sim \mathbf{z} \}$.

Let $[C(X)] = \{ [\mathbf{x}] \mid \mathbf{x} \text{ in } C(X) \}$. Let $[D]([\mathbf{z}], [\mathbf{v}]) = D(\mathbf{z}, \mathbf{v})$.

Metric Spaces

Let (X, d) be a metric space.

For each pair of Cauchy sequences \mathbf{z}, \mathbf{v} in the space, let

$$D(\mathbf{z}, \mathbf{v}) = \lim_{n \rightarrow \infty} d(\mathbf{z}[n], \mathbf{v}[n])$$

Let $C(X)$ = the Cauchy sequences in (X, d) , such as \mathbf{z} and \mathbf{v} .

Let $\mathbf{z} \sim \mathbf{v}$ iff $D(\mathbf{z}, \mathbf{v}) = 0$. Let $[\mathbf{z}] = [\mathbf{v}] = \{ \mathbf{j} \text{ in } C(X) \mid \mathbf{j} \sim \mathbf{z} \}$.

Let $[C(X)] = \{ [\mathbf{x}] \mid \mathbf{x} \text{ in } C(X) \}$. Let $[D]([\mathbf{z}], [\mathbf{v}]) = D(\mathbf{z}, \mathbf{v})$.

Then $([C(X)], [D])$ is a metric space called the “completion” of (X, d) .

Metric Spaces

Let $(X$

For each

 $D(\mathbf{z})$

Let $C(x)$

Let z

Let $[C$

Then (X, d)

space, let

n as **z** and **v**.

$$C(x) \mid \mathbf{j} \sim \mathbf{z} \}.$$
$$) = D(\mathbf{z}, \mathbf{v}) .$$

completion" of



Proof is trivial and is left to the reader

Table of Contents

- ~~1. First Order Logic // you already know this~~
- ~~2. Words and Languages // aka easy algebra~~
- ~~3. Finite Deterministic Automata~~
- ~~4. Regular Expressions & Languages~~
- ~~5. Monoids~~
- ~~6. Metric Spaces // the secret ingredient to ML~~

// Plus a short note at the end about how Π denotes functions ...

Finally, how Pin denotes functions.

Dr. Pin uses postfix notation ...

Finally, how Pin denotes functions.

Dr. Pin uses postfix notation ...

Suppose $f : X \rightarrow Y$, and $g : Y \rightarrow Z$.

Finally, how Pin denotes functions.

Dr. Pin uses postfix notation ...

Suppose $f : X \rightarrow Y$, and $g : Y \rightarrow Z$.

Then $\mathbf{g} \circ \mathbf{f} : X \rightarrow Z$

Finally, how Pin denotes functions.

Dr. Pin uses postfix notation ...

Suppose $f : X \rightarrow Y$, and $g : Y \rightarrow Z$.

Then $g \circ f : X \rightarrow Z$

$Y \rightarrow Z$ $X \rightarrow Y$

Finally, how Pin denotes functions.

Dr. Pin uses postfix notation ...

Suppose $f : X \rightarrow Y$, and $g : Y \rightarrow Z$.

Then $g \circ f : X \rightarrow Z$

$Y \rightarrow Z$ $X \rightarrow Y$

Kinda lame because the signatures don't line up.

Finally, how Pin denotes functions.

Dr. Pin uses postfix notation ...

Suppose $f : X \rightarrow Y$, and $g : Y \rightarrow Z$.

Then $g \circ f : X \rightarrow Z$

$Y \rightarrow Z$ $X \rightarrow Y$

Kinda lame because the signatures don't line up.

What if instead we wrote $x \ f \ g$?

$x \ (X \rightarrow Y) \ (Y \rightarrow Z)$

Finally, how Pin denotes functions.

Dr. Pin uses postfix notation ...

Suppose $f : X \rightarrow Y$, and $g : Y \rightarrow Z$.

Then $g \circ f : X \rightarrow Z$

$Y \rightarrow Z$ $X \rightarrow Y$

Kinda lame because the signatures don't line up.

What if instead we wrote $x \ f \ g$?

$x \ (\ X \rightarrow Y \) \ (\ Y \rightarrow Z \)$
 $X \ \text{-----}> \ Z$

Finally, how Pin denotes functions.

Dr. Pin uses postfix notation ...

Suppose $f : X \rightarrow Y$, and $g : Y \rightarrow Z$.

Then $g \circ f : X \rightarrow Z$

$Y \rightarrow Z$ $X \rightarrow Y$

Kinda lame because the signatures don't line up.

What if instead we wrote $x \ f \ g$?

$x \ (\ X \rightarrow Y \) \ (\ Y \rightarrow Z \)$
 $X \ \text{-----}> \ Z$

Much prettier!

The End.

Questions?