

# Primary Care

# Health Screening & Maintenance

| GENERAL SCREENING GUIDELINES [Evidence Grade] ( <sup>a</sup> USPSTF, <sup>b</sup> ADA, <sup>c</sup> AACE, <sup>d</sup> ACC/AHA, <sup>e</sup> ASCCP, <sup>f</sup> ACS.) |                                                                                                                                                                                                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------|----|----|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|-----------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|----|----------------------------------------------------------------------------|----------------------------------------------|----|----|-------|
| Age                                                                                                                                                                    | 18                                                                                                                                                                                                                                                                         | 19                                                                                          | 20 | 21 | 25 | 30                                                                                                                                                                                                                                                                                                                                                                                                                                    | 35 | 40                                                                                            | 45                                                                                                                       | 50 | 55                                                                         | 60                                           | 65 | 70 | 75+   |
| Cardiovascular Screening/Preventative Health Recommendations                                                                                                           |                                                                                                                                                                                                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| CVD Risk                                                                                                                                                               | Assess RFs Q4-6y [B] <sup>d</sup><br>(age, sex, Tchol/HDL, HTN, DM, CKD, smoking, FHx)                                                                                                                                                                                     |                                                                                             |    |    |    | Estimate risk w/ <a href="#">ASCVD calculator</a> Q4-6y [B] <sup>d</sup><br>If ASCVD risk ≥10% prescribe a statin* [B] <sup>a</sup> *If ASCVD risk 7.5-10% consider statin [C] <sup>a</sup> . Insufficient evidence for/against statin for age >75 [I].<br><a href="#">PREVENT</a> risk calc (30-79y) will soon replace the above. Provides 10yr and 30yr risk. Race free. Includes eGFR/UACR, BMI, A1c, and zip code (proxy for SDH) |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
|                                                                                                                                                                        |                                                                                                                                                                                                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| ASA for 1° ppx                                                                                                                                                         |                                                                                                                                                                                                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    | Consider in 40-59y w/ ASCVD risk ≥10% w/ no ↑risk of bleeding <sup>5</sup> [C] <sup>a,d</sup> |                                                                                                                          |    |                                                                            | Not rec for adults >60 [D] <sup>a</sup>      |    |    |       |
| DM                                                                                                                                                                     | If HTN or BMI ≥25 (≥23 Asian) w/ ≥1 DM RF <sup>^</sup> , or GDM [B] <sup>b</sup>                                                                                                                                                                                           |                                                                                             |    |    |    | Fasting plasma glucose or A1C [B] <sup>b</sup> Screen minimum Q3y. ↓interval if abnl result or new RF (e.g. weight gain) [C] <sup>b</sup> ✓ annually in pre-DM <sup>b</sup>                                                                                                                                                                                                                                                           |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| HTN                                                                                                                                                                    | Q3-5y; Q1y if elevated BP reading, obese, AA [A] <sup>a</sup>                                                                                                                                                                                                              |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    | Q1y [A] <sup>a</sup>                                                                          |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| HLD                                                                                                                                                                    | Insufficient evidence for screening [I] <sup>a</sup>                                                                                                                                                                                                                       | Men 20-45, women 20-55: screen Q5yr, ↑ if RF [C] <sup>c</sup><br>Q1y in DM [B] <sup>c</sup> |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    | M >45, F >55<br>Q1-2y if no RF* [A] <sup>c</sup> , Q1y in DM [B] <sup>c</sup>                 |                                                                                                                          |    |                                                                            | Screen Q1y [A] <sup>c</sup>                  |    |    |       |
| Obesity                                                                                                                                                                | Annual BMI [C] <sup>d</sup> → refer for or offer comprehensive lifestyle program if ≥25 (see <i>Weight &amp; Weight Loss</i> ) [B] <sup>d</sup>                                                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Diet                                                                                                                                                                   | Intensive behavioral counseling if CVD RF* [B] <sup>a</sup>                                                                                                                                                                                                                |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Exercise                                                                                                                                                               | 150min/wk mod.-intensity exercise or 75min/wk intense-exercise & 2d of muscle strength. activity [A] <sup>d</sup> ; fall prevention in >65 [B] <sup>a</sup>                                                                                                                |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Universal Cancer Screening                                                                                                                                             |                                                                                                                                                                                                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Colorectal CA                                                                                                                                                          | Start 10y prior to age of affected family member at dx**                                                                                                                                                                                                                   |                                                                                             |    |    |    | 45-49 needs colo Q10y, flex sig Q5y, or FIT testing Q1y [B] <sup>a</sup> ; if flex sig or FIT abnx, will need colo. If colo abnx, path determines f/u                                                                                                                                                                                                                                                                                 |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    | +/-** |
| Lung CA                                                                                                                                                                |                                                                                                                                                                                                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               | (Age 50-80) Q1y low-dose CT if 20 pack-yr or more (regardless of quitting date) or if current smoker [B] <sup>a***</sup> |    |                                                                            |                                              |    |    |       |
| Infectious Disease Screening                                                                                                                                           |                                                                                                                                                                                                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| HIV                                                                                                                                                                    | Screen all age 15-65 at least once; repeat based on risk assessment [A] <sup>a</sup>                                                                                                                                                                                       |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| HCV                                                                                                                                                                    | Screen all 18-79 at least once; repeat based on risk assessment [B] <sup>a</sup>                                                                                                                                                                                           |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| HBV                                                                                                                                                                    | Born in endemic region, getting HD or immunosuppressed, HIV+, people who inject drugs , MSM, close contact w/ HBV+ person [B] <sup>a</sup>                                                                                                                                 |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Latent TB                                                                                                                                                              | Screen if born or lived in high risk country (see USPSTF <a href="#">recs</a> ) or high risk setting (homeless, jail) [B] <sup>a</sup>                                                                                                                                     |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Syphilis                                                                                                                                                               | Screen in all individuals at increased risk of infection [A] <sup>a</sup>                                                                                                                                                                                                  |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Psych/SUD/Social Risk Factor Screening                                                                                                                                 |                                                                                                                                                                                                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Depression                                                                                                                                                             | Q1y [B] <sup>a</sup> by PHQ-2: in 2w how often (a) little interest/pleasure doing things & (b) down/depressed/hopeless ( <a href="#">JAMA 2020;323:2290</a> ). Not enough evidence to recommend for or against screening for suicide [I] <sup>a</sup>                      |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Anxiety                                                                                                                                                                | Screen regularly in adults ages 19-64, various screening tools such as GAD-2 or GAD-7 [B] <sup>a</sup>                                                                                                                                                                     |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Alcohol                                                                                                                                                                | Screen regularly with <a href="#">AUDIT-C</a> [B] <sup>a</sup>                                                                                                                                                                                                             |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Tobacco                                                                                                                                                                | Every encounter [A] <sup>a</sup> . Advise to quit, Assist doing so (plan, quit date, QuitWorks, meds), Arrange f/u. Strong evidence to start varenicline (Chantix) over other tx options, despite psych comorbidities or readiness ( <a href="#">AJRCCM. 2020;202:e5</a> ) |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Unhealthy Drug Use                                                                                                                                                     | Screening [B] <sup>a</sup> via asking questions about unhealthy drug use (not drug testing) is beneficial when services for accurate diagnosis, effective treatment, and appropriate care can be offered ( <a href="#">JAMA 2020;323:2310</a> )                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            |                                              |    |    |       |
| Inter-partner Violence                                                                                                                                                 | Screen regularly in women of reproductive age [B] <sup>a</sup> . Use <a href="#">HITS tool</a> . Assess immediate safety & consider HAVEN referral. ↑ rates in LGBT population.                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    | Consider screening for ongoing IPV, elder abuse screening [I] <sup>a</sup> |                                              |    |    |       |
| Fall Risk                                                                                                                                                              |                                                                                                                                                                                                                                                                            |                                                                                             |    |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                       |    |                                                                                               |                                                                                                                          |    |                                                                            | Yearly screening. PT. Vit D [B] <sup>a</sup> |    |    |       |

\*If ASCVD borderline: (5%-7.5%), use risk-enhancing factors [FHx, LDL-C ≥160, metabolic syndrome, CKD, S. Asian race/ethnicity, chronic inflammatory disease including HIV, menopause <40y, pre-eclampsia or pre-term delivery, abn biomarkers (e.g. hsCRP, Lp(a), apoB, ABL)] or CAC score to guide starting -statin (see *Outpatient CV Health*)

<sup>a</sup>History of previous GIB or PUD or bleeding from other sites, thrombocytopenia, coagulopathy, CKD, and concurrent use of other medications that increase bleeding risk (NSAID, steroids, DOACs, Warfarin)

<sup>d</sup>DM RFs: prior abnl testing (A1c≥5.7), FHx, AA/Latinx/Asian/NA ancestry, Hx GDM, PCOS, CVD, HTN, HDL<35 or TG>250, physical inactivity, conditions associated w/ insulin resistance (e.g. obesity)

<sup>\*\*</sup>Begin at age 40 or 10y prior to earliest dx in fam (whichever comes first) for ≥1<sup>st</sup> degree relative dx<60 or two 1<sup>st</sup> degree relatives of any age; Age 40: ≥1<sup>st</sup> degree relative dx≥60; 75+: based on life expectancy. Early screening for special populations: hx of high-grade polyp, IBD, prior colon or rectal ca, hx prior pelvic/gut radiation, genetics.

<sup>\*\*\*</sup>Consider screening cessation if limited life expectancy or screening would impede surgical candidacy.

| ADDITIONAL SCREENING GUIDELINES FOR MEN [Evidence grade] (aUSPSTF, fACS, gEndo, hPCOI) |                                                                                                                                                                                                                       |                                          |                                             |                                                          |                                                                            |    |                       |      |     |
|----------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------|---------------------------------------------|----------------------------------------------------------|----------------------------------------------------------------------------|----|-----------------------|------|-----|
| Age                                                                                    | 18-40                                                                                                                                                                                                                 | 40                                       | 45                                          | 50                                                       | 55                                                                         | 60 | 65                    | 70   | 75+ |
| AAA                                                                                    |                                                                                                                                                                                                                       |                                          |                                             |                                                          |                                                                            |    | If +tobacco hx [B]a * |      |     |
| Prostate CA**, ***                                                                     |                                                                                                                                                                                                                       | Discuss risk-benefit if<br>FHx** Q1-2ydf | Disc. risk-benefit if<br>FHx***, AA Q1-2ydf | Discuss risk-benefit if Q1-2yδ if life expectancy ≥ 10yf |                                                                            |    |                       |      |     |
|                                                                                        |                                                                                                                                                                                                                       |                                          |                                             |                                                          | PSA 55-69y if pt preference [C]a,<br><u>recommend against</u> if >70y [D]a |    |                       |      |     |
| Testicular CA                                                                          | Recommend AGAINST routine screening in all menΣ [D]a                                                                                                                                                                  |                                          |                                             |                                                          |                                                                            |    |                       |      |     |
| Anal CA#                                                                               | In MSM: anal PAP Q1y starting at 35 if HIV+ (space to Q3y after 3 neg Paps); consider Q2-3y starting at 45 if HIV.h                                                                                                   |                                          |                                             |                                                          |                                                                            |    |                       |      |     |
| Osteoporosis                                                                           |                                                                                                                                                                                                                       |                                          |                                             |                                                          | If RFag                                                                    |    |                       | DXAg |     |
| STIs                                                                                   | In MSM: Q1y, Q3-6mo if multiple/anonymous partners or sex in conjunction w/ drug use; hx re: sexual practice can guide screening (pharyngeal, anal, urine). Prescribe PrEP to patient at risk of HIV acquisition [A]a |                                          |                                             |                                                          |                                                                            |    |                       |      |     |

<sup>f</sup>Society for vascular surgery recommends screening in men and women <sup>\*\*</sup>Inform all patients of [uncertainties, risks, and potential benefits](#) before starting PSA screening

<sup>\*\*\*</sup>Discrepancy between societal guidelines, USPSTF: individual risk-benefits discussion for persons with prostates aged 55-69 [C], stop screening at age 70 [D]. ACS: for average-risk persons with prostate, start risk-benefits discussion at age 50, stop when life expectancy <10 years

<sup>6</sup>Screening frequency varies based on level. PSA < 2.5ng/mL = Q2y. PSA ≥ 2.5ng/mL = Q1y; [Amer Urological Assoc](#) recs Q2y interval

<sup>\*\*</sup>>1 first-degree relative with history of prostate cancer at early age <sup>\*\*\*</sup>First-degree relative diagnosed with prostate cancer at age <65

<sup>h</sup>Low body weight, prior fracture, smoking <sup>2</sup>If higher risk (h/o cryptorchidism or orchiopexy, testicular atrophy, or sx), evaluate, educate & consider periodic testicular exams

<sup>#</sup>Lacking evidence to inform screening initiation age & intervals. [NYS DOH A](#) includes follow-up algorithm. Stop when life expectancy <10y

# Primary Care

# Health Screening & Maintenance

## ADDITIONAL SCREENING GUIDELINES FOR NON-PREGNANT WOMEN [Evidence grade] <sup>(a)USPSTF, <sup>f</sup>ACS)</sup>

| Age                        | 18                                                                                                                                                                                          | 19 | 20 | 21                               | 25                                                                                                                                             | 30 | 35 | 40                                                                     | 45 | 50 | 55 | 60 | 65                   | 70 | 75+                                             |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|----|----------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|----|----|------------------------------------------------------------------------|----|----|----|----|----------------------|----|-------------------------------------------------|
| Breast CA*                 | Consider BRCA counseling if +FHx.<br>Screening tools <a href="#">available</a> [B] <sup>a</sup>                                                                                             |    |    |                                  |                                                                                                                                                |    |    | Q1-2y mammogram age 40-74. Stop if <10y life expectancy <sup>a,f</sup> |    |    |    |    |                      |    |                                                 |
| Cervical CA <sup>^δΣ</sup> |                                                                                                                                                                                             |    |    |                                  | Q5y 1° high risk HPV testing (preferred if available); if not available, Q5y<br>cytology + hrHPV co-testing OR Q3y cytology alone <sup>f</sup> |    |    |                                                                        |    |    |    |    |                      |    | If neg adequate<br>screening, stop <sup>a</sup> |
|                            |                                                                                                                                                                                             |    |    | Q3y cytology<br>[A] <sup>a</sup> | Q5y 1° hrHPV testing (preferred) <sup>f</sup> OR co-testing (hrHPV &<br>cytology) OR Q3y cytology alone [A] <sup>a,f</sup>                     |    |    |                                                                        |    |    |    |    |                      |    |                                                 |
| Ovarian CA                 | Recommend AGAINST routine screening in asx women [D] <sup>a</sup> ( <a href="#">Lancet 2021;397:2182</a> )                                                                                  |    |    |                                  |                                                                                                                                                |    |    |                                                                        |    |    |    |    |                      |    |                                                 |
| STIs                       | ≤24: GC/CT annually [B] <sup>a</sup>                                                                                                                                                        |    |    |                                  | Screen based on risk assessment [B] <sup>a</sup>                                                                                               |    |    |                                                                        |    |    |    |    |                      |    |                                                 |
| Contraception              | Discuss with everyone including trans ♂ (see <i>Women's Health</i> for contraception guidelines).<br>Start folic acid at reproductive age if planning/capable of pregnancy [A] <sup>a</sup> |    |    |                                  |                                                                                                                                                |    |    |                                                                        |    |    |    |    |                      |    |                                                 |
| Osteoporosis               | Consider earlier screening based on <a href="#">FRAX</a> assessment [B] <sup>a</sup> (see <i>Osteoporosis</i> )                                                                             |    |    |                                  |                                                                                                                                                |    |    |                                                                        |    |    |    |    | DXA [B] <sup>a</sup> |    |                                                 |

\*Discrepancy btw societal guidelines despite same evidence. USPSTF: 40-74 Q2y [B]; >75 no recommendation [I]. ACS: 40-44 option to start Q2y screening; 45-54 Q1y; ≥55 discuss transitioning to Q2y screening until life expectancy <10y. ACOG: offer mammography at 40; start screening at 50; discuss cessation at 75; screen Q1-2 years

<sup>a</sup>All screening methods still accepted to promote access to adequate cervical cancer screening. USPTF recommends age 21 start whereas [ACS](#) recommends age 25. [ACOG](#) recommends start at age 21 due to underscreening in younger populations.

<sup>Σ</sup>Subsequent screening dictated by age, prior screening results, and other factors. Use [ASCCP](#) for algorithm of next steps.

<sup>f</sup>Women w/ HIV require more frequent Paps & Paps preferred (HPV testing w/o cervical cytology is **not** approved). See [NIH guidelines](#)

## ADDITIONAL SCREENING GUIDELINES FOR TRANSGENDER INDIVIDUALS\*

| Age          | 18-40                                                                                                     | 40 | 45 | 50                                                                        | 55 | 60 | 65  | 70 | 75+ |
|--------------|-----------------------------------------------------------------------------------------------------------|----|----|---------------------------------------------------------------------------|----|----|-----|----|-----|
| CVD Risk     | As above; if patient has been on hormones for many years, use hormonal sex in the calculator              |    |    |                                                                           |    |    |     |    |     |
| Osteoporosis | Consider at any age if s/p gonadectomy and >5y w/o hormone tx                                             |    |    | Consider based on <a href="#">FRAX</a> (no guideline on which sex to use) |    |    | DXA |    |     |
| Breast CA    |                                                                                                           |    |    | Trans ♀: Q2y <i>after</i> 5-10y hormone use                               |    |    |     |    |     |
|              | Trans ♂: if natal tissue present (incl s/p partial mastectomy), screen per guidelines for non-trans ♀     |    |    |                                                                           |    |    |     |    |     |
| Cervical CA  | Trans ♂: if natal tissue present (incl s/p subtotal hysterectomy), screen per guidelines for non-trans ♀^ |    |    |                                                                           |    |    |     |    |     |
| Prostate CA  | Trans ♀: screen per guidelines for non-trans ♂. Use lower ULN for PSA if on estrogen                      |    |    |                                                                           |    |    |     |    |     |
| Anal CA      | Trans ♀: screen per guidelines for non-trans ♂.                                                           |    |    |                                                                           |    |    |     |    |     |

\*See Transgender Health

<sup>^</sup>Cells more likely to be "unsatisfactory" for interpretation on testosterone. On order, note that pt is on testosterone and/or amenorrheic. Self-collected vaginal swabs as sensitive as provider-collected swab ([PLoS 2018;13](#))

## VACCINES ([CDC](#), <sup>a</sup>[NYS DOH AI](#))

| Immunizations                     | 19-21                                                                                                                                                                                                                                                                                                                                                           | 22-26 | 27-49         | 50-64                                                                                                                                                                                                   | ≥65 |
|-----------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|---------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| COVID-19                          | 1 or more doses of updated (2024-2025) formula. See <a href="#">CDC guidelines</a> for unvaccinated and special populations                                                                                                                                                                                                                                     |       |               |                                                                                                                                                                                                         |     |
| Influenza                         | Annual flu vaccine<br><i>No live (intranasal) if immunocompromised (or contact), pregnant, asplenia, cochlear implant, CSF leak, ≥50; OK for 65+ to get regular adult flu vaccine if high dose not available</i>                                                                                                                                                |       |               | Annual high dose flu vaccine                                                                                                                                                                            |     |
| Tdap/Td                           | 1 dose Tdap then Td or Tdap booster every 10 years; extra Tdap dose during each pregnancy                                                                                                                                                                                                                                                                       |       |               |                                                                                                                                                                                                         |     |
| RSV                               | Seasonal administration in pregnancy                                                                                                                                                                                                                                                                                                                            |       |               | Recommended for adults 60+                                                                                                                                                                              |     |
| MMR*                              | 1-2 doses based on indication (if born 1957 or later)<br><i>1 dose in ♀ of childbearing age; 2 doses if HIV &amp; CD4≥200, healthcare personnel, or college students</i>                                                                                                                                                                                        |       |               |                                                                                                                                                                                                         |     |
| Varicella*                        | 2 doses (if born 1980 or later)                                                                                                                                                                                                                                                                                                                                 |       |               |                                                                                                                                                                                                         |     |
| Zoster*                           | Immunocompromised: 2 doses of RZV 1-2 mo apart<br><i>Administer RZV regardless of history of Zostavax (live, no longer available) administration</i>                                                                                                                                                                                                            |       |               | All: 2 doses of RZV 2-6 mo apart                                                                                                                                                                        |     |
| HPV**                             | 2-3 doses                                                                                                                                                                                                                                                                                                                                                       |       | 27-45y SDM*** |                                                                                                                                                                                                         |     |
|                                   | 3 dose series to all people with HIV up to 45y <sup>a</sup>                                                                                                                                                                                                                                                                                                     |       |               |                                                                                                                                                                                                         |     |
| PCV15/PPSV23<br>OR<br>PCV20       | Chronic heart, lung, kidney, or liver dz; DM; AUD; smoking; nephrotic syndrome; CSF leak; cochlear implant; immunocompromising conditions: PCV20 x1 OR PCV15 x1 followed by PPSV23 x1 ≥ 8 wks or 1 yr later                                                                                                                                                     |       |               | If never vaccinated, PCV20 x1 OR PCV15 x1 followed by PPSV23 x1 ≥1 yr later. If partially/fully vaccinated before age 50, may need to repeat vaccine. See <a href="#">CDC website</a> for full details. |     |
| Hep A                             | Travel to endemic country, HIV, MSM, <b>any</b> drug use (not just IVDU), chronic liver dz, undomiciled, work w/ HAV, close contact to international adoptee, settings for exposure, pregnancy at risk for infxn: 2 doses                                                                                                                                       |       |               |                                                                                                                                                                                                         |     |
| Hep B****                         | Universal vaccination for adults ages 19-59, as well as adults 60+ with risk factors for HBV (travel to endemic country, HCV, HIV, sexual exposure risk, IVDU, chronic liver dz, incarcerated, mucosal or percutaneous blood exposure (incl dialysis, DM, work exposure)<br>Recombivax HB: 3 doses (dose doubled if pre-dialysis or on HD); HEPLISAV-B: 2 doses |       |               |                                                                                                                                                                                                         |     |
| Meningococcus**<br>(MenACWY/MenB) | Functional/anatomical asplenia, HIV, ↓ complement (including by meds), travel to endemic country, freshmen in dorms, military recruits<br>1-2 doses + Q5Y if risk remains (add MenB for asplenia, ↓ complements + Q2-3Y if risk remains)                                                                                                                        |       |               |                                                                                                                                                                                                         |     |
| Hib                               | Functional or anticipated asplenia: 1 dose (if no prior); after HSCT: 3 doses (regardless of vaccination hx)                                                                                                                                                                                                                                                    |       |               |                                                                                                                                                                                                         |     |
| Monkeypox                         | Any person at risk of monkeypox (persons who are gay, bisexual, and other MSM, transgender or nonbinary people who in the past 6 mo. have had a new STD dx or >1 sexual partner): 2 doses                                                                                                                                                                       |       |               |                                                                                                                                                                                                         |     |

\*Hold live vaccines in pregnancy, malignancy, and immunocompromised (incl HIV w/ CD4 <200, cochlear implant). Ok if CD4 ≥200

\*\*Wait until after pregnancy

\*\*\*Shared decision making (SDM): most likely to benefit adults not yet exposed to HPV or w/ new sex partners

\*\*\*\*Conduct complete serologic testing (HBsAg, anti-HBs, and anti-HBc) on all adults over 18 at least once in a lifetime. Boosters not recommended for most people who have been vaccinated. For patients on HD, if anti-HBs <10 mIU/mL, a booster should be administered. For other immunocompromised people (HIV, stem-cell transplant recipients, chemotherapy) the need for booster doses is undetermined. Also consider booster in patients with anti-HBs levels <10 mIU/mL with an ongoing risk for exposure.