

Palmer amaranth (*Amaranthus palmeri*) adaptation to US Midwest agroecosystems

Maxwel C. Oliveira¹, Amit J. Jhala², Mark L. Bernards³, Chris Proctor²,
Strahinja Stepanovic², Rodrigo Werle^{1*}

¹ Department of Agronomy, University of Wisconsin-Madison, Madison, Wisconsin, United States

² Department of Agronomy and Horticulture, University of Nebraska-Lincoln, Lincoln, Nebraska, United States

³ School of Agriculture, Western Illinois University, Macomb, Illinois, United States

Correspondence*:

Rodrigo Werle

rwerle@uwisc.edu

2 ABSTRACT

Palmer amaranth (*Amaranthus palmeri* S. Watson) is one of the most troublesome agronomic weed species in the United States. Palmer amaranth is prevalent in the Southern Great Plains and Southeastern United States, and its range is expanding northward through natural dispersal and human intervention. Palmer amaranth dispersal warrants studies assessing species adaptation into new geographies. A study was conducted to investigate morphology, flowering and gender from cohorts of Palmer amaranth growing under corn, soybean, and bareground across five locations in the US Midwest. Results demonstrated that the first cohort of Palmer amaranth, established in June, produced 42% more biomass than plants from the second cohort (established in July). The first Palmer amaranth cohort produced 75.5 g plant⁻¹ in bareground, 28.3 g plant⁻¹ in soybean and 16.3 g plant⁻¹ in corn, whereas the second Palmer amaranth cohort produced 62.6, 6.3, and 1.4 g plant⁻¹ in bareground, soybean and corn, respectively. Palmer amaranth height was most impacted when growing in corn, and averaged 85.2 cm tall in the first cohort, and 38.2 cm tall in the second cohort in corn. Moreover, Palmer amaranth flowering window shifted according to crop and cohort timings. Palmer amaranth growing in intense competition, such as under low light in corn, resulted in the longest flowering window. Palmer amaranth gender was slightly influenced by day of year, weight and height. We documented a high degree of plasticity in Palmer amaranth, which will presumably favor its adaptation and expansion in cropping systems north of its current range. Therefore, preventing Palmer amaranth dispersal into new habitats is the most effective management strategy.

Keywords: Evolution, Flowering, Weed management, Pigweed

INTRODUCTION

Palmer amaranth (*Amaranthus palmeri* S. Watson) is currently ranked as one of the most economically detrimental weed species to cropping systems in the United States (Van Wychen, 2020). Unmanaged Palmer amaranth plants compete for water, light, and nutrients, which can drastically impact crop yields (Berger et al., 2015). For example, Palmer amaranth has been documented to reduce up to 91%, 68%, and

27 54% corn (Massinga et al., 2001), soybean (Klingaman and Oliver, 1994), and cotton (Morgan et al., 2001)
28 yields, respectively. Moreover, Palmer amaranth has shown a remarkable capacity to evolve resistance
29 to herbicides. To date, Palmer amaranth has evolved resistance to eight herbicide sites of action (Heap,
30 2021), increasing the weed management complexity (Lindsay et al., 2017) and posing an economical and
31 ecological risk to row-crop agriculture.

32 Palmer amaranth as a problem weed is a function of both inherent adaptations and selected management
33 practices. Palmer amaranth is a fast growing summer annual forb indigenous to the Sonoran Desert (Sauer,
34 1957). It became a serious problem weed in US agriculture in the 1990s (Ward et al., 2013). Palmer
35 amaranth weediness is likely a result of human-assisted selection combined with plant biology. Farm
36 mechanization, adoption of conservation agriculture (e.g., no-till), and intensive use of herbicides for weed
37 management are the main human-mediated selections of Palmer amaranth in cropping systems (Ward et
38 al., 2013). Palmer amaranth is a prolific seed producer with a C4 photosynthetic apparatus (Wang et al.,
39 1992). With dioecious nature, Palmer amaranth male and female plants are obligate outcrossers, increasing
40 the chances of exchanging adaptive traits among plants (Jhala et al., 2021; Oliveira et al., 2018). Also,
41 Palmer amaranth's small seeds (e.g., 1 mm) tend to thrive in no-tillage systems (Price et al., 2011), and
42 spread across locations through farm equipment (Sauer, 1972), seed mixes (Hartzler and Anderson, 2016),
43 wildlife (Farmer et al., 2017), etc., making it one of the most successful examples of weed adaptation to
44 current cropping systems.

45 Palmer amaranth's plasticity allows it to respond successfully to environmental changes. Palmer amaranth
46 demonstrates a high degree of plasticity to light, temperature, water availability, and human management
47 (Jha et al., 2010). Palmer amaranth has an extended germination period throughout the growing season
48 (Ward et al., 2013). Germination of Palmer amaranth was triggered by 18 C soil temperature at 5 cm
49 depth (Keeley et al., 1987), and optimal germination and biomass production occurred at 35/30 C day and
50 night temperatures (Guo and Al-Khatib, 2003). Palmer amaranth emergence can be influenced by tillage
51 and the use of preemergence herbicides (Chahal et al., 2021), and may result in weed germination shifts
52 within a population, as documented in *Bassia scoparia* (Sbatella and Wilson, 2010). In an experiment
53 where Palmer amaranth was subjected to continuous water stress, it survived and produced at least 14000
54 seeds plant⁻¹ (Chahal et al., 2018). Seeds from Palmer amaranth growing with limited water conditions
55 were heavier, less dormant, and prompt for germination (Matzrafi et al., 2021). Growing conditions and
56 management practices also influence Palmer amaranth sex dimorphism and flowering pattern (Korres et al.,
57 2017; Rumpa et al., 2019).

58 Palmer amaranth's current global range includes agronomic cropland in Italy (Milani et al., 2021),
59 soybean producing regions of Brazil and Argentina (Larran et al., 2017; Küpper et al., 2017), and the
60 Cotton Belt (Garetson et al., 2019; Bagavathiannan and Norsworthy, 2016) and southern Great Plains
61 of the United States Crespo et al. (2016). However, its range appears to be expanding steadily, and in
62 some cases rapidly, in the U.S. For example, in states with large Palmer amaranth infestations its range is
63 gradually pushing north as it displaces other weeds in agronomic row crops (R Werle, personal observation),
64 aided by wildlife (Farmer et al., 2017) and agricultural machinery (Ward et al., 2013). But there are other
65 examples where it has suddenly become a problem weed hundreds of miles away from known infestations.
66 New infestations in Michigan, Indiana and North Dakota appear to have resulted from Palmer amaranth
67 contaminated livestock feed. New infestations in Iowa and Minnesota were associated with planting
68 contaminated pollinator habitat seed (Yu et al., 2021). Palmer amaranth infestations have not been detected
69 in Canada, but Palmer amaranth seeds were found in sweet potato slips imported into the country (Page
70 et al., 2021). Global warming will create new opportunities for Palmer amaranth invasion. Although

71 agronomic crops in warm environments like Australia and Sub-Saharan Africa are currently at greatest risk
72 for Palmer amaranth invasion, warming temperatures will reduce barriers that may have limited Palmer
73 amaranths spread into cooler climates like Canada and Northern Europe (Kistner and Hatfield, 2018;
74 Briscoe Runquist et al., 2019).

75 There are many areas in the U.S. Corn Belt (North Central states) where Palmer amaranth is not yet
76 established, and its potential adaptability is untested. Nonetheless, the rapid expansion of Palmer amaranth
77 across the Northern U.S. is concerning and warrants investigations on its adaptability. We do know that
78 Palmer amaranth caused yield loss in Illinois soybean fields (Davis et al., 2015), but an Iowa study showed
79 that Palmer amaranth was not as well adapted as waterhemp (*Amaranthus tuberculatus*) to conditions in that
80 geography (Baker, 2021). We also know that Palmer amaranth plants that establish shortly after row-crop
81 planting have a much greater impact on crop yield than plants that emerge after the crop has produced
82 several leaves (MacRae et al., 2013). Understanding Palmer amaranth morphology and development under
83 different agroecosystems and across a wide geography can enhance our knowledge of its adaptability, and
84 may also aid in designing effective tactics to limit its range expansion and minimize its negative effects on
85 row crops. The objective of this study was to investigate the flowering pattern, gender, biomass production,
86 and height of Palmer amaranth cohorts growing in corn, soybean and bareground environments across five
87 locations in the North Central United States.

MATERIAL AND METHODS

88 Plant material and growing conditions

89 A Palmer amaranth accession (Kei3) from Perkins County, Nebraska with no glyphosate resistance was
90 selected for this study (Oliveira et al., 2021). Three weeks prior to the establishment of each cohort, seeds
91 were planted in plastic trays containing potting-mix. Emerged seedlings (1 cm) were transplanted into 200
92 cm^{-3} plastic pots (a plant pot $^{-1}$). Palmer amaranth seedlings were supplied with adequate water and kept
93 under greenhouse conditions at the University of Wisconsin-Madison, University of Nebraska-Lincoln, and
94 Western Illinois University; and kept outdoors at the Perkins extension office in Grant, NE until the 2-3
95 leaf stage (5 to 8 cm height) when they were transported to the field.

96 Field study

97 The experiment was conducted in 2018 and 2019 under field conditions at five locations: Arlington, WI
98 (43°18'N, 89°29'W), Clay Center, NE (40.57'N, 9814'W), Grant, NE (40.85'N, -101.70'W), Lincoln, NE
99 (41.16'N, 96.42'W), and Macomb, IL (40.49'N, -90.69'W).

100 Fields were conventionally tilled prior to crop planting. Corn and soybean were planted in 76-cm row
101 spacing (Table 1). Monthly mean air temperature and total precipitation were obtained using Daymet
102 weather data from June through September across the five locations in 2018 and 2019 (Correndo et al.,
103 2021) (Figure 1).

104 The field experimental units were three adjacent 9.1 m wide (12 rows at 76.2 cm row spacing) by 10.7 m
105 long. The experimental design were arranged in factorial design with three crops, two transplanting times
106 simulating two cohorts, repeated across five locations. Each field experimental unit was planted with corn,
107 soybean, or maintained as bareground. The two transplant timings were June 1 (first cohort) and July 1
108 (second cohort). Palmer amaranth seedlings (potting mix + two seedlings) were transplanted (6 cm deep
109 and 8 cm wide). Forty-eight plants were equidistantly placed (0.76 m apart) between rows within each
110 crop. After a week, one plant was eliminated and one was kept, resulting in 24 plants per experimental unit
111 and transplanting time (Figure 2). When needed, Palmer amaranth plants were supplied with water during
112 the first week after transplanting to assure seedling survival.

Table 1. Field study attributes from Arlington, WI, Clay Center, NE, Grant, NE, Lincoln, NE and Macomb, IL

Attributes	Arlington, WI	Clay Center, NE	Grant, NE	Lincoln, NE	Macomb, IL	
Bareground						
Corn	Weed control ^a Hybrid Seeding rate Weed control Stage at 1 st cohort Stage at 2 nd cohort Planting day Fertilization	S-metolachlor ^b fb glyphosate ^c / saflufenacil + imazethapyr + pyroxasulfone ^d NK0142 3120-EZ1 88956 S-metolachlor fb glyphosate V2-3 V6-7 April 30, 2018 / May 5, 2019 N (46-0-0) at 157 kg ha ⁻¹	DKC60-67 79000 S-metolachlor + trazine + mesotrione + bicyclopyrone ^e V6-7 V6-7 May 10, 2018/19	glyphosate G03C84-5122-EZ1 79000 glyphosate V6-7 V6-7 May 5, 2018 / May 14, 2019	P1197AM 86485 glyphosate + S-metolachlor ^f fb glyphosate ^g V2-3 V6-7 V6-7 April 27, 2018 / May 3, 2019 N (46-0-0) at 336 kg ha ⁻¹	glyphosate + S-metolachlor ^f fb glyphosate ^g DKC64-34RIB (2018) / G10T63-3000GT (2019) 88,000 (2018) / 79,000 (2019) glyphosate + S-metolachlor V4-5 V10-11 May 8, 2018 / April 26, 2019 UAN-32, 202 kg N ha ⁻¹
Soybean	Variety Seeding rate Weed control Stage at 1 st cohort Stage at 2 nd cohort Planting day	DSR-1950 296400 S-metolachlor / S-metolachlor V1-2 V5-6 May 5, 2018 / May 10, 2009	AG21X8 321237 saflufenacil + imazethapyr + pyroxasulfone V1-2 V5-6 May 14, 2018/19	AG28X7 220000 glyphosate V1-2 V5-6 May 5, 2018 / May 14, 2019	3017R2X (2018) / P31A22X (2019) 370650 glyphosate + S-metolachlor fb glyphosate V1-2 V5-6 May 7, 2018 / May 17, 2019	AG33X8 (2018) / CZ3601 (2019) 395000 (2018) / 370000 (2019) glyphosate + S-metolachlor V1-2 V7-8 May 8, 2018 / May 17, 2019
Soil	Type pH Organic matter (%)	Plano-silt-loam 6.6 3.5	Crete Silt Loam 6.5 3	2018: Mace silt loam / 2019: Kuma silt loam 2018: 6.1 / 2019: 5.1 2018: 2 / 2019: 1.7	Crete silty clay loam 5.2 3.4	

^aHoe weeding; ^bS-metolachlor, 1324 g ai ha⁻¹; ^cglyphosate, 1262 g ae ha⁻¹; ^dsaflufenacil + imazethapyr + pyroxasulfone, 215 g ai ha⁻¹; ^eS-metolachlor + trazine + mesotrione, + bicyclopyrone, 2409 g ai ha⁻¹; ^fS-metolachlor, 1070 g ai ha⁻¹; ^gglyphosate, 870 g ae ha⁻¹

113 After transplanting, Palmer amaranth flowering was monitored until the end of the study. When a plant
 114 flowered, the day was recorded, plant gender was identified (male or female), plant height was measured
 115 from soil surface to the top of plant. Also, aboveground plant, and all aboveground plant organs were
 116 harvested, then oven dried at 65 C until a constant weight was reached, and dry biomass (g plant⁻¹) was
 117 recorded. Plants were harvested at flowering because Palmer amaranth infestations do not occur at the
 118 Wisconsin or Illinois research locations. In our study, all locations followed the methodology of plant
 119 harvest at flowering initiation, except in Grant, NE. At the Grant, NE, location, all plants from the first
 120 cohort were harvested on July 6, 2018 or July 6, 2019, and all plants from the second cohort were harvested
 121 on August 17, 2018 or July 31, 2019, regardless of flowering status.

122 Statistical analyses

123 The statistical analyses were performed using R statistical software version 4.0.1 (R Core Team, 2021).

124 Analyses of Palmer amaranth height and biomass were performed with a linear mixed model using *lmer*
 125 function from “lme4” package (Bates et al., 2015). Plant height and biomass were log transformed to meet
 126 model assumption of normality. In the model, crop (bareground, corn, soybean) and cohort time (first and
 127 second) were the fixed effects and year nested with location were the random effects. Analysis of variance
 128 at α 0.05 was performed with *anova* function from “car” package (Fox and Weisberg, 2018). Marginal
 129 means and compact letter display were estimated with *emmeans* and *cld* from packages “emmeans” (Lenth
 130 et al., 2021) and “multcomp” (Hothorn et al., 2008), respectively.

131 Palmer amaranth cumulative flowering estimation was determined across all locations, except Grant,
 132 NE. Cumulative flowering estimation was determined using an asymmetrical three parameter log logistic
 133 Weibull model of the drc package (Ritz et al., 2015):

$$Y(x) = 0 + (d - 0)\exp(-\exp(b(\log(x) - e)))$$

134 In this model, Y is the cumulative flowering, d is the upper limit (set to 100), and e is the inflection point,
 135 and x is the day of year (doy).

136 The doy for 10, 50, and 90% cumulative flowering were determined using the *ED* function of drc package.
 137 Also, the 10, 50, and 90% Palmer amaranth cumulative flowering were compared among crops and cohorts
 138 using the *EDcomp* function of drc package. The *EDcomp* function compares the ratio of cumulative
 139 flowering using t-statistics, where P-value < 0.05 indicates that we fail to reject the null hypothesis.

140 A binary logistic regression was fitted to Palmer amaranth gender. Binary logistic regression is used for
141 predicting binary classes (Bangdiwala, 2018), such as the probability of a plant being female in a dioecious
142 species. Prior to the analysis, missing values were removed from the dataset (including all data from the
143 Grant location). The resulting dataset was split into 80% train and 20% test data. The 80% train is used
144 for model training and the 20% test is used for checking model performance on unseen dataset. Using the
145 80% train data, a generalized linear model (base R *glm* function) was fitted to binary response variable, the
146 probability of being female (0 to male and 1 to female). The independent variables were harvest day of year,
147 height, weight, and crop (without interaction). The model family was binomial with a logit function. The
148 model fit was assessed through pseudo R-squared values (McFadden, Cox and Snell, Cragg and Uhler) and
149 likelihood ratio using *nagelkerke* function from “rcompanion” package (Mangiafico, 2021). The marginal
150 effects computation was performed with Average Marginal Effects (AMEs) at every observed value of x
151 and averaged across the results (Leeper, 2017) using *margins* function from “margins” package (Leeper et
152 al., 2021). The 20% test data was predicted using the *predict* function with a cutoff estimation for male or
153 female using *performance* function from ROCR package (Sing et al., 2005). The model quality prediction
154 from the classification algorithm was measured with precision (*precision* function), recall (*recall* function)
155 and F1-score (*f_meas* function) using the “yardstick” package (Kuhn et al., 2021). The precision determines
156 the accuracy of positive predictions (female plants), recall determines the fraction of positives that were
157 correctly identified, and F1-score is a weighted harmonic mean of precision and recall with the best score
158 of 1 and the worst score of 0 (Raoniar, 2021). F1-score conveys the balance between the precision and the
159 recall (Yacoub and Axman, 2020). The area under the receiver operating curve (AUC-ROC) was also
160 estimated with *performance* function using the true positive and false positive rates. The higher the AUC,
161 better the model is at distinguishing between female and male Palmer amaranth.

RESULTS

162 Palmer amaranth height and biomass

163 Palmer amaranth plants accumulated more biomass when growing in bareground compared to plants
164 growing in soybean and corn (Figure 3A). Palmer amaranth plants in the first cohort produced 75.5, 28.3,
165 and 16.3 g plant⁻¹ in bareground, soybean and corn, respectively. Plants from the second cohort produced
166 62.6 g plant⁻¹ in bareground, followed by 6.3 g plant⁻¹ in soybean, and 1.4 g plant⁻¹ in corn.

167 Palmer amaranth height was less affected by cohort timing than was weight, with the exception of plants
168 growing in corn (Figure 3B). Plants from the first cohort were on average 69.2 cm tall in bareground, which
169 was not different from the 70.7 cm tall plants from the second cohort timing ($P = 0.74$). In addition, no
170 difference in Palmer amaranth height (69.3 cm) was detected from first cohort plants in soybean to first and
171 second cohort plants in bareground ($P > 0.75$). Palmer amaranth plants from the second cohort were nearly
172 10 cm shorter compared to the first cohort in soybeans ($P = 0.04$). The tallest (85.2 cm, first cohort) and
173 shortest (38.2 cm, second cohort) plants were observed in corn.

174 Palmer amaranth cumulative flowering

175 The initiation and duration of Palmer amaranth flowering was strongly influenced by cohort and
176 surrounding vegetation (Figure 4A, 4B). In the first cohort, floral initiation (10% flowering) occurred near
177 the end of June for all three treatments, at doy 180, 180.9, and 181.7 for soybean, bareground, and corn,
178 respectively. In the second cohort, floral initiation occurred earlier on plants growing in the bareground than
179 plants growing in soybean or corn (doy 203.8 versus doy 210.9 or 216.8, respectively). Palmer amaranth
180 growing in the bareground had the shortest duration of floral initiation (measured as the difference between
181 10% and 90% cumulative flowering) in both the first (34 days) and second (28 days) cohorts. Palmer

182 amaranth growing in soybean had a shorter duration of flowering than corn in the first cohort (40 days vs
 183 71 days), but a longer duration in the second cohort (50 vs 44 days).

184 Palmer amaranth cumulative flowering in the second cohort ranged from mid July to mid September
 185 (Figure 4B). Palmer amaranth growing in the bareground resulted in earlier flowering time compared to
 186 soybean and corn. Palmer amaranth growing in bareground reached 10%, 50%, and 90% flowering time at
 187 day 203.8, 214.4, and 232.2, respectively. Palmer amaranth growing in soybean reached 10% flowering at
 188 doy 210.9, which was 6 days prior to corn (P -value = 0.00). Similar trend was observed at 50% flowering,
 189 whereas Palmer amaranth reached 50% flowering in corn (doy 233.0) 4 days after soybeans (doy 228.9; P
 190 = 0.00). The 90% Palmer amaranth cumulative flowering occurred at same day in corn (260.9) and soybean
 191 (260.5; P = 0.66).

192 **Palmer amaranth gender**

193 The model goodness of fit was 0.23, 0.32, 0.40 using pseudo R-squared test from McFadden, Cox and
 194 Snell, and Cragg and Uhler, respectively. The likelihood ratio test showed a p-value of < 0.00. The average
 195 marginal effects showed that Palmer amaranth growing in corn resulted in 14.8% fewer females plants
 196 (Table 2). Moreover, increasing a cumulative flowering unit doy increases the probability of having a
 197 female plant by 0.4% (Table 2 and Figure 5A). A similar trend was observed for weight (Figure 5B) and
 198 height (Figure 5C), where the probability of being female increased 0.1% and 0.2% for each unit increase
 199 of weight and height, respectively (Table 2).

Table 2. Average marginal means of Palmer amaranth sex dimorphism logistic model. Factor pararamter values (crop and bareground) is shown compared to soybean.

Term	AME ^a	SE ^b	Lower	Upper	Z-score	P-value
crop_bareground	-0.048	0.054	-0.154	0.059	-0.876	0.381
crop_corn	-0.148	0.052	-0.250	-0.046	-2.842	0.004
doyh	0.004	0.001	0.003	0.006	4.959	0.000
height	0.002	0.001	0.001	0.003	2.953	0.003
weight	0.001	0.000	0.000	0.001	2.179	0.029

^a Average Marginal Effects. ^b Standard Error.

200 The model accuracy evaluation accuracy in the 20% test dataset was 0.62 with a cutoff value for female
 201 and male plants of 0.43. The model classification showed a precision of 0.64, recall of 0.66, and a F1-score
 202 of 0.65. In addition, the AUC was 0.64.

DISCUSSION

203 Our study confirmed that Palmer amaranth is well adapted to growing throughout the Midwestern United
 204 States, and is simply limited in its range by seed dispersal (Davis et al. 2015). We also confirmed Palmer
 205 amaranth's extraordinary plasticity to adapt to different agroecosystems. For example, Palmer amaranth
 206 mimicked crop architecture in competing for light (Figure 6), where plants growing in corn allocated
 207 resources to height, while plants growing in bareground allocated resources to numerous branches. Plants
 208 growing in the absence of a crop produced the greatest biomass, responding to a greater abundance of light,
 209 nutrient and water resources (Figure 3A, 3B). These results support the argument that Palmer amaranth
 210 can quickly evolve life-history traits to adapt to different cultural practices, similar to observations in a
 211 Palmer amaranth response to nitrogen study (Bravo et al., 2018). Our results highlight Palmer amaranth as

212 a threat to field crops as breeding more competitive crop varieties is likely to select more competitive weed
213 biotypes (Bravo et al., 2017).

214 Palmer amaranth growth and development in second cohort was limited due to the crop competitive
215 ability at advanced development stages. Plants were transplanted at greater crop height and width, which
216 reduced Palmer amaranth competitiveness. As a result, Palmer amaranth height and biomass was lower
217 compared to its first cohort. Moreover, Palmer amaranth growing without crop competition produced
218 the highest amounts of biomass. The Palmer amaranth strategy in bareground was to invest biomass in
219 growing plant width and height. Nonetheless, Palmer amaranth produced 17% less biomass in second
220 cohort compared to first cohort timing. In a bareground study, early emerged Palmer amaranth without
221 competition was 50% taller than late emerged plants (Webster and Grey, 2015). These results suggest that
222 crop competition is not the only factor limiting late Palmer amaranth establishment. The limited growth of
223 Palmer amaranth at second cohort is likely a reduced plant response to day length, light availability and
224 thermal units (e.g, growing degree days). The *Amaranthus* species are sensitive to photoperiod (Wu and
225 Owen, 2014). It is hypothesize that reduced day length or red/far-red effect contributed to smaller plants
226 at second cohort regardless the crop. A study in North Carolina and Illinois predicted that less than 10%
227 Palmer amaranth seedlings emergence occurred after June (Piskackova et al., 2021). In addition, Palmer
228 amaranth negative impact on soybean (Korres et al., 2020) and cotton (Webster and Grey, 2015) yields was
229 higher when plants were established near to crop planting. Therefore, early management is a key strategy
230 to minimize the damaging impact of Palmer amaranth to US Midwest cropping systems.

231 Seed production was not evaluated due to plant harvest at initiation of flowering. Nonetheless, it is well
232 documented a strong positive correlation between Palmer amaranth biomass and seed production (Schwartz
233 et al., 2016; Spaunhorst et al., 2018). In our study, plants growing from the first cohort accumulated 42%
234 more biomass when compared to the second cohort. Therefore, Palmer amaranth plants growing in the
235 second cohort is likely to produce less seeds regardless the crop. Our observation is consistent with the
236 findings that the first Palmer amaranth cohort produced 50% more seeds per plant than Palmer amaranth
237 plants established six weeks later in bareground (Webster and Grey, 2015). Still, seed production at the
238 second cohort will replenish the soil seedbank. Seed production and deposition in the seedbank is also a key
239 factor for species perpetuation (Menges, 1987). Palmer amaranth can produce a hundred thousands seeds
240 per plant (Schwartz et al., 2016; Keeley et al., 1987), which can stay viable in the soil seedbank for at least
241 36 months (Sosnoskie et al., 2013). Therefore, preventing Palmer amaranth seed production or/and seed
242 migration to non-native habitats is an essential strategy to minimize the species impact in agroecosystems.

243 An ecological approach to reduce Palmer amaranth seed production is understanding plant biology,
244 including flowering pattern. Floral initiation depends on the complex interaction between a plant's genetic
245 makeup and environmental conditions (Lang, 1965). We observed a significant disruption in flowering
246 caused by surrounding vegetation and cohort timing. Although all three treatments began flowering at the
247 same time in the first cohort (about doy 180), it took much less time for the bareground treatment to reach
248 90% flowering (34 days versus 40 for soybean and 71 for corn). The lack of competition (and stress) in the
249 bareground resulted in plants that flowered earlier, and were they not harvested, would have had longer
250 reproductive periods, thereby producing more seed. Similarly, in the second cohort, the window for all
251 plants to initiate flowering was much shorter for the bareground (28 days) compared to Palmer amaranth in
252 soybean (50 days) or corn (44 days). Significantly, plants in the bareground treatment flowered one week
253 earlier than soybean, and almost two weeks earlier than corn, again allowing for greater seed production
254 per plant had they not been harvested at initial flowering. In some cases, early flowering may confer an
255 evolutionary advantage, provided the plant has an indeterminate habit and flowering does not restrict plant

256 growth. In an Iowa study, Palmer amaranth initiated flowering two weeks prior to waterhemp (Baker, 2021),
257 and this characteristic may partially explain why Palmer amaranth can displace waterhemp where both
258 occur in the same agronomic field. In other cases, however, early flower initiation may be a response to
259 stressful conditions as a plant attempts to reproduce before running out of resources. In a dioecious species
260 like Palmer amaranth, exerting stress on plants to manipulate flowering may be beneficial in limiting seed
261 production (McFarlane et al., 2018; Schliekelman et al., 2005). For example, when growing under water
262 stress, there was a seven days flowering mismatch between male and female plants (Mesgaran et al., 2021),
263 which can minimize plant outcrossing, reduce seed production and the exchange of resistant alleles (Jhala
264 et al., 2021).

265 The mechanisms of gender-determination in plant species is intriguing and has aroused the curiosity of
266 many scientists, including Charles Darwin (Darwin, 1888). In our study, the gender model performance was
267 decent (AUC 0.64) considering the biology of plant flowering. A 1:1 male and female sex ratio is a general
268 an evolutionarily stable strategy for plant species perpetuation (Fisher, 1930). However, a slight deviation
269 from 1:1 sex ratio occurs in some dioecious species. For example, the dioecious *Halophila stipulacea*
270 is a female-biased plant in its native habitat, but naturalized *H. stipulacea* has a 1:1 ratio (Nguyen et al.,
271 2018). Naturalization of *H. stipulacea* reduced female-male ratio to expand into its non-native habitat
272 (Nguyen et al., 2018). Also, biotic and/or abiotic stress can influence plant gender determination. Palmer
273 amaranth male-to-female ratio was greater under high plant densities (Korres and Norsworthy, 2017) and
274 after herbicide application (Rumpa et al., 2019). We observed sexual dimorphism in Palmer amaranth
275 in response to surrounding vegetation and plant morphological attributes. Our model estimated that late
276 flowering, heavier and taller Palmer amaranth plants slightly deviated from 1:1 ratio in favor of female
277 plants. It was reported that female Palmer amaranth plants invested more in height, stem and biomass
278 while male plants invested more in leaf area and leaf dry weight under nutrient deficiency (Korres et al.,
279 2017). We observed more female plants in soybean and bareground compared to corn. Palmer amaranth
280 plants in the corn were more stressed by interspecific competition as evidenced by less biomass and
281 a lower weight:height ratio. Sexual dimorphism is documented in other dioecious species (Barrett and
282 Hough, 2013). For example, stronger female plant competition and greater male tolerance to herbivory was
283 reported in *Spinacia oleracea* (Pérez-Llorca and Sánchez Vilas, 2019). Research on candidate genes for
284 sex determination in *Amaranthus* species are currently underway but is far from complete (Montgomery et
285 al., 2021, 2019). Further studies are also needed to understand the ecological basis of Palmer amaranth
286 flowering, including plant behavior under climate change.

287 Our study demonstrated that Palmer amaranth is adapted to grow on arable land throughout the
288 Midwestern U.S. Palmer amaranth's range will continue to expand if current cropping practices are
289 continued. As waterhemp and Palmer amaranth begin to share the same habitat, it will increase weed
290 management complexity. Preventing Palmer amaranth seed dispersal must be a priority. Regional
291 collaboration is necessary to slow the spread of this aggressive and adaptable weed. Where Palmer
292 amaranth occurs, management tactics should focus on limiting Palmer amaranth establishment until row
293 crops can shade late-emerging plants. Increasing the diversity of crops in rotation, varying row crop planting
294 date, narrowing row width, and increasing residue cover through the use of cover crops are all tactics
295 that can minimize the growth and seed production of Palmer amaranth, and improve the sustainability of
296 Midwestern cropping systems.

DISCLOSURE/CONFLICT-OF-INTEREST STATEMENT

297 The authors declare that the research was conducted in the absence of any commercial or financial
298 relationships that could be construed as a potential conflict of interest.

DATA ACCESSIBILITY

299 The data and scripts used to analyse the data presented in this work can be found at Zenodo (Oliveira,
300 2021).

AUTHOR CONTRIBUTIONS

301 RW and MO: designed the experiments; AJ, CP, MB, MO, and SS: conducted the experiments; MO:
302 analyzed the data; MO: wrote the manuscript and all authors revised; AJ, CP, MB, MO, SS, and RW:
303 conceptualized the research. All authors reviewed the manuscript.

ACKNOWLEDGMENTS

304 Funding: This work received no specific grant from any funding agency, commercial, or not-for-profit
305 sectors.

REFERENCES

- 306 Bagavathiannan, M. V., and Norsworthy, J. K. (2016). Multiple-Herbicide Resistance Is Widespread in
307 Roadside Palmer Amaranth Populations. *PLOS ONE* 11, e0148748. doi:10.1371/journal.pone.0148748.
- 308 Baker, R. (2021). Comparative analysis of Palmer amaranth (*Amaranthus palmeri*) and waterhemp (*A.
309 Tuberculatus*) in Iowa. doi:10.31274/etd-20210609-11.
- 310 Bangdiwala, S. I. (2018). Regression: Binary logistic. *International Journal of Injury Control and Safety
311 Promotion* 25, 336–338. doi:10.1080/17457300.2018.1486503.
- 312 Barrett, S. C. H., and Hough, J. (2013). Sexual dimorphism in flowering plants. *Journal of Experimental
313 Botany* 64, 67–82. doi:10.1093/jxb/ers308.
- 314 Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting Linear Mixed-Effects Models Using
315 Lme4. *Journal of Statistical Software* 67, 1–48. doi:10.18637/jss.v067.i01.
- 316 Berger, S. T., Ferrell, J. A., Rowland, D. L., and Webster, T. M. (2015). Palmer Amaranth (*Amaranthus
317 palmeri*) Competition for Water in Cotton. *Weed Science* 63, 928–935. doi:10.1614/WS-D-15-00062.1.
- 318 Bravo, W., Leon, R. G., Ferrell, J. A., Mulvaney, M. J., and Wood, C. W. (2017). Differentiation of
319 Life-History Traits among Palmer Amaranth Populations (*Amaranthus palmeri*) and Its Relation to
320 Cropping Systems and Glyphosate Sensitivity. *Weed Science* 65, 339–349. doi:10.1017/wsc.2017.14.
- 321 Bravo, W., Leon, R. G., Ferrell, J. A., Mulvaney, M. J., and Wood, C. W. (2018). Evolutionary Adaptations
322 of Palmer Amaranth (*Amaranthus palmeri*) to Nitrogen Fertilization and Crop Rotation History Affect
323 Morphology and Nutrient-Use Efficiency. *Weed Science* 66, 180–189. doi:10.1017/wsc.2017.73.
- 324 Briscoe Runquist, R. D., Lake, T., Tiffin, P., and Moeller, D. A. (2019). Species distribution models
325 throughout the invasion history of Palmer amaranth predict regions at risk of future invasion and reveal
326 challenges with modeling rapidly shifting geographic ranges. *Sci Rep* 9, 2426. doi:10.1038/s41598-
327 018-38054-9.
- 328 Chahal, P. S., Barnes, E. R., and Jhala, A. J. (2021). Emergence pattern of Palmer amaranth (*Amaranthus
329 palmeri*) influenced by tillage timings and residual herbicides. *Weed Technology* 35, 433–439.
330 doi:10.1017/wet.2020.136.

- 331 Chahal, P. S., Irmak, S., Jugulam, M., and Jhala, A. J. (2018). Evaluating Effect of Degree of Water Stress
332 on Growth and Fecundity of Palmer amaranth (*Amaranthus palmeri*) Using Soil Moisture Sensors.
333 *Weed Science* 66, 738–745. doi:10.1017/wsc.2018.47.
- 334 Correndo, A. A., Moro Rosso, L. H., and Ciampitti, I. A. (2021). Retrieving and processing agro-
335 meteorological data from API-client sources using R software. *BMC Research Notes* 14, 205.
336 doi:10.1186/s13104-021-05622-8.
- 337 Crespo, R. J., Wingeier, A. B., Borman, C. J., and Bernards, M. L. (2016). Baseline Sensitivity of
338 Nebraska Waterhemp and Palmer Amaranth to Dicamba and 2,4-D. *Agronomy Journal* 108, 1649–1655.
339 doi:10.2134/agronj2015.0465.
- 340 Darwin, C. (1888). *The Different Forms of Flowers on Plants of the Same Species*. J. Murray Available at:
341 <http://books.google.com?id=7uMEAAAAYAAJ>.
- 342 Davis, A. S., Schutte, B. J., Hager, A. G., and Young, B. G. (2015). Palmer Amaranth (*Amaranthus palmeri*)
343 Damage Niche in Illinois Soybean Is Seed Limited. *Weed Science* 63, 658–668. doi:10.1614/WS-D-14-
344 00177.1.
- 345 Farmer, J. A., Webb, E. B., Pierce, R. A., and Bradley, K. W. (2017). Evaluating the potential for weed seed
346 dispersal based on waterfowl consumption and seed viability. *Pest Management Science* 73, 2592–2603.
347 doi:10.1002/ps.4710.
- 348 Fisher, R. A. (1930). The genetical theory of natural selection. *Eugen Rev* 22, 127–130. Available at:
349 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2984947/> [Accessed August 12,
350 2021].
- 351 Fox, J., and Weisberg, S. (2018). *An R Companion to Applied Regression*. SAGE Publications Available at:
352 <http://books.google.com?id=uPNrDwAAQBAJ>.
- 353 Garetson, R., Singh, V., Singh, S., Dotray, P., and Bagavathiannan, M. (2019). Distribution of herbicide-
354 resistant Palmer amaranth (*Amaranthus palmeri*) in row crop production systems in Texas. *Weed
355 Technology* 33, 355–365. doi:10.1017/wet.2019.14.
- 356 Guo, P., and Al-Khatib, K. (2003). Temperature effects on germination and growth of redroot pigweed
357 (*Amaranthus retroflexus*), Palmer amaranth (*A. Palmeri*), and common waterhemp (*A. rudis*). *Weed
358 Science* 51, 869–875. doi:10.1614/P2002-127.
- 359 Hartzler, B., and Anderson, M. (2016). Palmer amaranth: It's here, now what? 10.
- 360 Heap, I. (2021). Internation Herbicide-Resistant Weed Database. Available at: <http://www.weedscience.org/Home.aspx> [Accessed July 26, 2021].
- 362 Hothorn, T., Bretz, F., and Westfall, P. (2008). Simultaneous Inference in General Parametric Models.
363 *Biometrical Journal* 50, 346–363. doi:10.1002/bimj.200810425.
- 364 Jha, P., Norsworthy, J. K., Riley, M. B., and Bridges, W. (2010). Annual Changes in Temperature and Light
365 Requirements for Germination of Palmer Amaranth (*Amaranthus palmeri*) Seeds Retrieved from Soil.
366 *Weed Science* 58, 426–432. doi:10.1614/WS-D-09-00038.1.
- 367 Jhala, A. J., Norsworthy, J. K., Ganie, Z. A., Sosnoskie, L. M., Beckie, H. J., Mallory-Smith, C.
368 A., Liu, J., Wei, W., Wang, J., and Stoltenberg, D. E. (2021). Pollen-mediated gene flow and
369 transfer of resistance alleles from herbicide-resistant broadleaf weeds. *Weed Technology* 35, 173–187.
370 doi:10.1017/wet.2020.101.

- 371 Keeley, P. E., Carter, C. H., and Thullen, R. J. (1987). Influence of Planting Date on Growth of Palmer
372 Amaranth (*Amaranthus palmeri*). *Weed Science* 35, 199–204. doi:10.1017/S0043174500079054.
- 373 Kistner, E. J., and Hatfield, J. L. (2018). Potential Geographic Distribution of Palmer
374 Amaranth under Current and Future Climates. *Agricultural & Environmental Letters* 3, 170044.
375 doi:10.2134/ael2017.12.0044.
- 376 Klingaman, T. E., and Oliver, L. R. (1994). Palmer Amaranth (*Amaranthus palmeri*) Interference in
377 Soybeans (*Glycine max*). *Weed Science* 42, 523–527. doi:10.1017/S0043174500076888.
- 378 Korres, N. E., and Norsworthy, J. K. (2017). Palmer Amaranth (*Amaranthus palmeri*) Demographic and
379 Biological Characteristics in Wide-Row Soybean. *Weed Science* 65, 491–503. doi:10.1017/wsc.2017.12.
- 380 Korres, N. E., Norsworthy, J. K., FitzSimons, T., Roberts, T. L., and Oosterhuis, D. M. (2017). Differential
381 Response of Palmer Amaranth (*Amaranthus palmeri*) Gender to Abiotic Stress. *Weed Science* 65,
382 213–227. doi:10.1017/wsc.2016.34.
- 383 Korres, N. E., Norsworthy, J. K., Mauromoustakos, A., and Williams, M. M. (2020). Soybean density and
384 Palmer amaranth (*Amaranthus palmeri*) establishment time: Effects on weed biology, crop yield, and
385 economic returns. *Weed Science* 68, 467–475. doi:10.1017/wsc.2020.41.
- 386 Kuhn, M., Vaughan, D., and RStudio (2021). *Yardstick: Tidy Characterizations of Model Performance*.
387 Available at: <https://CRAN.R-project.org/package=yardstick> [Accessed August 24,
388 2021].
- 389 Kumar, V., Liu, R., and Stahlman, P. W. (2020). Differential sensitivity of Kansas Palmer amaranth
390 populations to multiple herbicides. *Agronomy Journal* 112, 2152–2163. doi:10.1002/agj2.20178.
- 391 Küpper, A., Borgato, E. A., Patterson, E. L., Netto, A. G., Nicolai, M., Carvalho, S. J. P. de, Nissen, S.
392 J., Gaines, T. A., and Christoffoleti, P. J. (2017). Multiple Resistance to Glyphosate and Acetolactate
393 Synthase Inhibitors in Palmer Amaranth (*Amaranthus palmeri*) Identified in Brazil. *Weed Science* 65,
394 317–326. doi:10.1017/wsc.2017.1.
- 395 Lang, A. (1965). “Physiology of flower initiation,” in *Differenzierung und Entwicklung / Differentiation
396 and Development* Handbuch der Pflanzenphysiologie / Encyclopedia of Plant Physiology., ed. A. Lang
397 (Berlin, Heidelberg: Springer), 1380–1536. doi:10.1007/978-3-642-50088-6_39.
- 398 Larran, A. S., Palmieri, V. E., Perotti, V. E., Lieber, L., Tuesca, D., and Permingeat, H. R. (2017). Target-site
399 resistance to acetolactate synthase (ALS)-inhibiting herbicides in *Amaranthus palmeri* from Argentina.
400 *Pest Management Science* 73, 2578–2584. doi:10.1002/ps.4662.
- 401 Leeper, T. J. (2017). Interpreting Regression Results using Average Marginal Effects with R’s margins. 31.
- 402 Leeper, T. J., Arnold, J., Arel-Bundock, V., and Long, J. A. (2021). *Margins: Marginal Effects
403 for Model Objects*. Available at: <https://CRAN.R-project.org/package=margins>
404 [Accessed August 24, 2021].
- 405 Lenth, R. V., Buerkner, P., Herve, M., Love, J., Riebl, H., and Singmann, H. (2021). *Emmeans: Estimated
406 Marginal Means, aka Least-Squares Means*. Available at: [https://CRAN.R-project.org/
package=emmeans](https://CRAN.R-project.org/
407 package=emmeans) [Accessed August 24, 2021].
- 408 Lindsay, K., Popp, M., Norsworthy, J., Bagavathiannan, M., Powles, S., and Lacoste, M. (2017). PAM:
409 Decision Support for Long-Term Palmer Amaranth (*Amaranthus palmeri*) Control. *Weed Technology*
410 31, 915–927. doi:10.1017/wet.2017.69.

- 411 MacRae, A. W., Webster, T. M., Sosnoskie, L. M., Culpepper, A. S., and Kichler, J. M. (2013). Cotton
412 Yield Loss Potential in Response to Length of Palmer Amaranth (*Amaranthus palmeri*) Interference.
413 17, 6.
- 414 Mangiafico, S. (2021). *Rcompanion: Functions to Support Extension Education Program Evaluation*.
415 Available at: <https://CRAN.R-project.org/package=rcompanion> [Accessed August
416 24, 2021].
- 417 Massinga, R. A., Currie, R. S., Horak, M. J., and Boyer, J. (2001). Interference of Palmer amaranth in corn.
418 *Weed Science* 49, 202–208. doi:10.1614/0043-1745(2001)049[0202:IOPAIC]2.0.CO;2.
- 419 Matzrafi, M., Osipitan, O. A., Ohadi, S., and Mesgaran, M. B. (2021). Under pressure: Maternal effects
420 promote drought tolerance in progeny seed of Palmer amaranth (*Amaranthus palmeri*). *Weed Science*
421 69, 31–38. doi:10.1017/wsc.2020.75.
- 422 McFarlane, G. R., Whitelaw, C. B. A., and Lillico, S. G. (2018). CRISPR-Based Gene Drives for Pest
423 Control. *Trends in Biotechnology* 36, 130–133. doi:10.1016/j.tibtech.2017.10.001.
- 424 Menges, R. M. (1987). Weed Seed Population Dynamics during Six Years of Weed Management Systems
425 in Crop Rotations on Irrigated Soil. *Weed Science* 35, 328–332. Available at: <http://www.jstor.org/stable/4044593>.
- 427 Mesgaran, M. B., Matzrafi, M., and Ohadi, S. (2021). Sex dimorphism in dioecious Palmer amaranth
428 (*Amaranthus palmeri*) in response to water stress. *Planta* 254, 17. doi:10.1007/s00425-021-03664-7.
- 429 Milani, A., Panizzo, S., Farinati, S., Iamonico, D., Sattin, M., Loddo, D., and Scarabel, L. (2021). Recent
430 Discovery of *Amaranthus palmeri* S. Watson in Italy: Characterization of ALS-Resistant Populations
431 and Sensitivity to Alternative Herbicides. *Sustainability* 13, 7003. doi:10.3390/su13137003.
- 432 Montgomery, J. S., Giacomini, D. A., Weigel, D., and Tranel, P. J. (2021). Male-specific Y-chromosomal
433 regions in waterhemp (*Amaranthus tuberculatus*) and Palmer amaranth (*Amaranthus palmeri*). *New
434 Phytologist* 229, 3522–3533. doi:10.1111/nph.17108.
- 435 Montgomery, J. S., Sadeque, A., Giacomini, D. A., Brown, P. J., and Tranel, P. J. (2019). Sex-specific
436 markers for waterhemp (*Amaranthus tuberculatus*) and Palmer amaranth (*Amaranthus palmeri*). *Weed
437 Science* 67, 412–418. doi:10.1017/wsc.2019.27.
- 438 Morgan, G. D., Baumann, P. A., and Chandler, J. M. (2001). Competitive Impact of Palmer Amaranth
439 (*Amaranthus palmeri*) on Cotton (*Gossypium hirsutum*) Development and Yield. *Weed Technology* 15,
440 408–412. doi:10.1614/0890-037X(2001)015[0408:CIOPAA]2.0.CO;2.
- 441 Nguyen, H. M., Kleitou, P., Kletou, D., Sapir, Y., and Winters, G. (2018). Differences in flowering sex
442 ratios between native and invasive populations of the seagrass *Halophila stipulacea*. *Botanica Marina*
443 61, 337–342. doi:10.1515/bot-2018-0015.
- 444 Oliveira, M. C. (2021). *Maxwelco/palmer_adaptation*. Zenodo Available at: <https://doi.org/10.5281/zenodo.5236831>.
- 446 Oliveira, M. C., Gaines, T. A., Patterson, E. L., Jhala, A. J., Irmak, S., Amundsen, K., and Knezevic, S.
447 Z. (2018). Interspecific and intraspecific transference of metabolism-based mesotrione resistance in
448 dioecious weedy *Amaranthus*. *The Plant Journal* 96, 1051–1063. doi:10.1111/tpj.14089.
- 449 Oliveira, M. C., Giacomini, D. A., Arsenijevic, N., Vieira, G., Tranel, P. J., and Werle, R. (2021).
450 Distribution and validation of genotypic and phenotypic glyphosate and PPO-inhibitor resistance in

- 451 Palmer amaranth (*Amaranthus palmeri*) from southwestern Nebraska. *Weed Technology* 35, 65–76.
452 doi:10.1017/wet.2020.74.
- 453 Page, E. R., Nurse, R. E., Meloche, S., Bosveld, K., Grainger, C., Obeid, K., Filotas, M., Simard, M.-J., and
454 Laforest, M. (2021). Import of Palmer amaranth (*Amaranthus palmeri* S. Wats.) Seed with sweet potato
455 (*Ipomea batatas* (L.) Lam) slips. *Can. J. Plant Sci.*, CJPS-2020-0321. doi:10.1139/CJPS-2020-0321.
- 456 Pérez-Llorca, M., and Sánchez Vilas, J. (2019). Sexual dimorphism in response to herbivory and
457 competition in the dioecious herb *Spinacia oleracea*. *Plant Ecol* 220, 57–68. doi:10.1007/s11258-
458 018-0902-7.
- 459 Piskackova, T. A. R., Reberg-Horton, S. C., Richardson, R. J., Jennings, K. M., Franca, L., Young, B. G.,
460 and Leon, R. G. (2021). Windows of action for controlling palmer amaranth (*Amaranthus palmeri*)
461 using emergence and phenology models. *Weed Research* 61, 188–198. doi:10.1111/wre.12470.
- 462 Price, A. J., Balkcom, K. S., Culpepper, S. A., Kelton, J. A., Nichols, R. L., and Schomberg, H. (2011).
463 Glyphosate-resistant Palmer amaranth: A threat to conservation tillage. *Journal of Soil and Water
464 Conservation* 66, 265–275. doi:10.2489/jswc.66.4.265.
- 465 Raoniar, R. (2021). Modelling Binary Logistic Regression Using R (research-oriented modelling and
466 interpretation). Available at: <https://bit.ly/3BhucN3> [Accessed August 11, 2021].
- 467 Ritz, C., Baty, F., Streibig, J. C., and Gerhard, D. (2015). Dose-Response Analysis Using R. *PLOS ONE*
468 10, e0146021. doi:10.1371/journal.pone.0146021.
- 469 Rumpa, M. M., Krausz, R. F., Gibson, D. J., and Gage, K. L. (2019). Effect of PPO-Inhibiting Herbicides
470 on the Growth and Sex Ratio of a Dioecious Weed Species *Amaranthus palmeri* (Palmer Amaranth).
471 *Agronomy* 9, 275. doi:10.3390/agronomy9060275.
- 472 Sauer, J. (1957). Recent Migration and Evolution of the Dioecious Amaranths. *Evolution* 11, 11–31.
473 doi:10.2307/2405808.
- 474 Sauer, J. D. (1972). The dioecious amaranths: A new species name and major range extensions. *Madroño*
475 21, 426–434. Available at: <http://www.jstor.org/stable/41423815>.
- 476 Sbatella, G. M., and Wilson, R. G. (2010). Isoxaflutole Shifts Kochia (*Kochia scoparia*) Populations in
477 Continuous Corn. *Weed Technology* 24, 392–396. doi:10.1614/WT-D-09-00023.1.
- 478 Schliekelman, P., Ellner, S., and Gould, F. (2005). Pest Control by Genetic Manipulation of Sex Ratio.
479 *Journal of Economic Entomology* 98, 18–34. doi:10.1093/jee/98.1.18.
- 480 Schwartz, L. M., Norsworthy, J. K., Young, B. G., Bradley, K. W., Kruger, G. R., Davis, V. M., Steckel,
481 L. E., and Walsh, M. J. (2016). Tall Waterhemp (*Amaranthus tuberculatus*) and Palmer amaranth
482 (*Amaranthus palmeri*) Seed Production and Retention at Soybean Maturity. *Weed Technology* 30,
483 284–290. doi:10.1614/WT-D-15-00130.1.
- 484 Sing, T., Sander, O., Beerenwinkel, N., and Lengauer, T. (2005). ROCR: Visualizing classifier performance
485 in R. *Bioinformatics* 21, 3940–3941. doi:10.1093/bioinformatics/bti623.
- 486 Sosnoskie, L. M., Webster, T. M., and Culpepper, A. S. (2013). Glyphosate Resistance Does Not
487 Affect Palmer Amaranth (*Amaranthus palmeri*) Seedbank Longevity. *Weed Science* 61, 283–288.
488 doi:10.1614/WS-D-12-00111.1.

- 489 Spaunhorst, D. J., Devkota, P., Johnson, W. G., Smeda, R. J., Meyer, C. J., and Norsworthy, J. K. (2018).
490 Phenology of Five Palmer amaranth (*Amaranthus palmeri*) Populations Grown in Northern Indiana and
491 Arkansas. *Weed Science* 66, 457–469. doi:10.1017/wsc.2018.12.
- 492 Team, R. C. (2021). *R: The R Project for Statistical Computing*. Vienna, Austria: R Foundation for Statistical
493 Computing Available at: <https://www.r-project.org/> [Accessed August 24, 2021].
- 494 Van Wychen, L. (2020). 2020 Survey of the most common and troublesome weeds in grass crops, pasture,
495 and turf in the United States and Canada. Available at: <https://wssa.net/wp-content/>
496 uploads/2020-Weed-Survey_grass-crops.xlsx.
- 497 Wang, J. L., Klessig, D. F., and Berry, J. O. (1992). Regulation of C4 Gene Expression in Developing
498 Amaranth Leaves. *The Plant Cell* 4, 173–184. doi:10.1105/tpc.4.2.173.
- 499 Ward, S. M., Webster, T. M., and Steckel, L. E. (2013). Palmer Amaranth (*Amaranthus palmeri*): A Review.
500 *Weed Technology* 27, 12–27. doi:10.1614/WT-D-12-00113.1.
- 501 Webster, T. M., and Grey, T. L. (2015). Glyphosate-Resistant Palmer Amaranth (*Amaranthus palmeri*)
502 Morphology, Growth, and Seed Production in Georgia. *Weed Science* 63, 264–272. doi:10.1614/WS-D-
503 14-00051.1.
- 504 Wu, C., and Owen, M. D. K. (2014). When Is the Best Time to Emerge: Reproductive Phenology and
505 Success of Natural Common Waterhemp (*Amaranthus rudis*) Cohorts in the Midwest United States?
506 *Weed Science* 62, 107–117. Available at: <http://www.jstor.org/stable/43700638>.
- 507 Yacoubi, R., and Axman, D. (2020). Probabilistic Extension of Precision, Recall, and F1 Score for
508 More Thorough Evaluation of Classification Models. in *Proceedings of the First Workshop on*
509 *Evaluation and Comparison of NLP Systems* (Online: Association for Computational Linguistics),
510 79–91. doi:10.18653/v1/2020.eval4nlp-1.9.
- 511 Yu, E., Blair, S., Hardel, M., Chandler, M., Thiede, D., Cortilet, A., Gunsolus, J., and Becker, R. (2021).
512 Timeline of Palmer amaranth (*Amaranthus palmeri*) invasion and eradication in Minnesota. *Weed*
513 *Technology*, 1–31. doi:10.1017/wet.2021.32.




Figure 1. Mean average temperature (C) and total monthly precipitation (mm) at Arlington, WI, Clay Center, NE, Grant, NE and Macomb, IL




Figure 2. Palmer amaranth adaptation study layout of a plant cohort timing in bareground, corn, and soybean. Twenty-four Palmer amaranth plants were placed 76.2 cm apart in each field experimental unit

FIGURES




Figure 3. Palmer amaranth biomass (A) and height (B) growing in corn, bareground, and soybean nested across Arlington, WI, Clay Center, NE, Grant, NE, Lincoln, NE and Macomb, IL




Figure 4. Cumulative flowering of Palmer amaranth at first and second transplant timing (A) and day of year of 10, 50, and 90 cumulative flowering at first and second cohort transplanting time (B) nested across Arlington, WI, Clay Center, NE, Grant, NE, Lincoln, NE and Macomb, IL




Figure 5. The probability (P) of being female Palmer amaranth by day of year (A), weight (B), and height (C). Black line represents the model estimation and shaded green the confidence intervals




Figure 6. MCO (180 cm) holds harvested Palmer amaranth plants at 40 days after first cohort transplanting (A) and 33 days after second cohort transplanting (B) time. From left to right in each image, Palmer amaranth growing in bareground, soybean and corn in Arlington, Wisconsin