

Palmer amaranth (*Amaranthus palmeri*) adaptation to US Midwest agroecosystems

Maxwel C Oliveira¹, Amit J Jhala², Mark Bernards³, Chris Proctor², Strahinja Stepanovic², Rodrigo Werle^{1*}

¹ Department of Agronomy, University of Wisconsin-Madison, Madison, Wisconsin, United States

² Department of Agronomy and Horticulture, University of Nebraska-Lincoln, Lincoln, Nebraska, United States

³ Department of Agronomy, Western Illinois University, Macomb, Illinois, United States

Correspondence*:

Rodrigo Werle

rwerle@uwisc.edu

2 ABSTRACT

Palmer amaranth (*Amaranthus palmeri* S. Watson) is one of the most troublesome agronomic weed species in the United States. Palmer amaranth is endemic to Southern United States but its range is expanding northward. Palmer amaranth dispersal warrants studies assessing species adaptation into new geographies. A study was conducted to investigate morphology, flowering and gender from cohorts of Palmer amaranth growing under corn, soybean, and bareground across five locations of US Midwest. Results demonstrated that the first cohort of Palmer amaranth, established in June, produced 42% more biomass than plants from the second cohort (established in July). The first Palmer amaranth cohort produced 75.5 g plant⁻¹ in bareground, 28.3 g plant⁻¹ in soybean and 16.3 g plant⁻¹ in corn, whereas the second Palmer amaranth cohort produced 62.6, 6.3, and 1.4 g plant⁻¹ in bareground, soybean and corn, respectively. Palmer amaranth height was more impacted when growing in corn. Palmer amaranth plants averaged 85.2 cm tall in the first cohort but 38.2 cm tall in the second cohort in corn. Moreover, Palmer amaranth flowering window shifted according to crop and cohort timings. Palmer amaranth growing in intense competition, such as under low light in corn, resulted in the longest flowering window. Also, Palmer amaranth gender was slightly influenced by day of year, weight and height. The model estimated that probability of being a female plant increased as biomass and height increased. Our results showed the fast adaptation and plasticity of Palmer amaranth to grow and adapt to cropping systems from the US Midwest. Palmer amaranth is likely to continue its expansion northward. Therefore, preventing plant dispersal into new habitats is the most effective management strategy. Reactive management to reduce Palmer amaranth impact on cropping systems should encompass diversity of tactics that minimize the species ability to establish into cropping systems, including crop rotation (beyond corn and soybean), early/late crop planting, row spacing, cover crops, and effective chemical control programs.

Keywords: Evolution, Flowering, Management, Pigweed, Weed

INTRODUCTION

27 Palmer amaranth (*Amaranthus palmeri* S. Watson) is currently ranked as one of the most economically
28 detrimental weed species to cropping systems in the United States (Van Wychen, 2020). Unmanaged
29 Palmer amaranth plants compete for water, light, and nutrients, which can drastically impact crop yields
30 (Berger et al., 2015). For example, Palmer amaranth has been documented to reduce up to 91%, 68%, and
31 54% corn (Massinga et al., 2001), soybean (Klingaman and Oliver, 1994), and cotton (Morgan et al., 2001)
32 yields, respectively. Moreover, Palmer amaranth has shown a remarkable capacity to evolve resistance
33 to herbicides. To date, Palmer amaranth has evolved resistance to eight herbicide sites of action (Heap,
34 2021), increasing the weed management complexity (Lindsay et al., 2017). Thus, Palmer amaranth poses
35 an economical and ecological risk to sustainable agriculture.

36 Palmer amaranth is a fast growing summer annual forb indigenous to the Sonoran Desert (Sauer, 1957).
37 The species would eventually emerge as a threat to US agriculture in the 1990s. Palmer amaranth weediness
38 is likely a result of human-assisted selection in combination with plant biology. Farm mechanization,
39 adoption of conservation agriculture (e.g., no-till), and intensive use of herbicides for weed management
40 are the main human-mediated selections of Palmer amaranth into cropping systems (Ward et al., 2013).
41 On the other hand, Palmer amaranth is a prolific seed producer with a C4 photosynthetic apparatus (Wang
42 et al., 1992). With a dioecious nature, Palmer amaranth male and female plants are obligate outcrosser
43 species, increasing the chances of exchanging adaptive traits among plants (Jhala et al., 2021; Oliveira et
44 al., 2018). Also, Palmer amaranth's small seeds (e.g, 1 mm) tend to thrive in no-tillage systems (Price et
45 al., 2011), and spread across locations through farm equipment (Sauer, 1972), seed mixes (Hartzler and
46 Anderson, 2016), wildlife (Farmer et al., 2017), etc. The dispersal capacity of Palmer amaranth makes the
47 species one of the most successful cases of weed adaptation to current cropping systems.

48 Light and temperature are the main environmental requirements for Palmer amaranth growth and
49 development (Jha et al., 2010). Palmer amaranth is reported with an extended germination period (Ward
50 et al., 2013). Germination of Palmer amaranth was triggered by 18 C soil temperature at 5 cm depth
51 (Keeley et al., 1987), and optimal germination and biomass production occurred at 35/30 C day and night
52 temperatures (Guo and Al-Khatib, 2003). In addition, Palmer amaranth emergence is human-mediated
53 by tillage timings and preemergence-applied herbicides (Chahal et al., 2021), which can result in weed
54 germination shifts (Sbatella and Wilson, 2010). Also, Palmer amaranth has shown to grow and develop
55 under reduced soil water content. In continuous water stress, Palmer amaranth survived and produced
56 at least 14000 seeds plant⁻¹ (Chahal et al., 2018). Seeds from Palmer amaranth growing with limited
57 water conditions were heavier, less dormant, and prompt for germination (Matzrafi et al., 2021). Growing
58 conditions and management practices also influence Palmer amaranth sex dimorphism and flowering
59 pattern (Korres et al., 2017; Rumpa et al., 2019). Therefore, Palmer amaranth has shown plasticity to
60 evolve and fast adapt under the current agroecosystem conditions. Future scenarios suggest that global
61 temperature warming can impact agriculture, and promote niches for Palmer amaranth invasion/adaptation
62 into new environments. Agronomic crops in Australia and Sub-Saharan Africa are currently at greatest risk
63 to Palmer invasion as climate warms (Kistner and Hatfield, 2018). Temperature is a key factor limiting
64 Palmer amaranth expansion to cooler geographies (Briscoe Runquist et al., 2019); however, under future
65 climate change Palmer amaranth is likely to expand northward into Canada and Northern Europe (Kistner
66 and Hatfield, 2018; Briscoe Runquist et al., 2019).

67 Palmer amaranth is already found in agronomic crops of South America (Larran et al., 2017; Küpper et al.,
68 2017) and Southern Europe (Milani et al., 2021). In the United States, Palmer amaranth is well established
69 in the Cotton Belt (Garetson et al., 2019; Bagavathiannan and Norsworthy, 2016) in the southern United

70 States but its range is expanding northward. For example, herbicide resistant Palmer amaranth is widespread
71 in Nebraska (Oliveira et al., 2021). There are some reported cases of Palmer amaranth in Michigan (Kohrt
72 et al., 2017) and Connecticut (Aulakh et al., 2021). Also, it is estimated that Palmer amaranth can cause
73 yield loss to soybean fields in Illinois (Davis et al., 2015), which is concerning as soybean along with corn
74 comprise most of US Midwest agronomic hectares. In Iowa, a study showed that Palmer amaranth is still
75 not well adapted compared to waterhemp (*Amaranthus tuberculatus*) (Baker, 2021), which is considered
76 the most troublesome species in the US Midwest (Tranel et al., 2011). Invasion and successful eradication
77 of Palmer amaranth is documented in Minnesota (Yu et al., 2021). Palmer amaranth infestations have not
78 been detected in Canada; however, Palmer amaranth seeds were detected in sweet potato slips in the country
79 (Page et al., 2021). Palmer amaranth is still not as well adapted and established to Northern as compared to
80 Southern North America. Therefore, Palmer amaranth range of expansion into new habitats can increase.
81 It seems certain the need to manage new Palmer amaranth infestations in agronomic crops throughout
82 northern United States is eminent in the near future. Strategies on Palmer amaranth management should
83 encompass the agroecosystem level but not focus only on attempts to eradicate the weed. Most tactics to
84 manage Palmer amaranth are based on technological solutions (Scott, 2011), which are short-term (e.g.,
85 herbicide and/or tillage) rather than long-term weed management. Palmer amaranth management should be
86 built around minimizing the species ability to adapt, grow and develop successfully in agroecosystems.

87 In the southeastern US, early growing Palmer amaranth is well known to have a higher impact on cotton
88 yields compared to late established plants (MacRae et al., 2013). In the northern states, Palmer amaranth
89 impact on the agroecosystem is recent. Studies investigating Palmer amaranth in northern states are limited
90 due to the plant classification as noxious weed species (Yu et al., 2021). Nonetheless, the continuous
91 Palmer amaranth dispersal and potential establishment across the northern United States is concerning and
92 warrants investigations on species morphology in northern environments. Understanding Palmer amaranth
93 biology and morphology under different agroecosystems can enhance our knowledge on species adaptation
94 and management practices. It can also aid in designing proactive and ecological tactics to limit the species
95 range expansion, reduce its negative impact on agroecosystems, and to develop resilient and sustainable
96 farming systems (MacLaren et al., 2020). Therefore, the objective of this study was to investigate the
97 flowering pattern, gender, biomass production, and height of Palmer amaranth cohorts growing under corn,
98 soybean and bareground across five locations in the United States Midwest.

MATERIAL AND METHODS

99 Plant material and growing conditions

100 A Palmer amaranth accession (Kei3) from Perkins County, Nebraska with no glyphosate resistance
101 according to Oliveira et al. (2021) was selected for this study. Three weeks prior to the establishment of
102 each cohort, seeds were planted in plastic trays containing potting-mix. Emerged seedlings (1 cm) were
103 transplanted into 200 cm⁻³ plastic pots (a plant pot⁻¹). Palmer amaranth seedlings were supplied with
104 adequate water and kept under greenhouse conditions at the University of Wisconsin-Madison, University
105 of Nebraska-Lincoln, and Western Illinois University; and kept outdoors at the Perkins extension office in
106 Grant, NE until the 2-3 leaf stage (5 to 8 cm height) when they were transported to the field.
107 **Field study**

108 The experiment was conducted in 2018 and 2019 under field conditions at five locations: Arlington, WI
109 (43°18'N, 89°29'W), Clay Center, NE (40.57'N, 9814'W), Grant, NE (40.85'N, -101.70'W), Lincoln, NE
110 (41.16'N, 96.42'W), and Macomb, IL (XXX'N, XXX'W).

111 Fields were conventionally tilled prior to crop planting. Corn and soybean were planted in 76-cm row
 112 spacing (Table 1). Monthly mean air temperature and total precipitation were obtained using Daymet
 113 weather data from June through September across the five locations in 2018 and 2019 (Correndo et al.,
 114 2021) (Figure 1)

Table 1. Field study attributes from Arlington, WI, Clay Center, NE, Grant, NE, Lincoln, NE and Macomb, IL

Attributes	Arlington, WI	Clay Center, NE	Grant, NE	Lincoln, NE	Macomb, IL
Bareground					
Corn	Weed control ^a	S-metolachor ^b fb glyphosate ^c / saflufenacil + imazethapyr + pyroxasulfone ^d	glyphosate	glyphosate + S-metolachor ^f fb glyphosate ^e	
	Hybrid	NK0142 3120-EZ1	DKC60-67	G03C84-5122-EZ1	P1197AM
	Seeding rate	88956	86487	79000	86485
	Weed control	S-metolachor fb glyphosate	S-metolachor + trazine + mesotrione + bicyclopyrone ^e	glyphosate	glyphosate + S-metolachor ^f fb glyphosate ^e
	Stage at 1 st cohort	V2-3	V6-7	V6-7	V2-3
	Stage at 2 nd cohort	V6-7	V6-7	V6-7	V6-7
	Planting day	April 30, 2018 / May 5, 2019	May 10, 2018/19	May 5, 2018 / May 14, 2019	April 27, 2018 / May 3, 2019
	Fertilization	N (46-0-0) at 157 kg ha ⁻¹			N (46-0-0) at 336 kg ha ⁻¹
Soybean	Variety	DSR-1950	AG21X8	AG28X7	2018: 3017R2X; 2019 P31A22X
	Seeding rate	296400	321237	220000	370650
	Weed control	glyphosate / S-metolachor	saflufenacil + imazethapyr + pyroxasulfone	glyphosate	glyphosate + S-metolachor fb glyphosate
	Stage at 1 st cohort	V1-2	V1-2	V1-2	V1-2
	Stage at 2 nd cohort	V5-6	V5-6	V5-6	V5-6
	Planting day	May 5, 2018 / May 10, 2009	May 14, 2018/19	May 5, 2018 / May 14, 2019	May 7, 2018 / May 17, 2019
Soil	Type	Plano-silt-loam	Crete Silt Loam	2018: Mace silt loam / 2019: Kuma silt loam	Crete silty clay loam
	pH	6.6	6.5	2018: 8.1 / 2019: 5.1	5.2
	Organic matter (%)	3.5	3	2018: 2 / 2019: 1.7	3.4

^aHoe weeding; ^bS-metolachor, 1324 g ai ha⁻¹; ^cglyphosate, 1262 g ae ha⁻¹; ^dsaflufenacil + imazethapyr + pyroxasulfone, 215 g ai ha⁻¹; ^eS-metolachor + trazine + mesotrione, + bicyclopyrone, 2409 g ai ha⁻¹; ^fS-metolachor, 1070 g ai ha⁻¹; ^gglyphosate, 870 g ae ha⁻¹

115 The field experimental units were three adjacent 9.1 m wide (12 rows at 76.2 cm row spacing) by 10.7
 116 m long. The experimental design were arranged in factorial design with three crops, two transplanting
 117 times simulating two cohorts, repeated across five locations. Each field experimental unit was planted with
 118 corn, soybean, or kept under bareground. The two transplant timings were June 1 (first cohort) and July 1
 119 (second cohort). Palmer amaranth seedlings (potting mix + two seedlings) were transplanted (6 cm deep
 120 and 8 cm wide). Forty-eight plants were equidistantly placed (0.76 m apart) between rows within each
 121 crop. After a week, one plant was eliminated and one was kept, resulting in 24 plants per experimental unit
 122 and transplanting time (Figure 2). When needed, Palmer amaranth plants were supplied with water during
 123 the first week after transplanting to assure seedling survival.

124 After transplanting, Palmer amaranth flowering was monitored until the end of the study. When a plant
 125 flowered, the day was recorded, plant gender was identified (male or female), plant height was measured
 126 from soil surface to the top of plant. Also, aboveground plant biomass was harvested near soil surface and
 127 oven dried at 65 C until reaching constant weight before weigh (g plant⁻¹) was recorded.

128 Plants had to be harvested at flowering because Palmer amaranth is neither endemic in Wisconsin nor in
 129 Illinois. In our study, all locations followed the methodology of plant harvest at flowering initiation, except
 130 in Grant, NE. In this location, all Palmer amaranth plants were harvest at once on July 06, 2018 and 2019
 131 (first cohort), and on August 17, 2018 and on July 31, 2019 (second cohort).

132 Statistical analyses

133 The statistical analyses were performed using R statistical software version 4.0.1 (Team, 2021).

134 Analyses of Palmer amaranth height and biomass were performed with a linear mixed model using *lmer*
 135 function from “lme4” package (Bates et al., 2015). Plant height and biomass were log transformed to meet
 136 model assumption of normality. In the model, crop (bareground, corn, soybean) and cohort time (first and
 137 second) were the fixed effects and year nested with location the random effects. Analysis of variance at α
 138 0.05 was performed with *anova* function from “car” package (Fox and Weisberg, 2018). Marginal means
 139 and compact letter display were estimated with *emmeans* and *cld* from packages “emmeans” (Lenth et al.,
 140 2021) and “multcomp” (Hothorn et al., 2008), respectively.

141 The Palmer amaranth flowering timing was estimated as cumulative flowering across all locations, except
142 Grant, NE. Palmer amaranth cumulative flowering estimation was determined using an asymmetrical three
143 parameter log logistic Weibull model of the drc package (Ritz et al., 2015).

$$Y(x) = 0 + (d - 0)\exp(-\exp(b(\log(x) - e)))$$

144 In this model, Y is the Palmer amaranth cumulative flowering, d is the upper limit (set to 100), and e is
145 the inflection point, and x day of year (doy).

146 The doy for 10, 50, and 90% Palmer amaranth cumulative flowering were determined using the *ED*
147 function of drc package. Also, the 10, 50, and 90% Palmer amaranth cumulative flowering were compared
148 among crops and cohorts using the *EDcomp* function of drc package. The EDcomp function compares the
149 ratio of cumulative flowering using t-statistics, where P-value < 0.05 indicates that we fail to reject the null
150 hypothesis.

151 A binary logistic regression was fitted to Palmer amaranth gender. Binary logistic regression is used
152 for predicting binary classes (Bangdiwala, 2018), such as the probability of a plant being female in a
153 dioecious species. Prior to the analysis, missing values were removed from the dataset. Also, data from
154 Grant was not used in this analysis due to the uniform plant harvesting at that location. The complete
155 dataset was splitted into 80% train and 20% test data. The 80% train is used for the model training and
156 the 20% test is used for checking model performance on unseen dataset. With 80% dataset, a generalized
157 linear model (base R *glm* function) was fitted to binary response variable, the probability of being female
158 (0 to male and 1 to female). The independent variables were day of year harvest, height, weight, and crop
159 (without interaction). The model family was binomial with a logit function. The model fit was assessed
160 through pseudo R-squared values (McFadden, Cox and Snell, Cragg and Uhler) and likelihood ratio using
161 *nagelkerke* function from “rcompanion” package (Mangiafico, 2021). The marginal effects computation
162 was performed with Average Marginal Effects (AMEs) at every observed value of x and average across the
163 results (Leeper, 2017) using *margins* function from “margins” package (Leeper et al., 2021). The 20% test
164 data was predicted using the *predict* function with a cutoff estimation for male or female using *performance*
165 function from ROCR package (Sing et al., 2005). The model quality prediction from the classification
166 algorithm was measured with precision (*precision* function), recall (*recall* function) and F1-score (*f_meas*
167 function) using the “yardstick” package (Kuhn et al., 2021). The precision determines the accuracy of
168 positive predictions (female plants), recall determines the fraction of positives that were correctly identified,
169 and F1-score is a weighted harmonic mean of precision and recall with the best score of 1 and the worst
170 score of 0 (Raoniar, 2021). F1-score conveys the balance between the precision and the recall (Yacoubi
171 and Axman, 2020). The area under the receiver operating curve (AUC-ROC) was also estimated with
172 *performance* function using the true positive and false positive rates. Higher the AUC, better the model is
173 at distinguishing between female and male Palmer amaranth.

RESULTS

174 Palmer amaranth height and biomass

175 Palmer amaranth plants accumulated more biomass when growing in bareground compared to plants
176 growing in soybean and corn (figure 3A). Palmer amaranth plants in the first cohort produced 75.5, 28.3,
177 and 16.3 g plant⁻¹ in bareground, soybean and corn, respectively. Plants from the second cohort produced
178 62.6 g plant⁻¹ in bareground, followed by 6.3 g plant⁻¹ in soybean, and 1.4 g plant⁻¹.

179 Palmer amaranth height was more uniform across cohort timings, except when growing in corn (figure
 180 3B). Palmer amaranth plants from the first cohort were on average 69.2 cm tall in bareground, which
 181 was not different from the 70.7 cm tall plants from the second cohort timing ($P = 0.74$). In addition, no
 182 difference in Palmer amaranth height (69.3 cm) was detected from first cohort plants in soybean to first and
 183 second cohort plants in bareground ($P > 0.75$). Palmer amaranth plants from the second cohort were nearly
 184 10 cm lower compared to the first cohort in soybeans ($P = 0.04$). The tallest (first cohort) and smallest
 185 (second cohort) Palmer amaranth plants were found in corn. Palmer amaranth reached 85.2 and 38.2 cm
 186 tall, respectively.

187 **Palmer amaranth cumulative flowering**

188 Palmer amaranth plants from the first cohort growing in corn resulted in a longer flowering window
 189 compared to plants growing in bareground and soybean (Figure 4A). The 10% cumulative Palmer amaranth
 190 flowering in soybean, bareground and corn occurred at the end of June. Palmer amaranth reached 10%
 191 flowering in soybean, bareground and corn at doy 180, 180.9 and 181.7, respectively. The 50% Palmer
 192 amaranth cumulative flowering occurred in July. Palmer amaranth reached 50% flowering in bareground,
 193 soybean and corn at doy 193.4, 194.8, and 206.6, respectively. Similar trend was observed at 90%
 194 Palmer amaranth cumulative flowering. Palmer amaranth reached 90% flowering at doy 252.6 in corn
 195 (early September), which was 38 and 32 days after reaching 90% flowering in bareground and soybean,
 196 respectively.

197 Palmer amaranth cumulative flowering in the second cohort ranged from mid July to mid September
 198 (Figure 4B). Palmer amaranth growing in the bareground resulted in earlier flowering time compared to
 199 soybean and corn. Palmer amaranth growing in bareground reached 10%, 50%, and 90% flowering time at
 200 day 203.8, 214.4, and 232.2, respectively. Palmer amaranth growing in soybean reached 10% flowering at
 201 doy 210.9, which was 6 days prior to corn (P -value = 0.00). Similar trend was observed at 50% flowering,
 202 whereas Palmer amaranth reached 50% flowering in corn (doy 233.0) 4 days after soybeans (doy 228.9; P
 203 = 0.00). The 90% Palmer amaranth cumulative flowering occurred at same day in corn (260.9) and soybean
 204 (260.5; P = 0.66).

205 **Palmer amaranth gender**

206 The model goodness of fit was 0.23, 0.32, 0.40 using pseudo R-squared test from McFadden, Cox and
 207 Snell, and Cragg and Uhler, respectively. The likelihood ratio test showed a p-value of < 0.00. The average
 208 marginal effects showed that Palmer amaranth growing in corn resulted in 14.8% less females plants (Table
 209 2). Moreover, increasing a unit doy increases the probability of having a female plant by 0.4% (Table 2 and
 210 Figure 4A). Similar trend is observed for weight (Figure 4B) as well as height (Figure 4C), whereas the
 211 probability of being female increase by 0.2% and 0.1% when a unit of weight (g) and height (cm) increases,
 212 respectively.

Table 2. Average marginal effects of Palmer amaranth gender logistic model. Factor parameter values (crop and bareground) is shown compared to soybean.

Term	AME	SE	Lower	Upper	Z-score	P-value
crop_bareground	-0.048	0.054	-0.154	0.059	-0.876	0.381
crop_corn	-0.148	0.052	-0.250	-0.046	-2.842	0.004
doyh	0.004	0.001	0.003	0.006	4.959	0.000
height	0.002	0.001	0.001	0.003	2.953	0.003
weight	0.001	0.000	0.000	0.001	2.179	0.029

^a Average Marginal Effects. ^b Standard Error.

213 The model accuracy evaluation accuracy in the 20% test dataset was 0.62 with a cutoff value for female
214 and male plants of 0.43. The model classification showed a precision of 0.64, recall of 0.66, and a F1-score
215 of 0.65. In addition, the AUC was 0.64.

DISCUSSION

216 Our study showed that Palmer amaranth biomass, height, flowering window and gender varied within
217 crops and cohort timings. In general, first cohort of Palmer amaranth plants were heavier and taller when
218 compared to the second cohort. At first cohort, resources (e.g., soil nutrients) and conditions (e.g., light)
219 were more timely available for the species. High biomass and taller Palmer amaranth plants are likely
220 a weed strategy to compete for light in between crop rows in absence of canopy. In such conditions,
221 Palmer amaranth showed an extraordinary plasticity to adapt upon the agroecosystem. This is evident when
222 comparing Palmer amaranth canopy shape. The Palmer amaranth competition (e.g., light) strategy was
223 to mimic the crop growth and development (Figure 6). These results suggests that Palmer amaranth can
224 quickly evolve life-history traits to adapt to cropping systems and cultural practices, which was also shown
225 in a study varying nitrogen fertilization (Bravo et al., 2018). Our results highlight Palmer amaranth as a
226 threat to field crops as breeding more competitive crop varieties is likely to select more competitive weed
227 biotypes (Bravo et al., 2017).

228 Palmer amaranth growth and development in second cohort was limited due to the crop competitive
229 ability at advanced development stages. Plants were transplanted at greater crop height and width, which
230 reduced Palmer amaranth competitiveness. As a result, Palmer amaranth height and biomass was lower
231 compared to its first cohort. Moreover, Palmer amaranth growing without crop competition produced
232 the highest amounts of biomass. The Palmer amaranth strategy in bareground was to invest biomass in
233 growing plant width and height. Nonetheless, Palmer amaranth produced 17% less biomass in second
234 cohort compared to first cohort timing. In a bareground study, early emerged Palmer amaranth without
235 competition was 50% taller than late emerged plants (Webster and Grey, 2015). These results suggest that
236 crop competition is not the only factor limiting late Palmer amaranth establishment. The limited growth of
237 Palmer amaranth at second cohort is likely a reduced plant response to day length, light availability and
238 thermal units (e.g, growing degree days). The *Amaranthus* species are sensitive to photoperiod (Wu and
239 Owen, 2014). It is hypothesize that reduced day length or red/far-red effect contributed to smaller plants
240 at second cohort regardless the crop. A study in North Carolina and Illinois predicted that less than 10%
241 Palmer amaranth seedlings emergence occurred after June (Piskackova et al., 2021). In addition, Palmer
242 amaranth negative impact on soybean (Korres et al., 2020) and cotton (Webster and Grey, 2015) yields was
243 higher when plants were established near to crop planting. Therefore, early management is a key strategy
244 to minimize the damaging impact of Palmer amaranth to US Midwest cropping systems.

245 Seed production was not evaluated due to plant harvest at initiation of flowering. Nonetheless, it is well
246 documented a strong positive correlation between Palmer amaranth biomass and seed production (Schwartz
247 et al., 2016; Spaunhorst et al., 2018). In our study, plants growing from the first cohort accumulated 42%
248 more biomass when compared to the second cohort. Therefore, Palmer amaranth plants growing in the
249 second cohort is likely to produce less seeds regardless the crop. Our observation is consistent with the
250 findings that the first Palmer amaranth cohort produced 50% more seeds per plant than Palmer amaranth
251 plants established six weeks later in bareground (Webster and Grey, 2015). Still, seed production at the
252 second cohort will replenish the soil seedbank. Seed production and deposition in the seedbank is also a
253 key factor for species perpetuation (Menges, 1987). Palmer amaranth can produce a hundred thousands
254 seeds per plant (Schwartz et al., 2016; Keeley et al., 1987), which can stay viable in the soil seedbank
255 for at least 36 months (Sosnoskie et al., 2013). Therefore, preventing Palmer amaranth seed production

256 or/and seed migration to non-native habitats is an essential strategy to minimize the species impact in
257 agroecosystems (Davis et al., 2015).

258 An ecological approach to reduce Palmer amaranth seed production is understanding plant biology,
259 including flowering pattern. Our study suggests that Palmer amaranth flowering was slightly influenced
260 by crops and cohort timings. Palmer amaranth growing in bareground and corn resulted in the overall
261 shortest and longest flowering window, respectively. The shortest flowering window of second cohorts
262 of Palmer amaranth growing in bareground highlighted the impact of photoperiod on flowering. When
263 growing in soybean, Palmer amaranth flowering window was similar to bareground for the first cohort but
264 similar to corn at second cohort timing. Plant flowering initiation is complex and depends on the complex
265 interaction between genetic makeup and the environmental conditions (Lang, 1965). We hypothesize that
266 when growing in high competition (e.g., second cohort), Palmer amaranth plants tend to initiate flowering
267 early, as well as having an extended flowering window. A study has shown that Palmer amaranth initiated
268 flowering two weeks prior to the native waterhemp in Iowa (Baker, 2021). Early flower initiation could also
269 be a plant strategy when growing in stressful conditions. For example, when growing under water stress,
270 early flowering in Palmer amaranth resulted in a mismatch between female and male plants by seven days
271 (Mesgaran et al., 2021). A mismatch in Palmer amaranth male and female flowering period can minimize
272 plant outcrossing, reducing plant seed production and/or exchange of resistant alleles (Jhala et al., 2021).
273 Sex dimorphism manipulation is considered a potential ecological pest control strategy (McFarlane et al.,
274 2018; Schliekelman et al., 2005).

275 The mechanisms of gender-determination in plant species is intriguing and arouse the curiosity of many
276 scientists, including Charles Darwin (Darwin, 1888). In our study, the gender model performance was
277 decent (AUC 0.64) considering the biology of plant flowering. A 1:1 male and female sex ratio is a general
278 evolutionary stable strategy for plant species perpetuation (Fisher, 1930). However, a slight deviation from
279 1:1 sex ratio might occur in some dioecious species. For example, the dioecious *Halophila stipulacea* is
280 a female-biased plant in its native habitat, but the naturalized *H. stipulacea* have a 1:1 ratio (Nguyen et
281 al., 2018). Naturalized of *H. stipulacea* reduced female-male ratio to expand into its non-native habitat
282 (Nguyen et al., 2018). Also, biotic and/or abiotic stress can influence plant gender determination. Palmer
283 amaranth male-to-female ratio was greater under high plant densities (Korres and Norsworthy, 2017) and
284 after herbicide application (Rumpa et al., 2019). Our model estimated that late flowering, heavier and
285 taller Palmer amaranth plants slightly deviated from 1:1 ratio in favor to female plants. It was reported that
286 female Palmer amaranth plants invested more in height, stem and biomass while male invested more in leaf
287 area and leaf dry weight under nutrient deficiency (Korres et al., 2017). Our model also estimated more
288 female plants in soybean and bareground compared to corn, which might be linked to plant competition
289 strategy in each crop. Our results showed the influence of life-history and ecological traits on sexual
290 dimorphism in Palmer amaranth. Sexual dimorphism is documented in other dioecious species (Barrett and
291 Hough, 2013). For example, stronger female plant competition and greater male tolerance to herbivory was
292 reported in *Spinacia oleracea* (Pérez-Llorca and Sánchez Vilas, 2019). Research on candidate genes for
293 sex determination in *Amaranthus* species are currently underway but it is far from complete (Montgomery
294 et al., 2021, 2019). Further studies are also needed to understand the ecological basis of Palmer amaranth
295 flowering, including the plant behavior under climate change.

296 Our study demonstrated the Palmer amaranth plasticity to grow and develop into arable land of US
297 Midwest. It is likely that Palmer amaranth range will continue to expanding into new geographies. The
298 migration of Palmer amaranth into the US Midwest will reshape the landscape as waterhemp and Palmer
299 amaranth will share the same habitat. The presence of Palmer amaranth and waterhemp will increase

300 the weed management complexity. Therefore, preventive management is a priority to minimizing Palmer
301 amaranth dispersal. Reactive management should focus on early-season management programs, which
302 would have a large negative effect on Palmer amaranth growth and development. Long-term tactics that
303 promote early-season crop advantage against Palmer amaranth, including diversity of crops in rotation,
304 early/late crop planting, plant width, and crop residue (e.g. cover crops) would minimize the negative
305 impact of Palmer amaranth to cropping systems. The aggressiveness and differential Palmer amaranth
306 adaptation to agroecosystem is striking and require regional collaboration to minimize the species impact
307 on sustainability and profitability of cropping systems.

DISCLOSURE/CONFLICT-OF-INTEREST STATEMENT

308 The authors declare that the research was conducted in the absence of any commercial or financial
309 relationships that could be construed as a potential conflict of interest.

DATA ACCESSIBILITY

310 The data and scripts used to analyse the data presented in this work can be found at Zenodo (Oliveira,
311 2021).

AUTHOR CONTRIBUTIONS

312 RW and MO: designed the experiments; AJ, CP, MB, MO, and SS: conducted the experiments; MO:
313 analyzed the data; MO: wrote the manuscript and all authors revised; AJ, CP, MB, MO, SS, and RW:
314 conceptualized the research. All authors reviewed the manuscript.

ACKNOWLEDGMENTS

315 Funding: This work received no specific grant from any funding agency, commercial, or not-for-profit
316 sectors.

REFERENCES

- 317 Aulakh, J. S., Chahal, P. S., Kumar, V., Price, A. J., and Guillard, K. (2021). Multiple herbicide-resistant
318 Palmer amaranth (*Amaranthus palmeri*) in Connecticut: Confirmation and response to POST herbicides.
319 *Weed Technology* 35, 457–463. doi:10.1017/wet.2021.6.
- 320 Bagavathiannan, M. V., and Norsworthy, J. K. (2016). Multiple-Herbicide Resistance Is Widespread in
321 Roadside Palmer Amaranth Populations. *PLOS ONE* 11, e0148748. doi:10.1371/journal.pone.0148748.
- 322 Baker, R. (2021). Comparative analysis of Palmer amaranth (*Amaranthus palmeri*) and waterhemp (*A.
323 Tuberculatus*) in Iowa. doi:10.31274/etd-20210609-11.
- 324 Bangdiwala, S. I. (2018). Regression: Binary logistic. *International Journal of Injury Control and Safety
325 Promotion* 25, 336–338. doi:10.1080/17457300.2018.1486503.
- 326 Barrett, S. C. H., and Hough, J. (2013). Sexual dimorphism in flowering plants. *Journal of Experimental
327 Botany* 64, 67–82. doi:10.1093/jxb/ers308.
- 328 Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting Linear Mixed-Effects Models Using
329 Lme4. *Journal of Statistical Software* 67, 1–48. doi:10.18637/jss.v067.i01.
- 330 Berger, S. T., Ferrell, J. A., Rowland, D. L., and Webster, T. M. (2015). Palmer Amaranth (*Amaranthus
331 palmeri*) Competition for Water in Cotton. *Weed Science* 63, 928–935. doi:10.1614/WS-D-15-00062.1.

- 332 Bravo, W., Leon, R. G., Ferrell, J. A., Mulvaney, M. J., and Wood, C. W. (2017). Differentiation of
333 Life-History Traits among Palmer Amaranth Populations (*Amaranthus palmeri*) and Its Relation to
334 Cropping Systems and Glyphosate Sensitivity. *Weed Science* 65, 339–349. doi:10.1017/wsc.2017.14.
- 335 Bravo, W., Leon, R. G., Ferrell, J. A., Mulvaney, M. J., and Wood, C. W. (2018). Evolutionary Adaptations
336 of Palmer Amaranth (*Amaranthus palmeri*) to Nitrogen Fertilization and Crop Rotation History Affect
337 Morphology and Nutrient-Use Efficiency. *Weed Science* 66, 180–189. doi:10.1017/wsc.2017.73.
- 338 Briscoe Runquist, R. D., Lake, T., Tiffin, P., and Moeller, D. A. (2019). Species distribution models
339 throughout the invasion history of Palmer amaranth predict regions at risk of future invasion and reveal
340 challenges with modeling rapidly shifting geographic ranges. *Sci Rep* 9, 2426. doi:10.1038/s41598-
341 018-38054-9.
- 342 Chahal, P. S., Barnes, E. R., and Jhala, A. J. (2021). Emergence pattern of Palmer amaranth (*Amaranthus*
343 *palmeri*) influenced by tillage timings and residual herbicides. *Weed Technology* 35, 433–439.
344 doi:10.1017/wet.2020.136.
- 345 Chahal, P. S., Irmak, S., Jugulam, M., and Jhala, A. J. (2018). Evaluating Effect of Degree of Water Stress
346 on Growth and Fecundity of Palmer amaranth (*Amaranthus palmeri*) Using Soil Moisture Sensors.
347 *Weed Science* 66, 738–745. doi:10.1017/wsc.2018.47.
- 348 Correndo, A. A., Moro Rosso, L. H., and Ciampitti, I. A. (2021). Retrieving and processing agro-
349 meteorological data from API-client sources using R software. *BMC Research Notes* 14, 205.
350 doi:10.1186/s13104-021-05622-8.
- 351 Darwin, C. (1888). *The Different Forms of Flowers on Plants of the Same Species*. J. Murray Available at:
352 <http://books.google.com?id=7uMEAAAAYAAJ>.
- 353 Davis, A. S., Schutte, B. J., Hager, A. G., and Young, B. G. (2015). Palmer Amaranth (*Amaranthus palmeri*)
354 Damage Niche in Illinois Soybean Is Seed Limited. *Weed Science* 63, 658–668. doi:10.1614/WS-D-14-
355 00177.1.
- 356 Farmer, J. A., Webb, E. B., Pierce, R. A., and Bradley, K. W. (2017). Evaluating the potential for weed seed
357 dispersal based on waterfowl consumption and seed viability. *Pest Management Science* 73, 2592–2603.
358 doi:10.1002/ps.4710.
- 359 Fisher, R. A. (1930). The genetical theory of natural selection. *Eugen Rev* 22, 127–130. Available at:
360 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2984947/> [Accessed August 12,
361 2021].
- 362 Fox, J., and Weisberg, S. (2018). *An R Companion to Applied Regression*. SAGE Publications Available at:
363 <http://books.google.com?id=uPNrDwAAQBAJ>.
- 364 Garetson, R., Singh, V., Singh, S., Dotray, P., and Bagavathiannan, M. (2019). Distribution of herbicide-
365 resistant Palmer amaranth (*Amaranthus palmeri*) in row crop production systems in Texas. *Weed*
366 *Technology* 33, 355–365. doi:10.1017/wet.2019.14.
- 367 Guo, P., and Al-Khatib, K. (2003). Temperature effects on germination and growth of redroot pigweed
368 (*Amaranthus retroflexus*), Palmer amaranth (*A. palmeri*), and common waterhemp (*A. rudis*). *Weed*
369 *Science* 51, 869–875. doi:10.1614/P2002-127.
- 370 Hartzler, B., and Anderson, M. (2016). Palmer amaranth: It's here, now what? 10.

- 371 Heap, I. (2021). Internation Herbicide-Resistant Weed Database. Available at: <http://www.weedscience.org/Home.aspx> [Accessed July 26, 2021].
- 372
- 373 Hothorn, T., Bretz, F., and Westfall, P. (2008). Simultaneous Inference in General Parametric Models. *Biometrical Journal* 50, 346–363. doi:10.1002/bimj.200810425.
- 374
- 375 Jha, P., Norsworthy, J. K., Riley, M. B., and Bridges, W. (2010). Annual Changes in Temperature and Light Requirements for Germination of Palmer Amaranth (*Amaranthus palmeri*) Seeds Retrieved from Soil. *Weed Science* 58, 426–432. doi:10.1614/WS-D-09-00038.1.
- 376
- 377
- 378 Jhala, A. J., Norsworthy, J. K., Ganie, Z. A., Sosnoskie, L. M., Beckie, H. J., Mallory-Smith, C. A., Liu, J., Wei, W., Wang, J., and Stoltzenberg, D. E. (2021). Pollen-mediated gene flow and transfer of resistance alleles from herbicide-resistant broadleaf weeds. *Weed Technology* 35, 173–187. doi:10.1017/wet.2020.101.
- 379
- 380
- 381
- 382 Keeley, P. E., Carter, C. H., and Thullen, R. J. (1987). Influence of Planting Date on Growth of Palmer Amaranth (*Amaranthus palmeri*). *Weed Science* 35, 199–204. doi:10.1017/S0043174500079054.
- 383
- 384 Kistner, E. J., and Hatfield, J. L. (2018). Potential Geographic Distribution of Palmer Amaranth under Current and Future Climates. *Agricultural & Environmental Letters* 3, 170044. doi:10.2134/ael2017.12.0044.
- 385
- 386
- 387 Klingaman, T. E., and Oliver, L. R. (1994). Palmer Amaranth (*Amaranthus palmeri*) Interference in Soybeans (*Glycine max*). *Weed Science* 42, 523–527. doi:10.1017/S0043174500076888.
- 388
- 389 Kohrt, J. R., Sprague, C. L., Nadakuduti, S. S., and Douches, D. (2017). Confirmation of a Three-Way (Glyphosate, ALS, and Atrazine) Herbicide-Resistant Population of Palmer Amaranth (*Amaranthus palmeri*) in Michigan. *Weed Science* 65, 327–338. doi:10.1017/wsc.2017.2.
- 390
- 391
- 392 Korres, N. E., and Norsworthy, J. K. (2017). Palmer Amaranth (*Amaranthus palmeri*) Demographic and Biological Characteristics in Wide-Row Soybean. *Weed Science* 65, 491–503. doi:10.1017/wsc.2017.12.
- 393
- 394 Korres, N. E., Norsworthy, J. K., FitzSimons, T., Roberts, T. L., and Oosterhuis, D. M. (2017). Differential Response of Palmer Amaranth (*Amaranthus palmeri*) Gender to Abiotic Stress. *Weed Science* 65, 213–227. doi:10.1017/wsc.2016.34.
- 395
- 396
- 397 Korres, N. E., Norsworthy, J. K., Mauromoustakos, A., and Williams, M. M. (2020). Soybean density and Palmer amaranth (*Amaranthus palmeri*) establishment time: Effects on weed biology, crop yield, and economic returns. *Weed Science* 68, 467–475. doi:10.1017/wsc.2020.41.
- 398
- 399
- 400 Kuhn, M., Vaughan, D., and RStudio (2021). *Yardstick: Tidy Characterizations of Model Performance*. Available at: <https://CRAN.R-project.org/package=yardstick> [Accessed August 24, 2021].
- 401
- 402
- 403 Küpper, A., Borgato, E. A., Patterson, E. L., Netto, A. G., Nicolai, M., Carvalho, S. J. P. de, Nissen, S. J., Gaines, T. A., and Christoffoleti, P. J. (2017). Multiple Resistance to Glyphosate and Acetylactate Synthase Inhibitors in Palmer Amaranth (*Amaranthus palmeri*) Identified in Brazil. *Weed Science* 65, 317–326. doi:10.1017/wsc.2017.1.
- 404
- 405
- 406
- 407 Lang, A. (1965). “Physiology of flower initiation,” in *Differenzierung und Entwicklung / Differentiation and Development* Handbuch der Pflanzenphysiologie / Encyclopedia of Plant Physiology., ed. A. Lang (Berlin, Heidelberg: Springer), 1380–1536. doi:10.1007/978-3-642-50088-6_39.
- 408
- 409

- 410 Larran, A. S., Palmieri, V. E., Perotti, V. E., Lieber, L., Tuesca, D., and Permingeat, H. R. (2017). Target-site
411 resistance to acetolactate synthase (ALS)-inhibiting herbicides in Amaranthus palmeri from Argentina.
412 *Pest Management Science* 73, 2578–2584. doi:10.1002/ps.4662.
- 413 Leeper, T. J. (2017). Interpreting Regression Results using Average Marginal Effects with R's margins. 31.
- 414 Leeper, T. J., Arnold, J., Arel-Bundock, V., and Long, J. A. (2021). *Margins: Marginal Effects*
415 *for Model Objects*. Available at: <https://CRAN.R-project.org/package=margins>
416 [Accessed August 24, 2021].
- 417 Lenth, R. V., Buerkner, P., Herve, M., Love, J., Riebl, H., and Singmann, H. (2021). *Emmeans: Estimated*
418 *Marginal Means, aka Least-Squares Means*. Available at: <https://CRAN.R-project.org/package=emmeans> [Accessed August 24, 2021].
- 420 Lindsay, K., Popp, M., Norsworthy, J., Bagavathiannan, M., Powles, S., and Lacoste, M. (2017). PAM:
421 Decision Support for Long-Term Palmer Amaranth (*Amaranthus palmeri*) Control. *Weed Technology*
422 31, 915–927. doi:10.1017/wet.2017.69.
- 423 MacLaren, C., Storkey, J., Menegat, A., Metcalfe, H., and Dehnen-Schmutz, K. (2020). An ecological
424 future for weed science to sustain crop production and the environment. A review. *Agron. Sustain. Dev.*
425 40, 24. doi:10.1007/s13593-020-00631-6.
- 426 MacRae, A. W., Webster, T. M., Sosnoskie, L. M., Culpepper, A. S., and Kichler, J. M. (2013). Cotton
427 Yield Loss Potential in Response to Length of Palmer Amaranth (*Amaranthus palmeri*) Interference.
428 17, 6.
- 429 Mangiafico, S. (2021). *Rcompanion: Functions to Support Extension Education Program Evaluation*.
430 Available at: <https://CRAN.R-project.org/package=rcompanion> [Accessed August
431 24, 2021].
- 432 Massinga, R. A., Currie, R. S., Horak, M. J., and Boyer, J. (2001). Interference of Palmer amaranth in corn.
433 *Weed Science* 49, 202–208. doi:10.1614/0043-1745(2001)049[0202:IOPAIC]2.0.CO;2.
- 434 Matzrafi, M., Osipitan, O. A., Ohadi, S., and Mesgaran, M. B. (2021). Under pressure: Maternal effects
435 promote drought tolerance in progeny seed of Palmer amaranth (*Amaranthus palmeri*). *Weed Science*
436 69, 31–38. doi:10.1017/wsc.2020.75.
- 437 McFarlane, G. R., Whitelaw, C. B. A., and Lillico, S. G. (2018). CRISPR-Based Gene Drives for Pest
438 Control. *Trends in Biotechnology* 36, 130–133. doi:10.1016/j.tibtech.2017.10.001.
- 439 Menges, R. M. (1987). Weed Seed Population Dynamics during Six Years of Weed Management Systems
440 in Crop Rotations on Irrigated Soil. *Weed Science* 35, 328–332. Available at: <http://www.jstor.org/stable/4044593>.
- 442 Mesgaran, M. B., Matzrafi, M., and Ohadi, S. (2021). Sex dimorphism in dioecious Palmer amaranth
443 (*Amaranthus palmeri*) in response to water stress. *Planta* 254, 17. doi:10.1007/s00425-021-03664-7.
- 444 Milani, A., Panozzo, S., Farinati, S., Iamonico, D., Sattin, M., Loddo, D., and Scarabel, L. (2021). Recent
445 Discovery of *Amaranthus palmeri* S. Watson in Italy: Characterization of ALS-Resistant Populations
446 and Sensitivity to Alternative Herbicides. *Sustainability* 13, 7003. doi:10.3390/su13137003.
- 447 Montgomery, J. S., Giacomini, D. A., Weigel, D., and Tranel, P. J. (2021). Male-specific Y-chromosomal
448 regions in waterhemp (*Amaranthus tuberculatus*) and Palmer amaranth (*Amaranthus palmeri*). *New*
449 *Phytologist* 229, 3522–3533. doi:10.1111/nph.17108.

- 450 Montgomery, J. S., Sadeque, A., Giacomini, D. A., Brown, P. J., and Tranel, P. J. (2019). Sex-specific
451 markers for waterhemp (*Amaranthus tuberculatus*) and Palmer amaranth (*Amaranthus palmeri*). *Weed
452 Science* 67, 412–418. doi:10.1017/wsc.2019.27.
- 453 Morgan, G. D., Baumann, P. A., and Chandler, J. M. (2001). Competitive Impact of Palmer Amaranth
454 (*Amaranthus palmeri*) on Cotton (*Gossypium hirsutum*) Development and Yield. *Weed Technology* 15,
455 408–412. doi:10.1614/0890-037X(2001)015[0408:CIOPAA]2.0.CO;2.
- 456 Nguyen, H. M., Kleitou, P., Kletou, D., Sapir, Y., and Winters, G. (2018). Differences in flowering sex
457 ratios between native and invasive populations of the seagrass *Halophila stipulacea*. *Botanica Marina*
458 61, 337–342. doi:10.1515/bot-2018-0015.
- 459 Oliveira, M. C. (2021). *Maxwelco/palmer_adaptation*. Zenodo Available at: <https://doi.org/10.5281/zenodo.5236831>.
- 460
- 461 Oliveira, M. C., Gaines, T. A., Patterson, E. L., Jhala, A. J., Irmak, S., Amundsen, K., and Knezevic, S.
462 Z. (2018). Interspecific and intraspecific transference of metabolism-based mesotrione resistance in
463 dioecious weedy *Amaranthus*. *The Plant Journal* 96, 1051–1063. doi:10.1111/tpj.14089.
- 464 Oliveira, M. C., Giacomini, D. A., Arsenijevic, N., Vieira, G., Tranel, P. J., and Werle, R. (2021).
465 Distribution and validation of genotypic and phenotypic glyphosate and PPO-inhibitor resistance in
466 Palmer amaranth (*Amaranthus palmeri*) from southwestern Nebraska. *Weed Technology* 35, 65–76.
467 doi:10.1017/wet.2020.74.
- 468 Page, E. R., Nurse, R. E., Meloche, S., Bosveld, K., Grainger, C., Obeid, K., Filotas, M., Simard, M.-J., and
469 Laforest, M. (2021). Import of Palmer amaranth (*Amaranthus palmeri* S. Wats.) Seed with sweet potato
470 (*Ipomea batatas* (L.) Lam) slips. *Can. J. Plant Sci.*, CJPS-2020-0321. doi:10.1139/CJPS-2020-0321.
- 471 Pérez-Llorca, M., and Sánchez Vilas, J. (2019). Sexual dimorphism in response to herbivory and
472 competition in the dioecious herb *Spinacia oleracea*. *Plant Ecol* 220, 57–68. doi:10.1007/s11258-
473 018-0902-7.
- 474 Piskackova, T. A. R., Reberg-Horton, S. C., Richardson, R. J., Jennings, K. M., Franca, L., Young, B. G.,
475 and Leon, R. G. (2021). Windows of action for controlling palmer amaranth (*Amaranthus palmeri*)
476 using emergence and phenology models. *Weed Research* 61, 188–198. doi:10.1111/wre.12470.
- 477 Price, A. J., Balkcom, K. S., Culpepper, S. A., Kelton, J. A., Nichols, R. L., and Schomberg, H. (2011).
478 Glyphosate-resistant Palmer amaranth: A threat to conservation tillage. *Journal of Soil and Water
479 Conservation* 66, 265–275. doi:10.2489/jswc.66.4.265.
- 480 Raoniar, R. (2021). Modelling Binary Logistic Regression Using R (research-oriented modelling and
481 interpretation). Available at: <https://bit.ly/3BhucN3> [Accessed August 11, 2021].
- 482 Ritz, C., Baty, F., Streibig, J. C., and Gerhard, D. (2015). Dose-Response Analysis Using R. *PLOS ONE*
483 10, e0146021. doi:10.1371/journal.pone.0146021.
- 484 Rumpa, M. M., Krausz, R. F., Gibson, D. J., and Gage, K. L. (2019). Effect of PPO-Inhibiting Herbicides
485 on the Growth and Sex Ratio of a Dioecious Weed Species *Amaranthus palmeri* (Palmer Amaranth).
486 *Agronomy* 9, 275. doi:10.3390/agronomy9060275.
- 487 Sauer, J. (1957). Recent Migration and Evolution of the Dioecious Amaranths. *Evolution* 11, 11–31.
488 doi:10.2307/2405808.

- 489 Sauer, J. D. (1972). The dioecious amaranths: A new species name and major range extensions. *Madroño*
490 21, 426–434. Available at: <http://www.jstor.org/stable/41423815>.
- 491 Sbatella, G. M., and Wilson, R. G. (2010). Isoxaflutole Shifts Kochia (*Kochia scoparia*) Populations in
492 Continuous Corn. *Weed Technology* 24, 392–396. doi:10.1614/WT-D-09-00023.1.
- 493 Schliekelman, P., Ellner, S., and Gould, F. (2005). Pest Control by Genetic Manipulation of Sex Ratio.
494 *Journal of Economic Entomology* 98, 18–34. doi:10.1093/jee/98.1.18.
- 495 Schwartz, L. M., Norsworthy, J. K., Young, B. G., Bradley, K. W., Kruger, G. R., Davis, V. M., Steckel,
496 L. E., and Walsh, M. J. (2016). Tall Waterhemp (*Amaranthus tuberculatus*) and Palmer amaranth
497 (*Amaranthus palmeri*) Seed Production and Retention at Soybean Maturity. *Weed Technology* 30,
498 284–290. doi:10.1614/WT-D-15-00130.1.
- 499 Scott, D. (2011). The Technological Fix Criticisms and the Agricultural Biotechnology Debate. *J Agric
500 Environ Ethics* 24, 207–226. doi:10.1007/s10806-010-9253-7.
- 501 Sing, T., Sander, O., Beerenwinkel, N., and Lengauer, T. (2005). ROCR: Visualizing classifier performance
502 in R. *Bioinformatics* 21, 3940–3941. doi:10.1093/bioinformatics/bti623.
- 503 Sosnoskie, L. M., Webster, T. M., and Culpepper, A. S. (2013). Glyphosate Resistance Does Not
504 Affect Palmer Amaranth (*Amaranthus palmeri*) Seedbank Longevity. *Weed Science* 61, 283–288.
505 doi:10.1614/WS-D-12-00111.1.
- 506 Spaunhorst, D. J., Devkota, P., Johnson, W. G., Smeda, R. J., Meyer, C. J., and Norsworthy, J. K. (2018).
507 Phenology of Five Palmer amaranth (*Amaranthus palmeri*) Populations Grown in Northern Indiana and
508 Arkansas. *Weed Science* 66, 457–469. doi:10.1017/wsc.2018.12.
- 509 Team, R. C. (2021). *R: The R Project for Statistical Computing*. Vienna, Austria: R Foundation for Statistical
510 Computing Available at: <https://www.r-project.org/> [Accessed August 24, 2021].
- 511 Tranel, P. J., Riggins, C. W., Bell, M. S., and Hager, A. G. (2011). Herbicide Resistances in *Amaranthus
512 tuberculatus*: A Call for New Options. *J. Agric. Food Chem.* 59, 5808–5812. doi:10.1021/jf103797n.
- 513 Van Wychen, L. (2020). 2020 Survey of the most common and troublesome weeds in grass crops, pasture,
514 and turf in the United States and Canada. Available at: [https://wssa.net/wp-content/
uploads/2020-Weed-Survey_grass-crops.xlsx](https://wssa.net/wp-content/
515 uploads/2020-Weed-Survey_grass-crops.xlsx).
- 516 Wang, J. L., Klessig, D. F., and Berry, J. O. (1992). Regulation of C4 Gene Expression in Developing
517 Amaranth Leaves. *The Plant Cell* 4, 173–184. doi:10.1105/tpc.4.2.173.
- 518 Ward, S. M., Webster, T. M., and Steckel, L. E. (2013). Palmer Amaranth (*Amaranthus palmeri*): A Review.
519 *Weed Technology* 27, 12–27. doi:10.1614/WT-D-12-00113.1.
- 520 Webster, T. M., and Grey, T. L. (2015). Glyphosate-Resistant Palmer Amaranth (*Amaranthus palmeri*)
521 Morphology, Growth, and Seed Production in Georgia. *Weed Science* 63, 264–272. doi:10.1614/WS-D-
522 14-00051.1.
- 523 Wu, C., and Owen, M. D. K. (2014). When Is the Best Time to Emerge: Reproductive Phenology and
524 Success of Natural Common Waterhemp (*Amaranthus rudis*) Cohorts in the Midwest United States?
525 *Weed Science* 62, 107–117. Available at: <http://www.jstor.org/stable/43700638>.
- 526 Yacoubi, R., and Axman, D. (2020). Probabilistic Extension of Precision, Recall, and F1 Score for
527 More Thorough Evaluation of Classification Models. in *Proceedings of the First Workshop on*

- 528 *Evaluation and Comparison of NLP Systems* (Online: Association for Computational Linguistics),
529 79–91. doi:10.18653/v1/2020.eval4nlp-1.9.
- 530 Yu, E., Blair, S., Hardel, M., Chandler, M., Thiede, D., Cortilet, A., Gunsolus, J., and Becker, R. (2021).
531 Timeline of Palmer amaranth (*Amaranthus palmeri*) invasion and eradication in Minnesota. *Weed
532 Technology*, 1–31. doi:10.1017/wet.2021.32.




Figure 1. Mean average temperature (C) and total monthly precipitation (mm) at Arlington, WI, Clay Center, NE, Grant, NE, Lincoln, NE and Macomb, IL




Figure 2. Palmer amaranth adaptation study layout of a plant cohort timing in bareground, corn, and soybean. Twenty-four Palmer amaranth plants were place 76.2 cm apart in each field experimental unit

FIGURES




Figure 3. Palmer amaranth biomass (A) and height (B) growing in corn, bareground, and soybean nested across Arlington, WI, Clay Center, NE, Grant, NE, Lincoln, NE and Macomb, IL




Figure 4. Cumulative flowering of Palmer amaranth at first and second transplant timing (A) and day of year of 10, 50, and 90 cumulative flowering at first and second cohort transplanting time (B) nested across Arlington, WI, Clay Center, NE, Grant, NE, Lincoln, NE and Macomb, IL




Figure 5. The probability (P) of being female Palmer amaranth by day of year (A), weight (B), and height (C). Black line represents the model estimation and shaded green the confidence intervals




Figure 6. MCO (180 cm) holds harvested Palmer amaranth plants at 40 days after first cohort transplanting (A) and 33 days after second cohort transplanting (B) time. From left to right in each image, Palmer amaranth growing in bareground, soybean and corn in Arlington, Wisconsin