

Linear Constant-Coefficient Difference Equations

Maxx Seminario
University of Nebraska-Lincoln
September 3, 2025

Introduction to Difference Equations

- **Important Class of LTI Systems:** Systems where input $x[n]$ and output $y[n]$ satisfy a difference equation

- **General Form:**

$$\sum_{k=0}^N a_k y[n-k] = \sum_{m=0}^M b_m x[n-m]$$

- **Parameters:**

- N : Order of the system (highest delay in output)
- a_k : Output coefficients (constant)
- b_m : Input coefficients (constant)
- M : Highest delay in input terms

- **Why Important?:**

- Provides computational algorithms for LTI systems
- Foundation for digital filter implementation
- Connects time-domain and system analysis

Example 1: The Accumulator System

ECEN 463/863

Maxx Seminario

Introduction

System

Block Diagram

Block Diagram

Block Diagram

Block Diagram

Block Diagram

Block Diagram

Problem: Find the difference equation for the accumulator system.

System Definition:

$$y[n] = \sum_{k=-\infty}^n x[k]$$

Approach:

- Rewrite the sum to separate current and past inputs
- Use the relationship between $y[n]$ and $y[n-1]$

Example 1: Accumulator Solution

ECEN 463/863

Maxx Seminario

Introduction

System

Block Diagram

Block Diagram

Block Diagram

Block Diagram

Block Diagram

Block Diagram

Step 1: Rewrite the accumulator equation:

$$y[n] = x[n] + \sum_{k=-\infty}^{n-1} x[k]$$

Step 2: Recognize that the sum is $y[n-1]$:

$$y[n-1] = \sum_{k=-\infty}^{n-1} x[k]$$


Step 3: Substitute to get the difference equation:

$$y[n] = x[n] + y[n-1]$$

Standard Form:

$$y[n] - y[n-1] = x[n]$$

Block Diagram: Recursive Accumulator



Recursive Implementation:

- Each output value computed using previously computed values
- $y[n] = x[n] + y[n-1]$: Add current input to previous output
- Requires initial condition (e.g., $y[-1] = 0$)

Example 2: Moving Average System

Problem: Find difference equation for causal moving average system.

System Definition (with $M_1 = 0$, so the system is causal):

$$y[n] = \frac{1}{M_2 + 1} \sum_{k=0}^{M_2} x[n - k]$$

Two Approaches:

- 1 **Direct (Non-recursive):** Use convolution form directly
- 2 **Recursive:** Express as cascade of simpler systems

Moving Average: Direct Implementation

Direct Form:

$$y[n] = \frac{1}{M_2 + 1} \sum_{k=0}^{M_2} x[n - k]$$


Standard Form:

$$y[n] = \sum_{k=0}^{M_2} \frac{1}{M_2 + 1} x[n - k]$$

Disadvantages of Direct Implementation:

- Requires $(M_2 + 1)$ multiplications per output sample
- Must store $(M_2 + 1)$ input samples in memory
- $O(M_2)$: Computational cost grows linearly with window size M_2

Block Diagram: Recursive Moving Average



Signal Flow:

- Input $x[n] \rightarrow$ Attenuator $\frac{1}{M_2+1}$
- Attenuated signal splits: direct path and $(M_2 + 1)$ sample delay
- Sum: $x_1[n] = \frac{1}{M_2+1} [x[n] - x[n - (M_2 + 1)]]$
- $x_1[n] \rightarrow$ Accumulator \rightarrow Output $y[n]$

Moving Average as Difference Equation

ECEN 463/863

Maxx Seminar

Moving Average
System

Intermediate signal:

$$x_1[n] = \frac{1}{M_2 + 1} [x[n] - x[n - M_2 - 1]]$$

Accumulator relation: $y[n] = x_1[n] + y[n - 1]$

Final recursive form:

$$y[n] - y[n - 1] = \frac{1}{M_2 + 1} [x[n] - x[n - M_2 - 1]]$$

Note: There is an unlimited number of distinct difference equations to represent an LTI I/O relation.

General Solution of Difference Equations

ECEN 463/863

Maxx Seminar

General Solution

Key Issue: Difference equation alone does not uniquely specify output!

General Solution Structure:

$$y[n] = y_p[n] + y_h[n]$$

- $y_p[n]$: Particular solution (satisfies original equation)
- $y_h[n]$: Homogeneous solution (satisfies homogeneous equation)

Homogeneous Equation ($x[n] = 0$):

$$\sum_{k=0}^N a_k y_h[n - k] = 0$$

Homogeneous Solution Form:

$$y_h[n] = \sum_{m=1}^N A_m z_m^n$$

where z_m are roots of characteristic polynomial $A(z) = \sum_{k=0}^N a_k z^{-k} = 0$

Auxiliary Conditions

ECEN 463/863

Maxx Seminario

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Need for Auxiliary Conditions:

- N undetermined coefficients A_m in homogeneous solution
- Need N auxiliary (boundary) conditions for unique solution

Types of Auxiliary Conditions:

- 1 **Fixed Values:** Specify $y[-1], y[-2], \dots, y[-N]$
- 2 **Initial Rest:** If $x[n] = 0$ for $n < n_0$, then $y[n] = 0$ for $n < n_0$

Forward and Backward Computation for Specific Class of Difference Equations

ECEN 463/863

Maxx Seminario

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Particular Solution

Specific Class of Difference Equations:

- Inputs $x[n]$ and Outputs $y[n]$ satisfy an N th-order linear constant coefficient difference equation:

$$\sum_{k=0}^N a_k y[n-k] = \sum_{k=0}^M b_k x[n-k]$$

$$a_0 y[n] + a_1 y[n-1] + \dots + a_N y[n-N] = b_0 x[n] + b_1 x[n-1] + \dots + b_M x[n-M]$$

Recursive Computation (forward):

$$y[n] = - \sum_{k=1}^N \frac{a_k}{a_0} y[n-k] + \sum_{k=0}^M \frac{b_k}{a_0} x[n-k]$$

Recursive Computation (backward):

$$y[n-N] = - \sum_{k=0}^{N-1} \frac{a_k}{a_N} y[n-k] + \sum_{k=0}^M \frac{b_k}{a_N} x[n-k]$$

Example 3: First-Order Difference Equation

ECEN 463/863
Maxx Seminario

Discrete-Time
Signals and
Systems
Particular
Solution

Problem: Solve the difference equation with given input and initial condition.

Difference Equation:
$$y[n] = ay[n - 1] + x[n]$$

Input:
$$x[n] = K\delta[n]$$

Auxiliary Condition:
$$y[-1] = c$$

Find: The complete solution $y[n]$ for all n .

Example 3: Solution for $n > 0$

ECEN 463/863
Maxx Seminario

Discrete-Time
Signals and
Systems
Particular
Solution

Method: Use recursive computation starting from $n = 0$.

For $n = 0$:

$$\begin{aligned} y[0] &= ay[-1] + x[0] & (1) \\ &= ay[-1] + K\delta[0] & (2) \\ &= ac + K \cdot 1 & (3) \\ &= ac + K & (4) \end{aligned}$$

For $n = 1$:

$$\begin{aligned} y[1] &= ay[0] + x[1] & (5) \\ &= a(ac + K) + K\delta[1] & (6) \\ &= a^2c + aK + K \cdot 0 & (7) \\ &= a^2c + aK & (8) \end{aligned}$$

Example 3: Solution for $n > 0$

For $n = 2$:

$$\begin{aligned} y[2] &= ay[1] + x[2] & (9) \\ &= a(a^2c + aK) + K\delta[2] & (10) \\ &= a^3c + a^2K + 0 & (11) \\ &= a^3c + a^2K & (12) \end{aligned}$$

For $n = 3$:

$$\begin{aligned} y[3] &= ay[2] + x[3] & (13) \\ &= a(a^3c + a^2K) + 0 & (14) \\ &= a^4c + a^3K & (15) \end{aligned}$$

Pattern Recognition: For $n \geq 0$: $y[n] = a^{n+1}c + a^nK$

Example 3: Solution for $n < 0$

For $n < 0$: Use backward recursion.

Rearrange difference equation:

$$y[n-1] = \frac{y[n] - x[n]}{a}$$

For $n = 0$ (find $y[-1]$):

$$y[-1] = \frac{y[0] - x[0]}{a} = \frac{(ac + K) - K}{a} = \frac{ac}{a} = c$$

This confirms our auxiliary condition

For $n = -1$ (find $y[-2]$):

$$y[-2] = \frac{y[-1] - x[-1]}{a} = \frac{c - 0}{a} = \frac{c}{a}$$

General pattern for $n < 0$:

$$y[n] = ca^{n+1} = \frac{c}{a^{n+1}}$$

Example 3: Complete Solution

Final Answer:

$$y[n] = \begin{cases} ca^{n+1}, & n < 0 \\ a^n(ac + K), & n \geq 0 \end{cases}$$

Alternative Compact Form:

$$y[n] = ca^{n+1} + Ka^n u[n]$$

where $u[n]$ is the unit step function.

Physical Interpretation:

- **Homogeneous part:** ca^{n+1} (due to initial condition)
- **Forced response:** $Ka^n u[n]$ (due to impulse input)
- **System behavior:** Exponential with base a
- **Stability:** System stable if $|a| < 1$

Summary: Linear Constant-Coefficient Difference Equations

■ **Definition:**

$$\sum_{k=0}^N a_k y[n-k] = \sum_{m=0}^M b_m x[n-m]$$

Describes the relationship between input $x[n]$ and output $y[n]$ in LTI systems.

■ **Key Components:**

- N and M : Order of the system
- a_k, b_m : Constant coefficients
- Recursive and direct computation methods

■ **General Solution:**

$$y[n] = y_p[n] + y_h[n]$$

- $y_p[n]$: Particular solution (due to input $x[n]$)
 - $y_h[n]$: Homogeneous solution (due to initial conditions)
 - Auxiliary conditions: Necessary to determine a unique / particular solutions.
- **Applications:** Digital filters, computational algorithms for LTI systems, and time-domain analysis.