

The Discrete Fourier Transform

Maxx Seminario

University of Nebraska-Lincoln

October 3, 2025

Overview: Representation of Sequences

- **Motivation:** How to represent arbitrary input sequences in frequency domain?
 - We know LTI system response to $e^{j\omega n}$ is $H(e^{j\omega})e^{j\omega n}$
 - Need to decompose arbitrary signals into complex exponentials
- **Key Questions:**
 - Can we represent any sequence as a sum of complex exponentials?
 - What are the conditions for such representations to exist?
 - How do we compute the representation?
- **Today's Topics:**
 - Fourier Transform pair for discrete-time signals
 - Convergence conditions
 - Symmetry properties
 - Fourier Transform theorems

The Discrete-Time Fourier Transform (DTFT)

Fourier Transform Pair:

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega}) e^{j\omega n} d\omega \quad (\text{Synthesis}) \quad (1)$$

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \quad (\text{Analysis}) \quad (2)$$

Key Interpretations:

- **Synthesis:** Represents $x[n]$ as superposition of complex sinusoids
- **Analysis:** Determines "how much" of each frequency is in $x[n]$
- $X(e^{j\omega})$ is generally complex-valued
- Integration can be over any interval of length 2π

Connection to LTI Systems:

$$H(e^{j\omega}) = \sum_{n=-\infty}^{\infty} h[n] e^{-j\omega n}$$

Frequency response = DTFT of impulse response

Complex Representation of DTFT

Rectangular Form:

$$X(e^{j\omega}) = X_R(e^{j\omega}) + jX_I(e^{j\omega})$$

Polar Form:

$$X(e^{j\omega}) = |X(e^{j\omega})| e^{j\angle X(e^{j\omega})}$$

Phase Considerations:

- Phase is not unique (can add multiples of 2π)
- **Principal value:** $\text{ARG}[X(e^{j\omega})] \in [-\pi, \pi]$
- **Continuous phase:** $\arg[X(e^{j\omega})]$ (unwrapped phase)

Periodicity:

- $X(e^{j\omega})$ is periodic in ω with period 2π
- Similar to Fourier series for periodic functions
- Eq. (2.131) is Fourier series of periodic function $X(e^{j\omega})$

Convergence of the DTFT

ECEN 463/863
Maxx Seminario

Question: When does $X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$ converge?
Absolute Summability (Sufficient Condition):

$$|X(e^{j\omega})| = \left| \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n} \right| \quad (3)$$

$$\leq \sum_{n=-\infty}^{\infty} |x[n]| |e^{-j\omega n}| \quad (4)$$

$$= \sum_{n=-\infty}^{\infty} |x[n]| < \infty \quad (5)$$

Implications:

- If $\sum_{n=-\infty}^{\infty} |x[n]| < \infty$, then $X(e^{j\omega})$ exists
- Series converges uniformly to a continuous function
- All stable sequences have Fourier transforms
- All FIR systems have finite, continuous frequency responses

Digital Signal Processing, Fall 2025

5 / 28 Digital Signal Processing, Fall 2025

Example: Suddenly-Applied Exponential

ECEN 463/863
Maxx Seminario

Sequence: $x[n] = a^n u[n]$
DTFT Calculation:

$$X(e^{j\omega}) = \sum_{n=0}^{\infty} a^n e^{-j\omega n} \quad (6)$$

$$= \sum_{n=0}^{\infty} (ae^{-j\omega})^n \quad (7)$$

$$= \frac{1}{1 - ae^{-j\omega}} \quad \text{if } |ae^{-j\omega}| < 1 \quad (8)$$

Convergence Condition: $|a| < 1$

Absolute Summability Check:

$$\sum_{n=0}^{\infty} |a|^n = \frac{1}{1 - |a|} < \infty \quad \text{if } |a| < 1$$

This confirms that absolute summability guarantees convergence.

6 / 28

Square Summability

ECEN 463/863
Maxx Seminario

Alternative Condition: Some sequences are not absolutely summable but are square summable:

$$\sum_{n=-\infty}^{\infty} |x[n]|^2 < \infty$$

Mean-Square Convergence:

$$\lim_{M \rightarrow \infty} \int_{-\pi}^{\pi} |X(e^{j\omega}) - X_M(e^{j\omega})|^2 d\omega = 0$$

where $X_M(e^{j\omega}) = \sum_{n=-M}^M x[n]e^{-j\omega n}$

Important Distinction:

- Error may not approach zero at each ω
- Total "energy" in error approaches zero
- Leads to Gibbs phenomenon at discontinuities

Digital Signal Processing, Fall 2025

7 / 28 Digital Signal Processing, Fall 2025

Example: Ideal Lowpass Filter

ECEN 463/863
Maxx Seminario

Frequency Response:

$$H_{lp}(e^{j\omega}) = \begin{cases} 1, & |\omega| < \omega_c \\ 0, & \omega_c < |\omega| \leq \pi \end{cases}$$

Impulse Response:

$$h_{lp}[n] = \frac{1}{2\pi} \int_{-\omega_c}^{\omega_c} e^{j\omega n} d\omega \quad (9)$$

$$= \frac{1}{2\pi j n} [e^{j\omega_c n} - e^{-j\omega_c n}] \quad (10)$$

$$= \frac{\sin(\omega_c n)}{\pi n}, \quad -\infty < n < \infty \quad (11)$$

Properties:


- **Noncausal:** $h_{lp}[n] \neq 0$ for $n < 0$
- **Not absolutely summable:** Decays as $1/n$
- **Square summable:** Mean-square convergence

8 / 28

Gibbs Phenomenon

Finite Sum Approximation:

$$H_M(e^{j\omega}) = \sum_{n=-M}^M \frac{\sin(\omega_c n)}{\pi n} e^{-j\omega n}$$



Key Points:

- Oscillations near discontinuity persist as $M \rightarrow \infty$
- Maximum overshoot $\approx 9\%$ (doesn't decrease)
- Oscillations become more rapid but localized

Fourier Transform of Special Sequences

Case 1: Constant Sequence

- Sequence: $x[n] = 1$ for all n
- Neither absolutely nor square summable
- DTFT (using impulse functions):

$$X(e^{j\omega}) = \sum_{r=-\infty}^{\infty} 2\pi\delta(\omega + 2\pi r)$$

Case 2: Complex Exponential

- Sequence: $x[n] = e^{j\omega_0 n}$, $-\pi < \omega_0 \leq \pi$
- DTFT:

$$X(e^{j\omega}) = \sum_{r=-\infty}^{\infty} 2\pi\delta(\omega - \omega_0 + 2\pi r)$$

- Verification: Substituting into synthesis equation yields $x[n] = e^{j\omega_0 n}$

Unit Step Sequence

Sequence: $u[n]$

DTFT:

$$U(e^{j\omega}) = \frac{1}{1 - e^{-j\omega}} + \sum_{r=-\infty}^{\infty} \pi\delta(\omega + 2\pi r)$$

Interpretation:

- First term: Response for $\omega \neq 0$
- Second term: DC component (impulses at $\omega = 2\pi r$)
- Neither absolutely nor square summable
- Requires generalized function theory

General Form: For sequences with discrete frequency components:

$$x[n] = \sum_k a_k e^{j\omega_k n} \Rightarrow X(e^{j\omega}) = \sum_{r=-\infty}^{\infty} \sum_k 2\pi a_k \delta(\omega - \omega_k + 2\pi r)$$

Conjugate Symmetry Definitions

For Sequences:

- **Conjugate-symmetric:** $x_e[n] = x_e^*[-n]$
- **Conjugate-antisymmetric:** $x_o[n] = -x_o^*[-n]$

Decomposition: Any sequence can be written as:

$$x[n] = x_e[n] + x_o[n] \quad (12)$$

$$x_e[n] = \frac{1}{2}(x[n] + x^*[-n]) \quad (13)$$

$$x_o[n] = \frac{1}{2}(x[n] - x^*[-n]) \quad (14)$$

For Real Sequences:

- **Even:** $x_e[n] = x_e[-n]$ (conjugate-symmetric when real)
- **Odd:** $x_o[n] = -x_o[-n]$ (conjugate-antisymmetric when real)

Symmetry Properties of DTFT

ECEN 463/863
Maxx Seminario

General Complex Sequences:

| Sequence | Fourier Transform |
|----------------------|--|
| $x^*[n]$ | $X^*(e^{-j\omega})$ |
| $x^*[-n]$ | $X^*(e^{j\omega})$ |
| $\text{Re}\{x[n]\}$ | $X_e(e^{j\omega})$ (conjugate-symmetric part) |
| $j\text{Im}\{x[n]\}$ | $X_o(e^{j\omega})$ (conjugate-antisymmetric part) |
| $x_e[n]$ | $X_R(e^{j\omega}) = \text{Re}\{X(e^{j\omega})\}$ |
| $x_o[n]$ | $jX_I(e^{j\omega}) = j\text{Im}\{X(e^{j\omega})\}$ |

Key Insight:

- Conjugate-symmetric sequences ↔ Real transforms
- Conjugate-antisymmetric sequences ↔ Imaginary transforms

Symmetry Properties for Real Sequences

ECEN 463/863
Maxx Seminario

If $x[n]$ is real:

| Property | Result |
|---------------------|--|
| DTFT | $X(e^{j\omega}) = X^*(e^{-j\omega})$ (conjugate symmetric) |
| Real part | $X_R(e^{j\omega}) = X_R(e^{-j\omega})$ (even function) |
| Imaginary part | $X_I(e^{j\omega}) = -X_I(e^{-j\omega})$ (odd function) |
| Magnitude | $ X(e^{j\omega}) = X(e^{-j\omega}) $ (even function) |
| Phase | $\angle X(e^{j\omega}) = -\angle X(e^{-j\omega})$ (odd function) |
| Even part of $x[n]$ | Transforms to $X_R(e^{j\omega})$ |
| Odd part of $x[n]$ | Transforms to $jX_I(e^{j\omega})$ |

Practical Implication: For real signals, we only need to compute DTFT for $\omega \in [0, \pi]$

Example: Symmetry Properties

ECEN 463/863
Maxx Seminario

Sequence: $x[n] = a^n u[n]$, $|a| < 1$ (real)

DTFT: $X(e^{j\omega}) = \frac{1}{1 - ae^{-j\omega}}$

Components:

$$X_R(e^{j\omega}) = \frac{1 - a \cos \omega}{1 + a^2 - 2a \cos \omega} \quad (\text{even}) \tag{15}$$

$$X_I(e^{j\omega}) = \frac{-a \sin \omega}{1 + a^2 - 2a \cos \omega} \quad (\text{odd}) \tag{16}$$

$$|X(e^{j\omega})| = \frac{1}{(1 + a^2 - 2a \cos \omega)^{1/2}} \quad (\text{even}) \tag{17}$$

$$\angle X(e^{j\omega}) = \tan^{-1} \left(\frac{-a \sin \omega}{1 - a \cos \omega} \right) \quad (\text{odd}) \tag{18}$$

Verification: Each property satisfies the symmetry conditions for real sequences

Overview of DTFT Theorems

ECEN 463/863
Maxx Seminario

Key Theorems:

- 1 **Linearity:** $ax[n] + by[n] \leftrightarrow aX(e^{j\omega}) + bY(e^{j\omega})$
- 2 **Time Shifting:** $x[n - n_d] \leftrightarrow e^{-j\omega n_d} X(e^{j\omega})$
- 3 **Frequency Shifting:** $e^{j\omega_0 n} x[n] \leftrightarrow X(e^{j(\omega - \omega_0)})$
- 4 **Time Reversal:** $x[-n] \leftrightarrow X(e^{-j\omega})$
- 5 **Differentiation:** $nx[n] \leftrightarrow j \frac{dX(e^{j\omega})}{d\omega}$
- 6 **Convolution:** $x[n] * y[n] \leftrightarrow X(e^{j\omega})Y(e^{j\omega})$
- 7 **Multiplication:** $x[n]y[n] \leftrightarrow \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\theta})Y(e^{j(\omega - \theta)})d\theta$

Why Important?

- Simplify analysis of complex signals
- Transform difficult operations into simple ones
- Foundation for filter design and signal processing

Linearity and Time Shifting

1. Linearity:

$$ax_1[n] + bx_2[n] \xleftrightarrow{\mathcal{F}} aX_1(e^{j\omega}) + bX_2(e^{j\omega})$$

2. Time Shifting:

$$x[n - n_d] \xleftrightarrow{\mathcal{F}} e^{-j\omega n_d} X(e^{j\omega})$$

Proof of Time Shifting:

$$\mathcal{F}\{x[n - n_d]\} = \sum_{n=-\infty}^{\infty} x[n - n_d] e^{-j\omega n} \quad (19)$$

$$= \sum_{m=-\infty}^{\infty} x[m] e^{-j\omega(m+n_d)} \quad (m = n - n_d) \quad (20)$$

$$= e^{-j\omega n_d} \sum_{m=-\infty}^{\infty} x[m] e^{-j\omega m} \quad (21)$$

$$= e^{-j\omega n_d} X(e^{j\omega}) \quad (22)$$

Frequency Shifting and Time Reversal

3. Frequency Shifting (Modulation):

$$e^{j\omega_0 n} x[n] \xleftrightarrow{\mathcal{F}} X(e^{j(\omega - \omega_0)})$$

Application: Modulation shifts spectrum by ω_0

4. Time Reversal:

$$x[-n] \xleftrightarrow{\mathcal{F}} X(e^{-j\omega})$$

For real sequences: $x[-n] \xleftrightarrow{\mathcal{F}} X^*(e^{j\omega})$

Example Application:

- Signal $x[n] = \cos(\omega_0 n) = \frac{1}{2}(e^{j\omega_0 n} + e^{-j\omega_0 n})$
- Using frequency shifting: spectrum has impulses at $\pm\omega_0$

Differentiation in Frequency

5. Differentiation Theorem:

$$nx[n] \xleftrightarrow{\mathcal{F}} j \frac{dX(e^{j\omega})}{d\omega}$$

Proof Sketch:

$$\frac{dX(e^{j\omega})}{d\omega} = \frac{d}{d\omega} \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \quad (23)$$

$$= \sum_{n=-\infty}^{\infty} x[n] (-jn) e^{-j\omega n} \quad (24)$$

$$= -j \sum_{n=-\infty}^{\infty} nx[n] e^{-j\omega n} \quad (25)$$

Applications:

- Finding moments of sequences
- Analyzing group delay of filters
- Computing derivatives of frequency response

Parseval's Theorem

Energy Conservation:

$$E = \sum_{n=-\infty}^{\infty} |x[n]|^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} |X(e^{j\omega})|^2 d\omega$$

Interpretation:

- Total energy in time domain = Total energy in frequency domain
- $|X(e^{j\omega})|^2$ is the **energy density spectrum**
- Shows how energy is distributed across frequencies

General Form:

$$\sum_{n=-\infty}^{\infty} x[n] y^*[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega}) Y^*(e^{j\omega}) d\omega$$

Application: Signal power calculation, filter energy analysis

The Convolution Theorem

ECEN 463/863
Maxx Seminario

Time Domain Convolution ↔ Frequency Domain Multiplication:

y[n] = x[n] * h[n] ↔ F Y(e^{j\omega}) = X(e^{j\omega})H(e^{j\omega})

Why This Works:

- Complex exponentials are eigenfunctions of LTI systems
- Input e^{j\omega n} produces output H(e^{j\omega})e^{j\omega n}
- By superposition, each frequency component is scaled by H(e^{j\omega})

Practical Impact:

- Convolution (complex operation) → Multiplication (simple)
- Foundation for frequency-domain filtering
- Basis for fast convolution algorithms

The Modulation (Windowing) Theorem

ECEN 463/863
Maxx Seminario

Time Domain Multiplication ↔ Frequency Domain Convolution:

y[n] = x[n]w[n] ↔ F Y(e^{j\omega}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\theta})W(e^{j(\omega-\theta)})d\theta

Key Points:

- This is **periodic convolution** (circular convolution)
- Integration over one period only
- Dual to the convolution theorem

Applications:

- **Windowing**: Multiplying by window function in time domain
- **Modulation**: Shifting spectrum by multiplication with e^{j\omega_0 n}
- **Spectral analysis**: Effect of finite-length observation

Example: Using Tables and Theorems

ECEN 463/863
Maxx Seminario

Problem: Find DTFT of x[n] = a^n u[n - 5]

Example: Inverse Transform Using Partial Fractions

ECEN 463/863
Maxx Seminario

Given: X(e^{j\omega}) = \frac{1}{(1 - ae^{-j\omega})(1 - be^{-j\omega})}

Example: Highpass Filter Design

ECEN 463/863
Maxx Seminario

Desired Frequency Response:

$$H(e^{j\omega}) = \begin{cases} e^{-j\omega n_d}, & \omega_c < |\omega| < \pi \\ 0, & |\omega| < \omega_c \end{cases}$$

Approach: Express as $H(e^{j\omega}) = e^{-j\omega n_d}(1 - H_{lp}(e^{j\omega}))$

Examples

Example: Solving Difference Equations

ECEN 463/863
Maxx Seminario

System: $y[n] - \frac{1}{2}y[n-1] = x[n] - \frac{1}{4}x[n-1]$
Finding Frequency Response:

- 1 Apply DTFT to both sides:
- 2 Factor and solve:
- 3 Partial fractions:
- 4 Inverse Transform:

Examples

Summary of DTFT

ECEN 463/863
Maxx Seminario

Key Concepts:

- 1 DTFT Pair:
 - Analysis: $X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$
 - Synthesis: $x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega})e^{j\omega n}d\omega$
- 2 Convergence:
 - Absolute summability \rightarrow uniform convergence
 - Square summability \rightarrow mean-square convergence
- 3 Symmetry Properties:
 - Real sequences have conjugate-symmetric DTFTs
 - Even/odd decomposition in both domains
- 4 Transform Theorems:
 - Convolution \leftrightarrow Multiplication
 - Time shift \leftrightarrow Linear phase
 - Differentiation, modulation, Parseval's theorem

Summary

Important Transform Pairs

ECEN 463/863
Maxx Seminario

| Sequence | DTFT |
|----------------------------------|--|
| $\delta[n]$ | 1 |
| $\delta[n - n_0]$ | $e^{-j\omega n_0}$ |
| $a^n u[n], a < 1$ | $\frac{1}{1 - ae^{-j\omega}}$ |
| $e^{j\omega_0 n}$ | $2\pi \sum_r \delta(\omega - \omega_0 + 2\pi r)$ |
| $\cos(\omega_0 n)$ | $\pi \sum_r [\delta(\omega - \omega_0 + 2\pi r) + \delta(\omega + \omega_0 + 2\pi r)]$ |
| $\frac{\sin(\omega_c n)}{\pi n}$ | Ideal lowpass filter |
| $u[n]$ | $\frac{1}{1 - e^{-j\omega}} + \pi \sum_r \delta(\omega + 2\pi r)$ |

Remember: DTFT is periodic with period 2π , so we only need to consider $\omega \in [-\pi, \pi]$

Summary