BIOMEDICAL LIMITED, ILORIN, No 1 Ohimege Road, Industrial Estate, Ilorin Kwara State, PMB 1499.	RAW MATERIAL ANALYTICALMETHODS	STP №	: BML/QA/STP/RMA/007/F02
		Material Code №	: RMA007
		Effective Date	: 4th Jan., 2024
		Supersedes	:
		Review Date	: 4th Dec., 2026
	FLUCONAZOLE	Revision No	: 01
		Ref. Book	BP 2023/IN-HOUSE
		Page	: 1 Of 11
Storage Conditions:	**Store in a well-closed container, protected from light**	**Department**	**: Quality**
Retest Frequency:	**12 months from date of release**	**Section**	**Quality Control**

ANALYTICAL METHODS

1 **DESCRIPTION**

1.1 Physically examine a representative sample of the material for Appearance and other characteristics: it is **White or almost white, hygroscopic, crystalline powder. Slightly soluble in water, freely soluble in methanol, soluble in acetone. It shows polymorphism**

2 **IDENTIFICATION**

2.1 **Light Absorption**

2.1.1 Accurately weigh and transfer about 20mg of the sample into a 100ml volumetric flask.

2.1.2 Add 60ml of distilled water, mechanically shake/sonicate till dissolution is complete

2.2.3 dilute to the mark with distilled water and mix.

2.1.4 Using a suitable spectrophotometer and 1-cm cells, measure the absorbances between the range of 230nm and 300nm using distilled water as blank **. The solution shows an absorption maxim at 261nm and 267 and a minimum at 264nm.**

2.2 **Melting point**

2.2.1 Determine the melting point using the capillary method:

3 **APPEARANCE OF SOLUTION**

3.1 Accurately weigh and transfer about 1g of the sample into a 20ml volumetric flask.

3.2 Add 15ml of methanol, mechanically shake/sonicate till dissolution is complete

3.3 Dilute to the mark with methanol and mix: **The** solution is clear and colourless .

4 **HEAVY METALS**

4.1 <u>Lead stock solution(100ppm)</u>

4.1.1 Accurately weigh and transfer about 160mg of lead into 250ml beaker.

4.1.2 Add 100ml of distilled water, 1ml of nitric acid and mix till dissolution is complete.

4.1.3 Rinse the 250ml beaker with several quantities of distilled water into the 1-litre volumetric

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		

flask.

4.1.4 Dilute with distilled water to 1000ml and mix. Prepare and store this solution in glass containers free from soluble lead salts.

4.2 Lead stock solution (10ppm)

4.2.1 Pipette 1.0ml of the 100ppm Lead standard solution into a 1000ml volumetric flask, dilute to the mark with methanol and mix.

4.3 Buffer Solution pH3.5

4.3.1 In a 250ml beaker, dissolve 25.0g of ammonium acetate in 25ml of water and add 38.0ml of 7M hydrochloric acid.

4.3.2 Adjust the pH if necessary, with 2M hydrochloric acid or 6M Ammonium hydroxide and dilute to 100.0ml with water.

4.4 Thioacetamide solution (4%)

4.4.1 Dissolve 4g of thioacetamide in a 100ml volumetric flask, make up with distilled water

4.42 Thioacetamide Reagent

To 2ml of thioacetamide solution in a 100ml volumetric flask, add 7.5ml of water, 37.5ml of M sodium hydroxide and 50ml of glycerol (85%). Heat in a water-bath for 20 seconds. Prepare immediately before use.

4.5 Sample Preparation

4.5.1 Dissolve 2.0 g in a mixture of 15 volumes of *water R* and 85 volumes of *methanol R*

4.5.2 dilute to 20.0 mL with the same mixture of solvents

4.5.3 To 12ml of the solution thus obtained (in a test tube), add 2ml of buffer solution pH3.5, 1.2ml of thioacetamide reagent and immediately mix.

4.6 Standard Preparation

4.6.1 Transfer 10 ml of 1ppm lead standard solution and 2 ml of the sample solution into a suitable test tube.

4.6.2 To the solution thus obtained, add 2ml of buffer solution pH3.5, 1.2ml of thioacetamide reagent and immediately mix.

4.7 Blank Preparation

4.7.1 A mixture of 10 mL of the solvent used for the substance to be examined and 2 mL of the prescribed solution of the substance to be examined in an organic solvent.

4.7.2 To the solution thus obtained, add 2ml of buffer solution pH3.5, 1.2ml of thioacetamide reagent and immediately mix.

4.8 Observation

4.8.1 Compared to the blank, after 2 minutes any brown colour in the solution obtained with the sample is not more intense than in the standard solution

5 Loss On Drying

5.1 Weigh two labelled petri dishes (W1) that has previously been dried at 105°C and cooled

5.2 Accurately weigh about 1g onto a petri-dish, ensuring they are evenly spread, then determine their weight together (W2).

5.3 Transfer into an Oven operating at 105°C.

5.4 Dry for 3 hours.

5.5 Transfer into a desiccator to cool to room temperature.

5.6 Weigh the petri-dish with its content (W3).

Calculation:

Calculate the moisture content as follows:

$$\% \text{ Moisture} = \frac{(W2 - W3) \times 100}{W2 - W1}$$

Where,

W1 = weight of empty petri-dish, g

W2 = weight of sample and petri-dish, g.

W3 = weight of petri-dish and the dried sample, g

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		

6 SULPHATED ASH

6.1 Weigh two crucibles (W1) that has previously been ignited and cooled.

6.2 Accurately weigh about 1g onto each crucible, ensuring they are evenly spread, then determine their weight together (W2).

6.3 Heat, gently at first on a Bunsen flame, until the sample is thoroughly charred.

6.4 Cool, then moisten the residue with 1mL of sulphuric acid, heat gently until white fumes are no longer evolved, and ignite at 800 ± 25°C until the carbon is consumed (about 2 hours).

6.5 Cool in a desiccator, weigh, and calculate the % ROI, if the %ROI exceed 0.1%, again moisten the residue with 1ml of sulphuric acid, heat and ignite as before.

6.6 Continue until constant weight is attained (until two successive weighing does not differ by more than 0.5mg).

6.7 Weigh the crucibles with its content (W3).

6.8 Calculation:

$$\% \text{ Sulphated Ash} = \frac{(W3 - W1) \times 100}{W2 - W1}$$

Where,

W1 = weight of empty crucibles, g

W2 = weight of sample and crucibles, g.

W3 = weight of crucibles and the dried sample, g

7 **RELATED SUBSTANCES** (Liquid chromatography)

7.1 Test solution Dissolve 0.100 g of the substance to be examined in the mobile phase, sonicate if necessary, and dilute to 10.0 mL with the mobile phase

7.2 Reference solution (a) Dilute 5.0 mL of the test solution to 100.0 mL with the mobile phase. Dilute 1.0 mL of this solution to 10.0 mL with the mobile phase

7.3 Reference solution (c) Dissolve 3.0 mg of fluconazole impurity B in the mobile phase, sonicate if necessary and, dilute to 100.0 mL with the mobile phase.

7.4 Reference solution (d) Dissolve 2.0 mg of fluconazole impurity C in the mobile phase and dilute to 20.0 mL with the mobile phase. To 1.0 mL of this solution add 1.0 mL of the test solution and dilute to10.0 mL with the mobile phase.

7.5 column size: l = 0.15 m, Ø = 4.6 mm;
stationary phase: octadecylsilyl silica gel for chromatography R1 (5 µm);
temperature: 40 °C
Mobile phase acetonitrile , 0.63 g/L solution of ammonium formate R (14:86 V/V).

Flow rate 1.0 mL/min.
Detection Spectrophotometer at 260 nm.
Injection 20 µL.
Run time 3.5 times the retention time of fluconazole

7.6 Identification of impurities
Use the chromatogram supplied with fluconazole for peak identification and the chromatogram obtained with reference solution (b) to identify the peak due to impurity A; use the chromatogram obtained with reference solution (c) to identify the peak due to impurity B and the chromatogram obtained with reference solution (d) to identify the peak due to impurity C

7.7 *Relative retention*
With reference to fluconazole (retention time = about 11 min): impurity B = about 0.4; impurity A = about 0.5; impurity C = about 0.8.

7.8 System suitability Reference solution (d):
— resolution: minimum 3.0 between the peaks due to impurity C and fluconazole.

7.9 Limits:
7.9.1 — impurity A: not more than 0.8 times the area of the principal peak in the chromatogram obtained with reference solution (a) (0.4 per cent);
7.9.2 — impurity B: not more than the area of the principal peak in the chromatogram obtained with reference solution (c) (0.3 per cent);

7.9.3 — impurity C: not more than the area of the corresponding peak in the chromatogram obtained with reference solution (d) (0.1 per cent);

7.9.4 — unspecified impurities: for each impurity, not more than 0.2 times the area of the principal peak in the chromatogram obtained with reference solution (a) (0.10 per cent);

7.9.5 — total: not more than 1.2 times the area of the principal peak in the chromatogram obtained with reference solution (a) (0.6 per cent);

8 **ASSAY**

8.1 **Reagents**

8.1.1 0.1M perchloric acid; prepared

8.1.2 0.1% crystal violet or naphthobenzein in glacial acetic acid
Accurately weigh and transfer about 0.1g to 100ml volumetric flask, dilute to the mark with glacial acetic acid and mix

8.2 **SAMPLE PREPARATION**

8.2.1 Accurately weigh about 125mg each into two separate, dry 250ml conical flasks.

8.2.2 Add 60 ml of glacial acetic acid to each flask, swirling from time to time till dissolution is complete.

8.2.3 Add two drops of 0.1% of crystal violet in glacial acetic acid and titrate with 0.1M perchloric acid in glacial acetic acid to an emerald green end-point.

8.2.4 Calculate the Fluconazole content as follows:

9 Microbiological Tests

9.1 <u>Limulus Amebocyte Lysate LAL Test</u>

9.1.1 Open the seals of the test tubes of lyophilized LAL Reagent

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		

9.1.2 Add 0.2ml LAL Reagent water, endotoxin standards or test sample into each tubes.

9.1.3 Swirl gently but thoroughly to rehydrate the LAL reagent. Do not shake or vortex as content will foam.

NOTE: cover the reaction tubes with pyrogen free aluminium foil

9.1.4 Incubate the reaction mixture immediately at 37°c for 60minutes. Avoiding vibration during incubation.

9.1.5 If large numbers of samples are tested in parallel, the tests should be batched and started at intervals that permit the reading of each within the time limit.

9.1.6 Sample to be tested should be stored under conditions that bacteriological activities are controlled

9.1.7 Read reaction tubes one by one.

9.1.8 Invert the tube in one smooth motion.

9.1.9 A positive result is noticed by the formation of a gel which does not collapse when the tube is inverted. A negative result is characterized by the absence of solid clot after inversion.

8.2

Testing Methodology of Bacterial Endotoxin (LAL)

9.2.1 Each specimen shall be tested in duplicate

9.2.2 NMC: Take 100µl(0.1ml) LRW

9.2.3 Take 50µl (0.05ml) LRW and 50µl (0.05ml) test dilution.

9.2.4 Take 50µl(0.05ml) test dilution and 50µl(0.05ml) CSE of 4λ

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		

9.2.5 Add the LAL reagents as mentioned below in the table 1

9.2.6 Mix gently and incubate the tubes in heating block at 37^0C ± 1 for 60min.

9.2.7 After incubation, remove the tubes gently from the heating block and slowly invert through 180^0C and note the observations.

9.2.8 If a firm gel has formed, record the results as positive.

9.2.9 A result is negative, if an intact gel is not formed.

9.2.10 Record the results and other details as per format no. "Analytical Test Data Sheet for Bacterial Endotoxin (LAL) Test" in case of sterile raw material.

8.3 **Pyrogen**

9.3.1 Perform the test in a separate area designated for Pyrogen testing and under environmental conditions similar to those under which the rabbits are housed.

9.3.2 Withhold all food from the rabbits used during the period of the test

9.3.3 Draw out the required number of rabbits from their living room according to the number of tests planned. Always select more rabbits than actually required to counter any exclusion of animals in the initial measuring.

9.3.4 All the rabbits which are to be used for the test should be more than 1.5kg weight, should be healthy and must not have been used for any test in previous three(3) days and in previous three (3)weeks if any test failed using that rabbit.

9.3.5 Put the selected rabbits to testing cages. Take the initial temperature of each rabbit by inserting the thermometer to in the anus rectum of the rabbit to a depth of about 7.5cm.

9.3.6 If the initial temperature is less than 38.8^0c or more than 39.8^ocexclude the animal from the test.

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		

9.3.7 Ensure all the needles and syringe to be used are new

9.3.8 Arrange the rabbits into groups of three taking into consideration that the initial temperature of the rabbits in a group should not differ by more than 1^0c.

9.3.9 Take another two reading of temperature in 30 min intervals between 1 and 3 hours subsequent to the injection.

9.3.10 Any rabbits showing more than 0.2^0c temperature difference between two successive readings in the last three readings can be excluded from the test.

9.3.11 Calculate the initial temperatures of the animals by taking the average of the temperature readings taken in the last 40 minutes.

9.3.12 Inject the solution intended to be tested to the marginal veins of the rabbits of a group slowly but not exceeding 10 minutes after the start of administration. The quantity of test solution depends upon the body weight of the rabbits and the test solution.

9.3.13 If the substance to be tested is not in aqueous sterilized solution form, consult the official monograph regarding the preparation of simulating solution and the volume of the preparation to be injected.

9.3.14 After injecting the solution, Record the temperature at 30 minutes intervals between 1 and 3 hours subsequent to the injection.

9.3.15 The test involves measuring the rise in temperature of rabbits following the intravenous injection of a test solution

8.4 MLT

9.4.1 Weigh 1g or 1ml of the raw material sample and dissolve aseptically in 9ml sterile soybean casein digest medium prepared as per SOP for media preparation (BML-QDG-MC-011).

9.4.2 Determine the Total Aerobic Microbial count (TAMC) and Total Yeast and Molds count (TYMC) by using pour plate method.

9.4.3 Testing to be performed under LAF of specified area

9.4.4 Arrange all the required material on the LAF before starting the test.

9.4.5 Aseptically Pipette 1ml of the solution from above in a sterile Petri dishes

9.4.6 Add approximately 15-20ml of sterile Soyabean Casein Digest Agar prepared as per SOP(BML-QDG-MC-011), (cooled approximately up to 45°C) in two petri plates for TAMC

9.4.7 Add approximately 15-20ml of sterile Saboraud Dextrose Agar prepared as per SOP (cooled approximately up to 45°C) in two petri- plates for TYMC

9.4.8 Gently swirl the contents inside the petri-dishes for proper mixing of the sample, allow medium to solidify, taking the precaution of not spilling the media from the plate

9.4.9 Simultaneously perform the negative control by plating 1 ml of diluents used instead of the sample for each medium. Add 15-20ml of the media previously maintained not more than 45°c and allow for solidification.

9.4.10 Incubate the plates at 30°C to 35°C for 3 - 5 days for soybean Casein Digest Agar for TAMC and plates of Saboraud Dextrose Agar at 20°C to 25°C for 5 days in an inverted

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		

position.

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	**QUALITY CONTROL SENIOR ANALYST**		
REVIEWED BY:	**QUALITY CONTROL HEAD**		
APPROVED BY:	**QUALITY ASSURANCE MANAGER**		