Magflex Suspension

Composition:

Each 5 ml contains:

Magnesium Trisilicate BP…… 250mg

Light Magnesium Carbonate BP…… 250mg

Sodium Bicarbonate BP…… 250mg

Indication:

Used for relief of Gastric ulcer, heartburn, indigestion, acid reflux, hyperacidity, and bloating.

Mechanism of Action:

Magnesium salts neutralize acid slowly and protect stomach lining. Sodium bicarbonate neutralizes acid quickly for fast relief.

Dosage:

Age	Dose
Adults ≥ 12 years	10 ml – 20 ml
Children < 12 years	5 ml – 10 ml
To be taken 3-4 times daily after meals and at bedtime	

Contra-indication:

Magflex Suspension should not be used if you have severe kidney problems, because Magnesium can build up in the body and lead to weakness, low blood pressure, or heart problems. Do not use this product if you have metabolic alkalosis, severe dehydration, or severe stomach pain of unknown cause. People on a low-sodium diet, or those with conditions like high blood pressure, heart failure, kidney disease, or liver cirrhosis should avoid this medicine due to its sodium content. Avoid use if you are allergic to any of the ingredient.

Adverse effects:

Possible diarrhoea, bloating, belching, stomach cramps, or rare nausea.

Patient Advice & Precautions

1. Keep all medicine tightly closed after use and out of children's reach.
2. It should not be taken together with certain medicines such as antibiotics, iron, thyroid medication, or bisphosphonates.
3. At least, 1–2 hours interval should be exercised between Magflex and uptake of other drugs.
4. Pregnant or breast-feeding women, elderly patients, and those with very low stomach acid should use this medicine only on the advice of a healthcare professional. If symptoms persist, worsen, or new symptoms appear, stop using the product and consult your doctor.

Storage:

Store at temperature not more than 30°C, protected from sunlight and moisture.

Presentation:

Available in 100 ml and 200 ml PET bottle.

SHAKE WELL BEFORE USE.

Manufactured by:

Biomedical Limited
1, Ohimege Road, Gaa-Imam, Industrial Estate,
Ilorin, Kwara State, Nigeria.

Magflex Suspension

Composition:

Chaque 5 ml contient:

Trisilicate de Magnésium BP..... 250 mg

Carbonate de Magnésium léger BP ..... 250 mg

Bicarbonate de Sodium BP ..... 250 mg

Indication:

Utilisé pour le soulagement de l'ulcère gastrique, des brûlures d'estomac, de l'indigestion, du reflux acide, de l'hyperacidité et des ballonnements.

Mécanisme d'action:

Les sels de magnésium neutralisent lentement l'acidité et protègent la muqueuse de l'estomac. Le bicarbonate de sodium neutralise l'acidité rapidement, offrant un soulagement rapide.

Dosage:

Age	Dose
Adultes ≥ 12 ans	10 ml à 20 ml
Enfants < 12 ans	5 ml à 10 ml
À prendre 3 à 4 fois par jour, après les repas et au coucher.	

Contre-indications:

Magflex Suspension ne doit pas être utilisé en cas de problèmes rénaux sévères, car le Magnésium peut s'accumuler dans l'organisme et entraîner faiblesse, hypotension ou problèmes cardiaques. Ne pas utiliser en cas d'alcalose métabolique, de déshydratation sévère ou de douleur abdominale intense d'origine inconnue. Les personnes suivant un régime pauvre en sodium, ou souffrant d'hypertension, d'insuffisance cardiaque, de maladie rénale ou de cirrhose du foie doivent éviter ce médicament en raison de sa teneur en sodium.
Éviter l'utilisation en cas d'allergie à l'un des composants.

Effets indésirables:

Possibilité de diarrhée, ballonnements, éructations, crampes abdominales ou, rarement, nausées.

Avertissements & Précautions:

1. Garder le médicament bien fermé après usage et hors de portée des enfants.
2. Ne pas prendre en même temps que certains médicaments tels que les antibiotiques, le fer, les médicaments thyroïdiens ou les bisphosphonates.
3. Laisser au moins 1 à 2 heures entre Magflex et les autres médicaments.
4. Les femmes enceintes ou allaitantes, les patients âgés et ceux ayant une très faible acidité gastrique doivent utiliser ce médicament uniquement sur les conseils d'un professionnel de santé. Si les symptômes persistent, s'aggravent ou si de nouveaux symptômes apparaissent, arrêtez d'utiliser le produit et consultez votre médecin.

Espace de rangement:

Conserver une température ne depassant pas 30°C.Protéger de la lumière du soleil et de l'humidité.

Présentation:

Disponible en flacons PET de 100 ml et 200 ml**.**

BIEN AGITER AVANT USAGE.

Fabriqué par:

Biomedical Limited
1, Route d'Ohimege, Zone Industrielle de Gaa-Imam,
Ilorin, Kwara State, Nigeria.