BIOMEDICAL LIMITED, ILORIN, No 1 Ohimege Road, Industrial Estate, Ilorin Kwara State, PMB 1499.	RAW MATERIAL ANALYTICAL METHODS	STP №	: BML/QA/STP/RMA/003/F02
		Material Code №	: RMA003
		Effective Date	: 4th Jan., 2024
		Supersedes	:
		Review Date	: 4th Dec., 2026
	POTASSIUM CHLORIDE	Revision No	: 01
		Ref. Book	BP 2023/IN-HOUSE
		Page	: 1 Of 20
Storage Conditions:	**Store in a well-closed container, protected from light**	**Department**	**: Quality**
Retest Frequency:	**12 months from date of release**	**Section**	**Quality Control**

ANALYTICAL METHODS

1 DESCRIPTION

1.1 Physically examine a representative sample of the material for Appearance and other characteristics: it is **A** White or almost white, crystalline powder or colourless crystals Freely soluble in water, practically insoluble anhydrous ethanol**.**

2 IDENTIFICATION

2.1 Potassium test

Sodium Sulphide Solution

2.1.1 Weigh about 12g of sodium sulphide into a 250ml beaker.
Add 60ml of 75%w/v glycerol solution, mechanically shake/sonicate till dissolution is complete.
Transfer onto steam-bath and heat, with mixing, till dissolution is complete.
Allow cooling, transfer to a 100ml volumetric flask and dilute to the mark with 75% glycerol and mix.

2.1.2 15% w/v solution of sodium carbonate; Accurately weigh and transfer about 15g sodium carbonate to 100ml volumetric flask, dilute to the mark with distilled water and mix

2.1.3 Procedure
2.1.4 Accurately weigh and transfer about 10g of the sample into a 100ml volumetric flask.
2.1.5 Add 60ml of distilled water, mechanically shake till dissolution is complete.
2.1.6 Dilute to the mark with distilled water and mix
2.1.7 Pipette 2ml of the solution into a test tube.
2.1.8 Add 1ml of 15% w/v solution of sodium carbonate and heat to boiling; No **precipitate is produced.**

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		

2.1.9 Add while still hot, 0.05ml of 12%w/v sodium sulphide in 75% glycerol and mix: No precipitate is produced.

2.1.10 Cool in ice and add 2 ml of a 15% w/v solution of (+)-tartaric acid and allow to stand: **A white, crystalline precipitate is produced.**

2.3 Chloride test

2.3.1 Reagents

2M nitric acid

Pipette 12.6ml of concentrated Nitric acid to 100ml volumetric flask containing 60ml distilled water, dilute to the mark with distilled water and mix

4.25% w/v silver nitrate

Accurately weigh and transfer about 4.25g to 100ml volumetric flask, dilute to the mark with distilled water and mix

10M ammonia

Dilute about 41g of con ammonia to 100ml volumetric flask, dilute to the mark with distilled water and mix

2.3.2 Sample preparation

2.3.3 Dissolve 1g of sample in 2 ml of distilled water.

2.3.4 Acidify with 2M nitric acid, add 0.4 ml of 4.25% w/v silver nitrate solution.

2.3.5 Shake and allow to stand; **a curdy, white precipitate is produced**

2.3.6 Filter and wash the precipitate with three 1-ml quantities of water.

2.3.7 Carry out this operation rapidly in subdued light, disregarding the fact that the supernatant solution may not become perfectly clear: Suspend the precipitate in 2 ml of water and add 1.5ml of 10M ammonia; **the precipitate dissolves easily with the possible exception of a few large particles which dissolve slowly.**

3 APPEARANCE OF SOLUTION

3.1 Accurately weigh and transfer about 20g of sample into a 100ml volumetric flask.

3.2 Add 60ml of freshly distilled and cooled water; mechanically shake/sonicate till dissolution is complete.

3.3 Dilute to the mark with distilled water and mix.**Solution S is clear and colourless**

3.4 Observe the solution against bright northerly light: Dry for 2 hours.

4 ACIDITY OR ALKALINITY ACIDITY OR ALKALINITY.

4.1 **Reagents**

4.1.1 0.1%w/v bromothymol blue indicator.ethanol; ; Accurately weigh and transfer about 0.1g bromothymol blue to 100ml volumetric flask, dilute to the mark with ethanol and mix

4.1.2 0.01M hydrochloric acid; dilute 10ml of 0.01M hydrochloric acid to 100ml with distilled water

4.1.3 0.01M sodium hydroxide ; dilute 10ml of 0.1M sodium hydroxide to 100ml with distilled water

4.2 **Sample preparation**

4.2.1 Accurately weigh and transfer about 20g of sample into a 100ml volumetric flask.

4.2.2 Add 60ml of distilled water and mechanically shake/sonicate till dissolution is complete. Dilute to the mark with same solvent and mix

4.2.3 Pipette 20.0ml of this solution into a 100ml conical flask, add 0.1ml of 0.1%w/v bromothymol blue indicator.

4.2.4 Titrate with 0.01M hydrochloric acid or 0.01M sodium hydroxide solution to a change in the colour of the indicator: **Not more than 0.5 ml of 0.01M hydrochloric acid or 0.01M sodium hydroxide is required to change the colour of the indicator.**

5 BROMIDES

5.1 Phenol Red Solution

5.1.1 Accurately weigh and transfer about 0.105g of phenol red into 100ml volumetric flask

5.1.2 Add 3.0ml of 0.1M sodium hydroxide, 25ml of 96% ethanol and mechanically shake/sonicate till dissolution is complete.

5.1.3 Dilute to the mark with distilled and mix.

5.2 Standard Preparation

5.2.1 Accurately weigh and transfer about 15mg of potassium bromide into a 1000ml volumetric flask.

5.2.2 Add 600ml of distilled water and mechanically shake/sonicate till dissolution is complete.

5.2.3 Dilute to the mark with distilled and mix.

5.3 Sample Preparation

5.3.1 Accurately weigh and transfer about 20g of sample into a 100ml volumetric flask.

5.3.2 Add 60ml of distilled water and mechanically shake/sonicate till dissolution is complete.

5.3.3 Dilute to the mark with distilled water and mix.

Procedure

5.3.4 Pipette 1.0ml each of the sample and standard preparations into separate 10ml volumetric flasks.

5.3.5 Add to each 10ml flask, 6.0ml of distilled water, 2.0ml of phenol red solution, 1.0ml of 0.01%w/v chloramines solution and mix immediately.

5.3.6 After exactly 2 minutes, add to each 10ml flask, 0.15 ml of 0.1M sodium thiosulphate and mix.

5.3.7 Using a suitable spectrophotometer and 1-cm cells, measure the absorbance of the sample and standard preparations at 590nm against distilled water: **The absorbance of the sample solution is not greater than that of a standard preparation (50 ppm).**

6 IODIDES

6.1 Starch Solution, Iodide-free

6.1.1 Weigh about 1g of soluble starch into 250ml beaker

6.1.2 Add 5ml of distilled water and and triturate with glass rod.

6.1.3 With continuous stirring, 100ml of boiling water.

6.1.4 After the addition of the boiling water, continue to stir for further 5 minutes.

6.2 Sodium Nitrite-Sulphuric Mixture **(Prepare 10 mniutes before use!!!)**

6.2.1 Mix 1.0ml of 1.5%w/v sodium nitrite, 2ml of 0.5M sulphuric acid, 25mlof iodide-free starch solution and 25ml of distilled water

6.3 Procedure

6.3.1 Accurately weigh about 5g of sample into a 250ml beaker

6.3.2 Moisten the sample by the dropwisely adding the freshly prepared sodium nitrite-sulphuric mixture

6.3.3 Allow 5 minutes and examine the sample in daylight: **The substance shows no blue colour.**

7 SULPHATES

7.1 1000ppm SULPHATE STANDARD PREPARATION

7.1.1 Accurately weigh and transfer about 0.181g of analytic grade potassium sulphate into a 100ml volumetric flask.

7.1,2 Add 60ml of 30% v/v ethanol in water and sonicate till dissolution is complete.

7.1.3 Dilute to the mark with 30% v/v ethanol in water and mix “(1000ppm sulphate standard stock solution).

7.1.4 Pipette 20.0ml of this solution into another 100ml volumetric flask, dilute to the mark with 30% v/v ethanol in water and mix (200ppm sulphate working standard).

7.2 SAMPLE PREPARATION

7.2.1 Accurately weigh and transfer about 10g of sample into a 100ml volumetric flask.

7.2.2 Add 60ml of distilled water and mechanically shake/sonicate till dissolution is complete.

7.2.3 Dilute to the mark with same solvent and mix.

7.3 Procedure

7.3.1 Pipette 15.0ml each of this solution and 1ml of acetic acid into 25ml volumetric flasks.

7.3.2 To another flask, 13.0ml of distilled water, 2ml of 150ppm sulphate standard solution, and 1ml of acetic acid into 25ml volumetric flasks.

7.3.3 To each flask, add 1ml of 25% w/v solution of barium chloride solution and 1ml of 2M nitric acid.

7.3.4 Dilute to the mark with 30% v/v ethanol in water and mix.

7.3.5 Shake each flask and allow standing for 5 minutes: **Any opalescence produced is not more intense than that of Sulphate standard (200ppm SO_4).**

8 ALUMINIUM

8.1 Acetate Buffer pH 6.0

8.1.1 Weigh about 40g of ammonium acetate into a 250ml beaker

8.1.2 Add 100ml of distilled water, 1.6ml of glacial acetic acid and mix till dissolution is complete.

8.1.3 Adjust the pH to 6.0, if necessary, using 10M ammonia or 5M acetic acid

8.1.4 Dilute to the 200ml- mark with distilled water and mix.

8.2 200ppm Aluminium Standard Solution

8.2.1 Weigh about 0.352g of [Aluminium Potassium Sulphate (AlK(SO4)2,12H2O) into a 100ml volumetric flask.

8.2.2 Add 30ml of distilled water, 10 ml of 1M sulphuric acid and mechanically shake/sonicate till dissolution is complete.

8.2.3 Dilute to the 100ml- mark with distilled water and mix. **Store this solution for not more than 3 months.**

8.3 2ppm Aluminium Standard Solution **(Prepare Immediately Before Use!)**

8.3.1 Pipette 1.0ml of 200ppm aluminium solution into 100ml volumetric flask.

8.3.2 Dilute to the 100ml- mark with distilled water and mix. **Store this solution for not more than 3 months.**

8.4 0.5% w/v Solution Of 8-Hydroxyquinoline

8.4.1 Weigh and transfer about 1g of 8-hydroxyquinoline into a dry 200ml volumetric flask.

8.4.2 Add 100ml of chloroform, mechanically shake/sonicate till dissolution is complete.

8.4.3 Dilute to the 200ml- mark with chloroform and mix.

8.5 SAMPLE PREPARATION

8.5.1 Accurately weigh and transfer about 4g of sample into a 100ml volumetric flask.

8.5.2 Add 60ml of distilled water and mechanically shake/sonicate till dissolution is complete.

8.5.3 Add 10ml of acetate buffer pH 6.0, dilute to the mark with distilled water and mix.

8.6 Procedure

8.6.1 Into three 250 ml Separating funnels, pipette and add in the following order:

Reagent s	Sample	Standard	Blank
Sample Solution	100.0ml	-	-
2ppm Aluminium Standard Solution	-	2.0ml	-
Water	-	98.0ml	100.0ml
Acetate Buffer pH 6.0	10.0ml	10.0ml	10.0ml

8.6.2 Separately extract each of the sample, standard and blank preparations in the above table with successive quantities of 20, 20 and 10 ml of a 0.5% w/v solution of 8-hydroxyquinoline in chloroform, filtering the extracts through chloroform-wet Whatman #1 filter paper into separate, 'sample', 'standard', 'blank' labeled, dry 50ml volumetric flasks

8.6.3 Using a suitable spectrophotometer and 1-cm cells, measure the absorbance of the test solution and that of the standard solution against the prepared blank at wavelength of 392 nm and 518 nm: **The absorbance of the test solution is not greater than that of the standard solution.**

9 **BARIUM**

9.1 Pipette 5.0ml of solution S and 5ml of distilled water into test tube and mix:

9.2 Add 1ml of 1M sulphuric acid and mix

9.3 After at least 15 minutes, any opalescence in the sample solution is not more intense than that in a mixture of 5ml of solution S and 6ml of distilled water

10 IRON

10.1 <u>20ppm Iron Standard Stock</u>

10.1.1 Accurately weigh and transfer about 140.4mg of Ferrous Ammonium Sulphatehexahydrate
$[Fe(NH_4)_2(SO_4)_6H_2O]$ into a one-litre volumetric flask.

10.1.2 Add 300ml of distilled water, 25ml of concentrated sulphuric acid, mechanically shake/sonicate till dissolution is complete.

10.2 <u>Sample Preparation</u>

10.2.1 Accurately weigh and transfer about 10g of sample into a 100ml volumetric flask.

10.2.2 Add 60ml of distilled water and mechanically shake/sonicate till dissolution is complete.

10.2.3 Dilute to the mark with same solvent and mix.

10.2.4 Procedure

10.2.5 Pipette 10.0ml each of the sample and standard preparations into separate 25ml volumetric flask.

10.2.6 Add 2ml of 20%w/v citric acid, 0.1ml of mercpatoacetic acid, mix and compare sample and standard preparations

10.2.7 Dilute to the mark with distilled water and mix.

10.2.8 Using a suitable measuring cylinder, transfer 20ml of each preparation into separate Nessler cylinders and compare the intensity of the colour developed: **Pink colouration ob solution is not more intense than the standard solution.**

11 MAGNESIUM AND ALKALI METALS

11.1 **Reagent**

11.1.1 0.01M disodium edetate prepared as per SOP NO QCC016-05

11.1.2 0.1M zinc sulphate prepared as per SOP NO QCC016-05

11.2 **Sample preparation**

11.2.1 Into a 500ml beaker, measure (with a cylinder), 200ml of distilled water.

11.2.2 Add 0.1 g of hydroxylamine hydrochloride, 10 ml of ammonia buffer pH 10.0, 1 ml of 0.1M zinc sulphate and about 15 mg of mordant black 11 triturate..

11.2.3 Heat to about 40° and titrate with 0.01M disodium edetate VS until the violet colour changes to a full blue.

11.2.4 Accurately weigh and transfer about 10g of sample into the beaker and mix until dissolution is complete.

11.2.5 If the colour of the solution changes to violet, titrate with 0.01M disodium edetate VS until the full blue colour is again produced.

11.2.6 The volume of 0.01M sodium edetate used does not exceed 2.5ml.

12 SODIUM

12.1 Potassium Antimonate V

12.1.1 Weigh about 2g of potassium antimonite V into a 250ml beaker.

12.1.2 Add 95ml of hot water, drop a bar magnet and mix (using a magnetic stirrer) till dissolution is complete

12.1.3 Cool rapidly to room temperature in a water-bath at 25°C

12.1.4 Add 50ml of 5%w/v solution of potassium hydroxide and 1ml of 2M sodium hydroxide and mix.

12.1.5 Allow standing for 24 hours, filter through Whatman #1 filter-paper into another 250ml beaker

12.1.5 Dilute to 150ml with distilled water and mix.

12.2 1000ppm Standard sodium Stock Solution

12.2.1 Accurately weigh and transfer about 308mg - 310mg f anhydrous sodium sulphate into a 100ml volumetric flask.

12.2.2 Add 60ml of distilled water and mechanically shake/sonicate till dissolution is complete.

12.2.3 Dilute to the mark with distilled water and mix.

12.3 Sample Preparation

12.3.1 Accurately weigh and transfer about 20g of sample into a 100ml volumetric flask.

12.3.2 Add 60ml of distilled water and mechanically shake/sonicate till dissolution is complete.

12.3.3 Dilute to the mark with distilled water and mix.

12.4 Procedure

12.4.1 Pipette 10ml each of sample and standard preparations into separate 50 ml test tube

12.4.2 Add 5ml of a potassium antimonite V and heat to boiling.

12.4.3 Allow cooling in ice and if necessary scratch the inside of the tube with a glass rod

12.4.4 Swirl the tube: **Any opalescence produced in the sample preparation is not more intense than in the reference solution (0.1%)**

13 **LOSS ON DRYING**

13.1 Weigh one labelled petri dish (W1) that has previously been dried at 100°C and cooled.

13.2 Accurately weigh about 1g onto a petri-dish, ensuring they are evenly spread, then determine their weight together (W2).

13.3 Transfer into an Oven operating at 105°C.

13.4 Dry for 3 hours.

13.5 Transfer into a desiccators to cool room temperature.

13.6 Weigh the petri-dish with its content (W3).

Calculate the moisture content as follows:

$$\% \text{ Moisture} = \frac{(W2 - W3) \times 100}{W2 - W1}$$

Where,

W1 = weight of empty petri-dish, g)

W2 = weight of sample and petri-dish, g.

W3 = weight of petri-dish and the dried sample, g

14 **HEAVY METALS**

14.1 Lead stock solution(100ppm)

14.1.1 Accurately weigh and transfer about 160mg of Lead into a 1-litre volumetric flask.

14.1.2 Add 200ml of distilled water, 1ml of concentrated nitric acid, cover and shake till dissolution is complete.

14.1.3 Dilute with distilled water to 1000ml and mix. Prepare and store this solution in glass containers free from soluble lead salts.

14.2 Lead stock solution (1ppm)

14.2.1 Pipette1.0ml of the 100ppm lead standard solution into a 100ml volumetric, dilute to the mark with distilled water and mix.

14.3 Buffer Solution pH3.5

14.3.1 In a 250ml beaker, dissolve 25.0g of ammonium acetate in 25ml of water and add 38.0ml of 7M hydrochloric acid.

14.3.2 Adjust the pH if necessary with 2M hydrochloric acid or 6M ammonium hydroxide and dilute to 100.0ml with water.

14.4 Thioacetamide solution(4%)

14.4.1 Dissolve 4g of thioacetamide in a 100ml volumetric flask, make up with distilled water

Thioacetamide Reagent

14.4.2 To 2ml of thioacetamide solution in a 100ml volumetric flask, add 7.5ml of water, 37.5ml of M sodium hydroxide and 50ml of glycerol (85%). Heat in a water-bath for 20 seconds. Prepare immediately before use.

14.5 Sample Preparation

14.5.1 Accurately weigh and transfer about 20g of sample into a 100ml volumetric flask.

14.5.2 Add 60ml of distilled water and mechanically shake/sonicate till dissolution is complete.

14.5.3 Dilute to the mark with same solvent and mix.

14.5.4 To 12ml of the solution thus obtained (in a test tube), add 2ml of buffer solution pH3.5, 1.2ml of thioacetamide reagent and immediately mix.

14.6 Standard Preparation

14.6.1 Transfer 10 ml of 1ppm *lead standard solution* and 2 ml of the sample solution into a suitable test tube.

14.6.2 To the solution thus obtained, add 2ml of buffer solution pH3.5, 1.2ml of thioacetamide reagent and immediately mix.

14.7 Blank Preparation

14.7.1 Prepare a blank using a mixture of 10ml of water and 2ml of sample solution.

14.7.2 To the solution thus obtained, add 2ml of buffer solution pH3.5, 1.2ml of thioacetamide reagent and immediately mix.

14.8 OBSERVATION

Compared to the blank, after 2 minutes any brown colour in the solution obtained with the sample is not more intense than in the standard solution

15 ASSAY

15.1 **Reagents**

15.1.1 5%w/v potassium chromate solution Accurately weigh and transfer about 5.0g potassium chromate to 100ml volumetric flask, dilute to the mark with distilled water and mix

0.1M Silver Nitrate

Sample preparation

15.1.2 Accurately weigh about 60mg each into two separate 250ml conical flasks.

15.1.3 Dilute to 50ml with water and sonicate/swirl till dissolution is complete.

16 **Microbiological Tests**

16.1 **<u>Limulus Amebocyte Lysate LAL Test</u>**

16.1.1 Open the seals of the test tubes of lyophilized LAL Reagents

16.1.2 Add 0.2ml LAL Reagent water, endotoxin standards or test sample into each tubes.

16.1.3 Swirl gently but thoroughly to rehydrate the LAL reagent. Do not shake or vortex as content will foam.

NOTE: cover the reaction tubes with pyrogen free aluminium foil

16.1.4 Incubate the reaction mixture immediately at 37°c for 60minutes. Avoiding vibration during incubation.

16.1.5 If large numbers of samples are tested in parallel, the tests should be batched and started at intervals that permit the reading of each within the time limit.

16.1.6 Sample to be tested should be stored under conditions that bacteriological activities are controlled

16.1.7 Read reaction tubes one by one.

16.1.8 Invert the tube in one smooth motion.

16.1.9 A positive result is noticed by the formation of a gel which does not collapse when the tube is inverted. A negative result is characterized by the absence of solid clot after inversion.

Testing Methodology of Bacterial Endotoxin (LAL)

16.2.1 Each specimen shall be tested in duplicate

16.2.2 NMC: Take 100µl(0.1ml) LRW

16.2.3 Take 50µl (0.05ml) LRW and 50µl (0.05ml) test dilution.

16.2.4 Take 50µl(0.05ml) test dilution and 50µl(0.05ml) CSE of 4λ

16.2.5 Add the LAL reagents as mentioned below in the table 1

16.2.6 Mix gently and incubate the tubes in heating block at 37^0C ± 1 for 60min.

16.2.7 After incubation, remove the tubes gently from the heating block and slowly invert through 180^0C and note the observations.

16.2.8 If a firm gel has formed, record the results as positive.

16.2.9 A result is negative, if an intact gel is not formed.

16.2.10 Record the results and other details as per format no. "Analytical Test Data Sheet for Bacterial Endotoxin (LAL) Test" in case of sterile raw material.

Pyrogen

16.3 16.3.1 Perform the test in a separate area designated for Pyrogen testing and under environmental conditions similar to those under which the rabbits are housed.

16.3.2 Withhold all food from the rabbits used during the period of the test

16.3.3 Draw out the required number of rabbits from their living room according to the number of tests planned. Always select more rabbits than actually required to counter any exclusion of animals in the initial measuring.

16.3.4 All the rabbits which are to be used for the test should be more than 1.5kg weight, should be healthy and must not have been used for any test in previous three(3) days and in previous three (3)weeks if any test failed using that rabbit.

16.3.5 Put the selected rabbits to testing cages. Take the initial temperature of each rabbit by inserting the thermometer to in the anus rectum of the rabbit to a depth of about 7.5cm.

16.3.6 If the initial temperature is less than 38.8^{0}c or more than 39.8^{o}cexclude the animal from the test.

16.3.7 Ensure all the needles and syringe to be used are new

16.3.8 Arrange the rabbits into groups of three taking into consideration that the initial temperature of the rabbits in a group should not differ by more than 1^{0}c.

16.3.9 Take another two reading of temperature in 30 min intervals between 1 and 3 hours subsequent to the injection.

16.3.10 Any rabbits showing more than 0.2^{0}c temperature difference between two successive readings in the last three readings can be excluded from the test.

16.3.11 Calculate the initial temperatures of the animals by taking the average of the temperature readings taken in the last 40 minutes.

16.3.12 Inject the solution intended to be tested to the marginal veins of the rabbits of a group slowly but not exceeding 10 minutes after the start

of administration. The quantity of test solution depends upon the body weight of the rabbits and the test solution.

16.3.13 If the substance to be tested is not in aqueous sterilized solution form, consult the official monograph regarding the preparation of simulating solution and the volume of the preparation to be injected.

16.3.14 After injecting the solution, Record the temperature at 30 minutes intervals between 1 and 3 hours subsequent to the injection.

16.3.15 The test involves measuring the rise in temperature of rabbits following the intravenous injection of a test solution.

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		

16.4 **MLT**

16.4.1 Weigh 1g or 1ml of the raw material sample and dissolve aseptically in 9ml sterile soybean casein digest medium prepared as per SOP for media preparation (BML-QDG-MC-011).

16.4.2 Determine the Total Aerobic Microbial count (TAMC) and Total Yeast and Molds count (TYMC) by using pour plate method.

16.4.3 Testing to be performed under LAF of specified area

16.4.4 Arrange all the required material on the LAF before starting the test.

16.4.5 Aseptically Pipette 1ml of the solution from above in a sterile Petri dishes

16.4.6 Add approximately 15-20ml of sterile Soyabean Casein Digest Agar prepared as per SOP(BML-QDG-MC-011), (cooled approximately up to 45°C) in two petri plates for TAMC

16.4.7 Add approximately 15-20ml of sterile Saboraud Dextrose Agar prepared as per SOP (cooled approximately up to 45°C) in two petri-plates for TYMC

16.4.8 Gently swirl the contents inside the petri-dishes for proper mixing of the sample, allow medium to solidify, taking the precaution of not spilling the media from the plate

16.4.9 Simultaneously perform the negative control by plating 1 ml of diluents used instead of the sample for each medium. Add 15-20ml of the media previously maintained not more than 45°c and allow for solidification.

16.4.10 Incubate the plates at 30°C to 35°C for 3 - 5 days for soybean Casein Digest Agar for TAMC and plates of Saboraud Dextrose Agar at 20^{0}C to 25°C for 5 days in an inverted position.