BIOMEDICAL LIMITED ILORIN	STANDARD TESTING PROCEDURE **TITLE:** STANDARD CHEMICAL TESTING PROCEDURE FOR BIOBABE SYRUP (60ml)		
DEPARTMENT:	**QUALITY**	**SECTION:**	**QUALITY CONTROL**
STP NO.:	**BML/QA/STP/FPS/012**	**PAGE NO.:**	**Page 1 of 9**
STP Ref.:	**BML-QDG-QA-105**	**REF. BOOK**	**BP 2024/IN-HOUSE**
EFFECTIVE DATE:	**04/01/2024**	**REVISION NO:**	**01**
REVIEW DATE:	**04/01/2026**	**SUPERSEDE:**	**00**

S/N	TEST	DESCRIPTIONS OF PROCEDURE
1.	**pH**	1. Take sample sufficient to immerse the electrode-head, and dip the electrode in the sample. 2. Wait until the reading stabilises, then take readings.
2.	**Diphenhydramine HCl** **Label Claim: 7mg/5ml**	**Method 1** Reagents: 1. Buffer solution, pH 3.0: 100mL of 0.1N potassium hydrogen phthalate and 40.6mL of 0.1N hydrochloric acid are mixed and diluted to 200mL. OR Dissolve 2.042g of potassium hydrogen phthalate in water and made up to 100mL. 2. Bromocresol green (BCG) solution: dissolve 69.8mg of the dye in 2mL of 0.1N sodium hydroxide and dilute to 1000mL with water. OR 34.9mg of the dye is dissolved in 100mL of 0.1M NaOH and diluted to 500mL with water. ***Standard solution:*** 20mcg/mL of the substance in water.

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		

		Sample solution: Accurately weigh a quantity of the sample equivalent to 20mg of the compound and dissolve in water (20mcg/mL). *5mL of syrup made to 100mL, 28.6mL of the resulting solution is made up to 100mL.* **Procedure:** • To 15ml each of the sample and standard solutions, add 60ml of dye solution. • Add 1 to 2 drops of 1N hydrochloric acid till yellow colour is obtained. • Add 10ml of buffer solution. • Extract the complex with three 15ml portion of chloroform. • Combine the chloroform layer dilute to 50ml. • Add few crystals of anhydrous Sodium Sulphate and shake. • Measure the absorbance of each sample and standard solutions at 415nm using chloroform as a blank. • Calculate the results by comparison

		% Assay of Diphenhydramine HCl = $\frac{absorbance\ of\ sample}{absorbance\ of\ standard}$ x 100 **Note:** 1. *Under acidic conditions, most substances with ternary amino group or quaternary ammonium salts make yellow complexes with the dye, extractable with chloroform.* 2. *Ammonium chloride, menthol, sodium citrate, paracetamol, saccharine, vitamin C, phenylephrine do not interfere.*
3.	**Paracetamol** **Label Claim:** **120mg/5ml**	**METHOD 1:** ***Standard solution***: 10mcg/mL in 0.1M hydrochloric acid. • Weigh accurately and dissolve 0.0249g of standard paracetamol in 0.1M HCl, and make up the volume to 50mL • From the resulting solution, take 1mL to 50mL with the same solvent. ***Sample solution***: • Dissolve an equivalent of 0.5g PCM [20ml Syrup] in 0.1M HCl and make up to

100ml std. flask.

- Dilute 1ml of resulting solution with 0.1M HCl to 50ml std. Flask.
- Take 5ml of final solution and dilute to 50 ml STD flask.
- Read absorbance at 244nm and compare results with standard

% Assay of Paracetamol = $\frac{absorbance\ of\ sample}{absorbance\ of\ standard}$ x 100

REFERENCE: Sethi's Quantitative Analysis of Pharmaceutical Formulations. Vol. 2

METHOD 2:

Standard Solution:

1mg/100mL

- Weigh accurately and dissolve 120mg of standard paracetamol in 0.1M HCl, and make up the volume to 100mL
- From the resulting solution, take 1mL to 100mL with the same solvent.

Sample Solution:

- Dissolve an equivalent of 120mg of the syrup (5mL) of the syrup make up the volume to 100mL with 0.1M HCl.

- From the resulting solution, take 1mL to 100mL with the same solvent
- Read absorbance at 244nm and compare results with standard

NOTE: in acidic solution, paracetamol has lambda max 246nm, whereas in alkaline solution, it has lambda max 257nm. This bathochromic shift 246→257 nm is due to molecular re-arrangement in the solvent.

% Assay of Paracetamol = $\frac{absorbance\ of\ sample}{absorbance\ of\ standard}$ x 100

REFERENCE:

Sethi's Quantitative Analysis of Pharmaceutical Formulations. Vol. 2

METHOD 3:

Standard Solution:

1mg/100mL

- Weigh accurately and dissolve 100mg of standard paracetamol in 0.1M NaOH, and make up the volume to 100mL
- From the resulting solution, take 1mL to 100mL with the same solvent.

		Sample Solution: • Dissolve an equivalent of 100mg of the syrup (4mL) of the syrup in 0.1mL and make up the volume to 100mL. • From the resulting solution, take 1mL to 100mL with the same solvent • Read absorbance at 257nm and compare results with standard **NOTE:** in acidic solution, paracetamol has lambda max 246nm, whereas in alkaline solution, it has lambda max 257nm. This bathochromic shift 246→257 nm is due to molecular re-arrangement in the solvent. **% Assay of Paracetamol** = $\frac{absorbance\ of\ sample}{absorbance\ of\ standard}$x 100 **REFERENCE:** Sethi's Quantitative Analysis of Pharmaceutical Formulations. Vol. 2
4.	**Microbial Enumeration Test**	- Determination of total aerobic microbial count (TAMC) and total combined yeasts and molds count (TYMC) by the pour plate method. - Dispense 1 ml of diluted sample in the center of the Petri plate using a sterile micropipette or calibrated pipette. - Open the lid of the prepared media, flame its mouth.

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		

		- Pour about 15 mL of sterilized molten media at the appropriate temperature above the sample. - Close the lid of the plate then mix the sample in the media properly by gently swirling the plate. The plate is generally swirled in an “S” or “8” shape. - Allow the media to completely solidify. - Incubate the plate in an inverted position under suitable incubation conditions (Incubate the plates of Soybean Casein Digest Agar at 30°C-35°C for 3-5days for TAMC and the plates of Sabouraud Dextrose Agar at 20°C to 25°C for 5-7 days. **<u>TEST FOR ABSENCE OF PATHOGEN</u>** - Measure 10ml /10g of the samples, dilute 1n 90ml of nutrient broth - Transfer 10ml into 90ml of Tryptone soya broth - Incubate at 30-35^{0}c for 18-24hours - Transfer 1 ml of the dilution into 100ml of MacConkey broth and incubate at 42-44°c in 18-72hours - Subculture on MacConkey Agar or Eosin Methylene blue Agar and incubate at 30-35°c - For negative control, proceeding same as above except without sample, streak a portion of incubated sterile MacConkey broth medium on the surface of Eosin Methylene Blue Agar.

- Incubate plates at 30 to 35° for 48 hours.

A. Result Interpretation

- Following the incubation, developed colonies are observed. Each colony will account for one viable microbial cell or one colony-forming unit.
- If all the colonies are of the same type, then we can predict that the sample contained only one type of microbial genera.
- However, if there are different genera or species producing similar types of colonies, it is required to carry out identification test.
- If the colonies are of different morphology, then we can conclude that the sample contained a mixed population.

Count the colonies and calculate CFU/mL by using the formula:

$$\text{CFU/mL} = \frac{\text{Total number of colonies obtained x dilution factor}}{\text{Volume of sample used (aliquot)}}$$

- This will give the total number of viable microbial cells present in the given sample.
- For optimum count, the number of colonies must be between 20 – 200 CFU/mL. Beyond this limit, the whole procedure must be repeated.
- If the number of the colony is less than 20, it is suggested to use the sample of lower dilution, whereas, if the total number of colonies exceeds 200, it is suggested to use the sample of higher dilution on successive repeats.

		- The result should be recorded following the Standard testing procedure SOP **BML-QDG-QA-105** - If the colonies are fused or the whole plate is covered with a single colony, then report as "too numerous to count" (TNTC) and repeat the process of taking the sample at a higher dilution.

*LAF--- Laminar Air Flow

CURRENT REVISION	SUPERSEDE	REASON FOR REVIEW
00	NIL	FIRST ISSUE
01	00	Review of the procedures Change in header and footer format

	DESIGNATION	SIGNATURE	DATE
PREPARED BY:	QUALITY CONTROL SENIOR ANALYST		
REVIEWED BY:	QUALITY CONTROL HEAD		
APPROVED BY:	QUALITY ASSURANCE MANAGER		