

STA2201H Methods of Applied Statistics II

Monica Alexander

Week 9: More hierarchical models

Notes

- ▶ Midterm
- ▶ A3 out this week. Optional (changed assessment structure)
- ▶ Also have to hand in research proposal and EDA

Research project

- ▶ Pick a dataset and research question and write a short paper
- ▶ Some examples from last year on Quercus (posted with permission)
- ▶ No real rules, but Stan/Bayesian (hierarchical) models **strongly** encouraged
 - ▶ GLMs
 - ▶ Hierarchical models
 - ▶ Temporal smoothing
 - ▶ Penalized splines
 - ▶ Measurement error/ data models
- ▶ Research proposal (DUE WITH A3):
 - ▶ What is your question and why is it interesting/important?
 - ▶ (brief) what is known? Hypotheses?
 - ▶ What's your dataset and main outcome/covariates of interest?
 - ▶ EDA: characteristics of dataset, any key patterns in outcome/relationships with other covariates
 - ▶ Should be done in the one Rmd/quarto file (hand in both Rmd/quarto and pdf)

Overview

- ▶ Funnels of hell and reparameterization
- ▶ GLMs in a hierarchical context

Reading

- ▶ Lesaffre and Lawson, ‘Bayesian Biostatistics’. Lip cancer example is from here.
- ▶ GH Chapters 14-15
- ▶ BDA Chapters 15-16
- ▶ A nice overview of hierarchical funnels:
<https://arxiv.org/pdf/1312.0906.pdf>

Hierarchical models and funnels

Back to radon

$$y_i | \alpha_{j[i]}, \beta_{j[i]} \sim N\left(\alpha_{j[i]} + \beta_{j[i]} x_i, \sigma_y^2\right)$$

$$\alpha_j \sim N(\mu_\alpha, \sigma_\alpha^2)$$

$$\beta_j \sim N(\mu_\beta, \sigma_\beta^2)$$

- ▶ y_i is log radon level
- ▶ α_j is county-specific intercept
- ▶ β_j is county-specific slope
- ▶ x_i is floor 1/ basement dummy

Let's fit this in Stan

Just a moment

$$\begin{aligned}y_i | \alpha_{j[i]}, \beta_{j[i]} &\sim N\left(\alpha_{j[i]} + \beta_{j[i]}x_i, \sigma_y^2\right) \\ \alpha_j &\sim N(\mu_\alpha, \sigma_\alpha^2) \\ \beta_j &\sim N(\mu_\beta, \sigma_\beta^2)\end{aligned}$$

- ▶ What's the interpretation of σ_β^2 ?
- ▶ What happens to the β s when σ_β^2 is small?

Fit in Stan

$$\alpha_1 = \alpha_j$$
$$\alpha_2 = \beta_j$$

```
transformed parameters {
  vector[N] y_hat;

  for (i in 1:N)
    y_hat[i] = a1[county[i]] + a2[county[i]] * x[i];
}

model {
  mu_a1 ~ normal(0, 1);
  a1 ~ normal(mu_a1, sigma_a1);
  mu_a2 ~ normal(0, 1);
  a2 ~ normal(mu_a2, sigma_a2);

  y ~ normal(y_hat, sigma_y);
}
```


Fit in Stan

Fit to MN only. Here's some of the results:


| | mean | se_mean | n_eff | Rhat |
|-------------|------------|-------------|------------|----------|
| ## a1[1] | 1.1774943 | 0.005621286 | 2137.37787 | 1.000478 |
| ## a1[35] | 1.0571842 | 0.006286536 | 1510.52071 | 1.001698 |
| ## a2[1] | -0.6357303 | 0.006656921 | 1904.38698 | 1.000949 |
| ## a2[35] | -0.6264791 | 0.007461098 | 1091.30463 | 1.005544 |
| ## sigma_a1 | 0.3404846 | 0.001891062 | 644.68028 | 1.001531 |
| ## sigma_a2 | 0.2795236 | 0.030739752 | 19.36188 | 1.213995 |

Let's look at some diagnostics

Pick a county (number 35) and plot the trace and density




β_{35}




What about the variance parameter

What's happening here?



σ_β

Scatterplot of β_{35} and σ_β



Funnels of hell

- ▶ The density of these models looks like a ‘funnel’, with a region of high density but low volume below a region of low density and high volume
- ▶ This property of the posterior is a characteristic of the model and not a problem in itself, but makes sampling hard
- ▶ Especially a problem for Gibbs, but still a problem for HMC
- ▶ The narrower the space, the smaller the step size has to be
- ▶ Larger step sizes more likely to get rejected, so the sampler can get stuck

We fit a centered parameterization

$$\begin{aligned}y_i | \alpha_{j[i]}, \beta_{j[i]} &\sim N\left(\alpha_{j[i]} + \beta_{j[i]}x_i, \sigma_y^2\right) \\ \alpha_j &\sim N(\mu_\alpha, \sigma_\alpha^2) \\ \beta_j &\sim N(\mu_\beta, \sigma_\beta^2)\end{aligned}$$

- ▶ e.g. β_j is directly dependent on hyperparameters μ_β and σ_β^2

Non-centered parameterization

$$y_i | \alpha_{j[i]}, \beta_{j[i]} \sim N\left(\alpha_{j[i]} + \beta_{j[i]} x_i, \sigma_y^2\right)$$

$$\begin{aligned} \beta_j &= \mu_\beta + \eta_j \cdot \sigma_\beta \\ \eta_j &\sim N(0, 1) \end{aligned}$$

...

- ▶ β_j is now a global mean + some offset, scaled by the β -specific standard deviation.
- ▶ In a non-centered parameterization we do not try to fit the group-level parameters directly, rather we fit a latent Gaussian variable from which we can recover the group-level parameters with a scaling and a translation
- ▶ The variables we are actually sampling are uncorrelated

Non-centered Stan model

```
transformed parameters {
  vector[85] a1;
  vector[85] a2;
  vector[N] y_hat;


  a1 = mu_a1 + sigma_a1 * eta1;
  a2 = mu_a2 + sigma_a2 * eta2;

  for (i in 1:N)
    y_hat[i] = a1[county[i]] + a2[county[i]] * x[i];
}


model {
  mu_a1 ~ normal(0, 1);
  mu_a2 ~ normal(0, 1);
  eta1 ~ normal(0, 1);
  eta2 ~ normal(0, 1);
  y ~ normal(y_hat, sigma_y);
}
```

sigma-a ~ normal(0, 1)
sigma-y ~ normal(0, 1)


Diagnostics for σ_β



Better exploration of the funnel



Less dependence between sampled parameters



Summary

- ▶ Funnel structure of posterior density is a consequence of hierarchical model structure
- ▶ Not just the shape but the mass of density
- ▶ Mostly a problem when you don't have much data! More shrinkage = more mass in narrow bit of funnel

GLMs in a hierarchical context

GLMs in a hierarchical context

- ▶ Last week we introduced hierarchical models in the setting where the data are assumed to be normally distributed
- ▶ But can easily extended to model other types of non-normal data hierarchically

Let's revisit the Poisson case:

$$y_i | \lambda_i \sim \text{Poisson}(\lambda_i)$$

where $\lambda_i > 0$, may depend on covariates, and vary by group membership $j[i]$.

Lip cancer

- ▶ Let's look at an example of estimating the risk of lip cancer by region in the former German Democratic Republic (Lesaffre and Lawson, chapter 9)
- ▶ In 1989, 2,342 deaths were recorded from lip cancer among males in 195 regions of GDR.
- ▶ For each region i we observed the number of deaths y_i
- ▶ We also know the expected count e_i , based on
 - ▶ age-specific mortality rates for whole country
 - ▶ age distribution of each region
- ▶ We also know the percentage of the population working outside
- ▶ Goal: estimate **relative risk** for each region θ_i

Lip cancer

ratio

- ▶ Define the standardized mortality rate $SMR_i = y_i/e_i$
- ▶ This is an estimate of the underlying relative risk θ_i , which captures the relative difference in risk of dying for region i as compared to the reference population.
- ▶ Problem of small populations, some of the SMRs are based on very low counts, so are very uncertain
 - ▶ 15% based on counts < 5
 - ▶ 62% based on counts < 10


Set-up

$$y_i \sim \text{Poisson}(\theta_i \cdot e_i)$$


log $\theta_i = ??$

- ▶ From last week, in a broad sense, what are our three options for θ_i ?

What the data look like



What the data look like



Lip cancer

100 countries; 1.

$$y_i \sim \text{Poisson}(\theta_i \cdot e_i)$$
$$\log \theta_i \sim \text{normal}(\mu_i)$$
$$y_i \sim \text{Poisson}(\theta_i \cdot e_i)$$
$$\log \theta_i \sim \text{normal}(\mu_i)$$

- 2.
1. Model θ_i s separately
 2. Find one θ for all regions
 3. Use hierarchical model to exchange information about θ_i s across regions

$$y_i \sim \text{Poisson}(\theta_i \cdot e_i)$$

$$\log \theta_i = \alpha_i + \beta(x_i^c)$$

$$\alpha_i \sim N(\mu, \sigma_\mu^2)$$

Where x_i^c is the (centered) percent of male population engaged in agriculture/forestry and fisheries in region i

3.

$$y_i \sim \text{Poisson}(\theta_i \cdot e_i)$$
$$\log \theta_i \sim \text{normal}(\mu, \sigma^2)$$
$$\sigma \sim N(0, 1)$$


Lip cancer

Full model

$$\begin{aligned}y_i &\sim \text{Poisson } (\theta_i \cdot e_i) \\ \log \theta_i &= \alpha_i + \beta(x_i^c) \\ \alpha_i &\sim N(\mu, \sigma_\mu^2) \\ \mu, \beta &\sim N(0, 1) \\ \sigma_\mu &\sim N_+(0, 1)\end{aligned}$$

Fitting in Stan

```
data {  
    int<lower=1> N;  
    vector[N] x;  
    vector[N] offset;  
    int<lower=0> deaths[N];  
    int<lower=0> region[N];  
}  
parameters {  
    vector[N] alpha;  
    real mu;  
    real beta;  
    real<lower=0> sigma_mu;  
}  
model {  
    vector[N] log_lambda;  
    for (i in 1:N){  
        log_lambda[i] = alpha[i] + beta*x[i] + offset[i];  
    }  
  
    alpha ~ normal(mu, sigma_mu);  
  
    mu ~ normal(0, 1);  
    beta ~ normal(0,1);  
    sigma_mu ~ normal(0, 1);  
  
    deaths ~ poisson_log(log_lambda);  
}
```



Stan interlude

Note in the previous slide I wrote

```
deaths ~ poisson_log(log_lambda)
```

for the likelihood.

There are a million (well, at least two) other options, see here:

https://mc-stan.org/docs/2_18/functions-reference/poisson.html

- ▶ target += poisson_lpmf(deaths | lambda)
i.e. "Increment target log probability density with..."
- ▶ deaths ~ poisson(lambda) this is shorthand for the above,
to make people used to BUGS/JAGS happier.

In the above example, I used `poisson_log` so I didn't have to exponentiate my expression, but you do you.

Also note that I used an 'offset' in the same way that we did in `glm`, but you could equally write in terms of the expected deaths
e.g. `lambda[i] = theta[i]*expected[i]`, etc etc


Interpretation of coefficient on proportion working outside?

$$\text{exp}(\beta)$$

```
##                               mean      se_mean    n_eff   Rhat  
## mu      0.08643664 0.0005393231 4491.138 0.9999066  
## beta    1.97390770 0.0068382166 2423.780 1.0004250  
## sigma_mu 0.38661177 0.0006415591 2319.308 0.9994168
```

$$y_i \sim \text{Poisson}(\theta_i \cdot e_i)$$
$$\log \theta_i = \alpha_0 + \beta x_i$$

Observed SMR v estimated θ_i



Overdispersion

- ▶ Recall that in many applications, counts are likely to be overdispersed
- ▶ Actually not bad in lip cancer case (how do you tell?)
- ▶ Going back k weeks, what were our two main options for dealing with overdispersion?

Overdispersion with a quasi-Poisson set-up

In the lip cancer case we had

$$y_i \sim \text{Poisson} (\exp(\alpha_i + \beta x_i^c) \cdot e_i)$$

...

For the overdispersed case we would have


$$y_i \sim \text{Poisson} (\exp(\alpha_i + \beta x_i^c + \varepsilon_i) \cdot e_i)$$

...

$$\varepsilon_i \sim N(0, \sigma_\varepsilon^2)$$

...

How to compare?



Logistic regression

Logistic regression in a hierarchical context

Can easily extend these hierarchical models to model binary outcomes and the probability of an event happening by various groups of interest, e.g. in the simplest form we would have

$$y_i | \pi_i \sim \text{Bern}(\pi_i), \text{ OR}$$

$$y_i | \pi_i \sim \text{Bin}(n_i, \pi_i), \text{ if total number of trials is } n_i$$

and then

$$\begin{aligned}\text{logit}(\pi_i) &= \log\left(\frac{\pi_i}{1 - \pi_i}\right) \\ &= \text{some function of covariates, e.g.}\end{aligned}$$

$$\text{logit}(\pi_i) = \beta_0 + \beta_1 x_i$$

and then could put some hierarchical structure on the β 's for instance.

Fertility intentions

- ▶ Using data from the 2015-2017 National Survey of Family Growth
- ▶ Gathers information on family life, marriage and divorce, pregnancy, infertility, use of contraception, and men's and women's health in the US.
- ▶ Some microdata are public:
<https://www.cdc.gov/nchs/nsfg/index.htm>

Fertility intentions

Interested in looking at fertility intentions, i.e. do women intend to have more children (and if so, how many)


- ▶ predictor of childbearing
- ▶ gaps tell you something about unmet need, due to social, economic conditions, cultural changes

We are interested in the probability that a women wants more children.


- ▶ Individual covariates: age, education, marital status
- ▶ Possible extensions: state/region, parity (current number of children)

Some clear patterns

Observed proportion of women wanting more kids



Sample size differences



Non-nested hierarchical model

- ▶ So far we have considered the simplest hierarchical structure of individuals i in groups j
- ▶ In the polls example, individuals have different group memberships (based on age and education) that we may want to pool across.

E.g. a reasonable model set-up would be

- ▶ to have hierarchies for age and education
- ▶ an extended model that includes geography may also have county nested within state, for example

Just one hierarchical model

A hierarchical model with just age would be

$$y_i | \pi_i \sim \text{Bern}(\hat{\pi}_i)$$

Probability of having more children

$$\Pr(y_i = 1) = \pi_i = \text{logit}^{-1}(\alpha_j[i])$$

$$\alpha_j \sim N(\mu_\alpha, \sigma_\alpha^2), \text{ for } j = 1, \dots, 5$$

$$\text{logit}(\pi_i) = \alpha_j[i]$$

this age group
of women

Add in other effects


$$y_i | \pi_i \sim \text{Bern}(\pi_i)$$
$$\pi_i = \text{logit}^{-1}(\beta_0 + \beta_1 \text{formerly married}_i + \beta_2 \text{married}_i + \alpha_{j[i]}^{\text{age}} + \alpha_{k[i]}^{\text{edu}})$$
$$\alpha_j^{\text{age}} \sim N(0, \sigma_{\text{age}}^2), \text{ for } j = 1, \dots, 5$$
$$\alpha_k^{\text{edu}} \sim N(0, \sigma_{\text{edu}}^2), \text{ for } k = 1, \dots, 4$$
$$\beta_1 \sim N(0, 1)$$
$$\beta_2 \sim N(0, 1)$$

- ▶ Indexes: individual i , age j , education k
- ▶ Notice that the effects for age, educ, are centered around zero and now there is a 'global' intercept β_0
 - ▶ any non-zero means for the α s could be folded into the global intercept
- ▶ The baseline category is never married

Some results

α_j age


Estimated age effects




Some results

α_{educ}
 β

Estimated education effects




Remember we had no observations from one cell



Estimates from model

$$\rightarrow \text{logit } \pi_i = \beta_0 + \beta_1 \dots + \cancel{\beta_2} \cancel{\text{age}} + \alpha_{\text{age}} + \alpha_{\text{age}} \pi_i + \cancel{\alpha_{\text{age}}} \cancel{\pi_i}$$

Estimated probabilities of wanting more children
20–24 year olds with Bachelor+



age
20-24
educ
Bachelor

$$y_i \sim \text{Bern}(\pi_i)$$

What about getting estimates for the 45-49 year olds?

How do we generate these?

$$y_i | \pi_i \sim \text{Bern}(\pi_i)$$

$$\pi_i = \text{logit}^{-1} \left(\beta_0 + \beta_1 \text{formerly married}_i + \beta_2 \text{married}_i + \alpha_{j[i]}^{\text{age}} + \alpha_{k[i]}^{\text{edu}} \right)$$


$$\alpha_j^{\text{age}} \sim N(0, \sigma_{\text{age}}^2), \text{ for } j = 1, \dots, 5$$

$$\alpha_k^{\text{edu}} \sim N(0, \sigma_{\text{edu}}^2), \text{ for } k = 1, \dots, 4$$

$$\hat{\alpha}_{45-49}^{\text{age}} \sim N(0, \sigma_{\text{age}}^2)$$

Results, compared to 40-44 year olds

Estimated probabilities of wanting more children
Married with Bachelor+



Issues

We draw a new $\tilde{\alpha}_j^{\text{age}}$ based on

$$\tilde{\alpha}_j^{\text{age}} \sim N(0, \sigma_{\text{age}}^2)$$

This assumes the age effects are conditionally independent of each other and centered at zero.

But we can do better!

An alternative specification on age

Consider the model as before but with one change:

$$y_i | \pi_i \sim \text{Bern}(\pi_i)$$

$$\pi_i = \text{logit}^{-1} \left(\beta_0 + \beta_1 \text{formerly married}_i + \beta_2 \text{married}_i + \alpha_{j[i]}^{\text{age}} + \alpha_{k[i]}^{\text{edu}} \right)$$


$$\alpha_j^{\text{age}} \sim N \left(\alpha_{j-1}^{\text{age}}, \sigma_{\text{age}}^2 \right), \text{ for } j = 2, \dots, 5$$

$$\alpha_k^{\text{edu}} \sim N \left(0, \sigma_{\text{edu}}^2 \right), \text{ for } k = 1, \dots, 4$$

- ▶ This is assuming that the age effect in group j is similar to that in the previous age group (plus some noise)
- ▶ We are placing a random walk on the age effects; assuming some structure over age
- ▶ Note: need to put a prior on the first age group here

Now look at the estimated probabilities for 45-49 year olds

Estimated probabilities of wanting more children
Married with Bachelor+, alternative model




Summary

- ▶ Can model non-normal data in hierarchical setting
- ▶ Can have different hierarchies going on at the same time
- ▶ In the fertility intentions case, hierarchical models makes sense here because cell counts get very small: want to stabilize estimates.
- ▶ We can think about structured priors on the hierarchical effects (e.g. over age)
 - ▶ this idea extends to other dimensions, e.g. particularly thinking about geographic space

Measurement error

Measurement error

- ▶ We have talked about hierarchical models and how to fit them in Stan
- ▶ Often consider something like

$$y_i | \mu_i, \sigma^2 \sim N(\mu_i, \sigma^2)$$


with a model on μ_i . But what is σ^2 ?


- ▶ So far, have assumed it's constant for all observations i and estimated in the model
- ▶ But often have some info based on how the data were collected
- ▶ Can incorporate info about measurement error into our models

Motivating example

- ▶ Interested in estimating and projecting neonatal mortality in Sri Lanka over time
- ▶ Data available are from different sources which have differing degrees of error
- ▶ Best: vital registration systems (VR) but this hasn't always existed in Sri Lanka
- ▶ Also rely on survey data

Motivating example

Ratio of neonatal to other child mortality (logged), Sri Lanka



Child mortality in Sri Lanka

Goals: estimate expected level of ratio over time

Issues:

- ▶ overlapping observations
- ▶ missing years
- ▶ different data sources
- ▶ different errors

Let's start off simple

$$t = 1955 \text{ to } 2015$$

For starters with the Sri Lankan data, let's model just a linear function over time

$$y_t \sim N(\mu_t, \sigma^2)$$

with

$$\mu_t = \alpha + \beta(t - t_c)$$

t_c is the mid-year of the study period.

But there's an issue with the indexes here!



Allowing for overlapping observations and missing data

A pretty straightforward extension:

$$y_i \sim N(\mu_{t[i]}, \sigma^2)$$

with

$$\mu_t = \alpha + \beta(t - t_c)$$

where $t[i]$ is the same indexing as in hierarchical case: the year t which observation i relates to.

Fit in Stan

```
data {
    int<lower=0> N; //number of observations
    int<lower=0> T; //number of years
    int<lower=0> mid_year;
    vector[N] y; //log ratio
    vector[T] years; // vector of unique years (1955 ... 2015)
    int<lower=0> year_i[N]; // year index for observation i
}

parameters {
    real alpha;
    real beta;
    real<lower=0> sigma;
}
```

Fit in Stan

```
transformed parameters{
  vector[T] mu;

  for(t in 1:T){
    mu[t] = alpha + beta*(years[t] - mid_year);
  }
}

model {
  vector[N] y_hat;


  y ~ normal(mu[year_i], sigma);

  alpha ~ normal(0, 1);
  beta ~ normal(0,1);
  sigma ~ normal(0, 1);
}
```

Results

Yin/N(m, β)

Ratio of neonatal to other child mortality (logged), Sri Lanka
Linear fit



Incorporating measurement error


$$y_i \sim N(\mu_i, \sigma^2)$$

↑
estimate
new
constant

Adding in the measurement error (standard errors based on sampling in this case) involves swapping out the estimated σ^2 with data:

$$y_i | \mu_i \sim N(\mu_i, s_i^2)$$

where s_i is the observed standard error.



Stan


```
model {  
    y ~ normal(mu[year_i], se);  
    alpha ~ normal(0, 1);  
    beta ~ normal(0,1);  
}
```

vector[N] se;

Result

Ratio of neonatal to other child mortality (logged), Sri Lanka

Linear fit, red = ME, black = no ME



Take-aways

- ▶ Easy to account for missing data with right index set-up
- ▶ Accounting for measurement error useful when have data from different sources

This was a pretty simple linear model, can we do better?