

# The Theory of the Universal Wave Function and the Origins of the Cosmos

Maya Benowitz\*  
(Dated: October 27, 2024)

I assume quantum mechanics applies to the entire universe and solve the Schrödinger equation exactly, obtaining a unique closed-form wave function of the universe. Quantum information and quantum geometry emerge as one from the bootstrap. The geometry of quantum information is central to the problem of time in quantum cosmology, which is resolved by a flat space holographic duality between a timeless theory in Euclidean 3-space without a spin-2 field and a theory in  $3 + 1$  spacetime dimensions with an emergent spin-2 field. Among the observable consequences are two spectacular predictions: the dark energy is a quantum scalar curvature field that decays with the scale factor like  $a^{-2}$ , and all black holes are dark energy composite objects with a mass function that grows like  $a^3$ .

## I. MOTIVATION

The cosmos seems to have been born into a spatially flat and thermal state of low entropy, originating from a curvature singularity beyond the effective field theory (EFT) description of space, time, matter, and energy [1]. This grand puzzle demands new physics. This research program aims to discover a unifying principle of the physics of creation and its observable consequences.

The mythos of our shared humanity is a liminal space of creation stories. The origin of the universe is the ultimate question and a timeless puzzle. The philosophy of this research program is a form of monism that hypothesizes all of reality can be derived from a single unifying mathematical principle—an ancient idea. But physicists need more than philosophy and mathematics. We need empirical data, and the empirical data is conclusive: the universe is quantum.

Understanding the nature of the cosmological singularity requires a complete theory of quantum gravity. Indeed, the conceptual problems in the foundations of physics at extremely early times do not begin at the Planck scale [2]. The Copenhagen interpretation of quantum mechanics (QM), which assumes a Heisenberg cut between quantum and classical systems for wave function collapse, breaks down when the scale of the universe is on the order of an atom. QM in the 'laboratory frame' applied to the early universe as a whole requires the troubling assumption of an external classical observer to prepare the initial state of the cosmos. I argue Copenhagen can not be the final formulation of quantum mechanics. Indeed, a compendium of clues points elsewhere.

One of the most consequential experiments in the foundations of physics is the Aharonov–Bohm effect [3–6]. How can a charged particle's gauge potential possibly couple to the complex phase of a nonphysical probability wave? Is the measured *change* in phase not an indirect measurement of the physical existence of the wave function? There is still more evidence of the reality of the quantum state. The Pusey–Barrett–Rudolph (PBR)

no-go theorem has been experimentally confirmed: either the quantum state is an objective physical state, or the assumption of preparation independence is violated, and the experimenter has no free-will in the choice of measurement settings [7–9]. But if the wave function is physical, where do probability amplitudes and the Born rule come from? The measurement problem, the problem of the external observer, and the problem of time in quantum cosmology are all deeply related [10]. These foundational conceptual issues can not be ignored in a fundamental theory. To understand the quantum origins of space and time, we must first understand the true nature of quantum mechanics.

The most cogent formulation of QM with a physical wave function is Everett's seminal work: *The Theory of a Universal Wave Function* [11, 12]. Everett profoundly impacted Wheeler and DeWitt, inspiring the Wheeler–DeWitt (WdW) equation and the field of quantum cosmology. The development of the canonical quantum gravity program of WdW and what became popularized as the many worlds interpretation diverged. Work in the foundations of QM became orthogonal to quantum gravity.

On the one hand, Everettian QM is done almost exclusively in the non-relativistic limit. Space and time are not just classical. They are Newtonian! The creation of new worlds (i.e., branching) happens with respect to a universal clock in strictly finite-dimensional Hilbert spaces. What happens if we replace the Schrödinger equation with a relativistic wave equation in the Everettian framework and work in a covariant fashion with an infinite-dimensional Hilbert space?

On the other hand, WdW *assumes* general relativity (GR) as an input to quantize, and out comes an ill-defined infinite dimensional partial differential equation with pathological consequences. There are no known unitary solutions of WdW [13]. What is more, the Hamiltonian vanishes on physical states with compact geometries in general covariant theories, meaning there is an infinite number of possible boundary conditions to impose on the equation and, therefore, an infinite number of possible solutions. Vacuum solutions are not unique. But if the universal wave function contains all physical information of the universe, there can only be One.

---

\* mayabenowitz@gmail.com

General covariance is necessary, but is it sufficient? To the best of human knowledge, nature is quantum from the beginning. Quantization schemes—like coordinate systems—are an artifact of formalism. This guiding principle of *quantum covariance* is a radical constraint. But without intuition of the quantum world at the moment of creation, how is one to even begin?

The standard lore in the foundations of physics tells us degrees of freedom at higher energies (smaller distances) are more fundamental than those at lower energies (larger distances). At the heart of the Wilsonian EFT framework is the assumption that the macroscopic dynamics of a system are independent of the microscopic details. But is this true for gravity? Is the observation of the cosmological constant entering local galaxy dynamics through a universal acceleration scale  $a_0 \approx cH_0 \approx c^2\Lambda^{1/2}$  [14–16] merely a numerical coincidence or a profound clue into the nature of quantum gravity? How is it that something as small as a galaxy “knows” about the de Sitter (dS) radius associated with the whole of the observable universe? UV/IR mixing appears to be a model-independent feature of quantum gravity, with examples emerging across string theory and beyond [13, 17–32]. This begs the question: does gravity violate the separation of scales?

Perhaps moving deep into the IR, at the scale of the entire universe, is no less fundamental than working in the UV. Indeed, this perspective is supported by the data. EFT predicts the dark energy is dominated by the UV vacuum energy (the Planck energy). Yet, it is observed to be astonishingly smaller (by 120 orders of magnitude), meaning it sets the IR cutoff that determines the largest scale anyone shall ever see. Motivated by this compendium of clues, I assume reductionism works in both directions. This *universal democracy*, a generalization of Chew’s “nuclear democracy” to cosmology, suggests the quantum degrees of freedom of the entire universe *are* the fundamental UV degrees of freedom. Intriguingly, this view was discovered at the end of the second superstring revolution with holography [33–39]. In some deeply mysterious way, the fundamental degrees of freedom exist at the boundary of spacetime—at infinity—where a covariant definition of energy is well-defined in both quantum mechanics (QM) and GR.

The guiding principles of quantum covariance and universal democracy motivate the quantum-first program recently proposed by Carroll, the idea that spacetime and quantum fields emerge from the wavefunction of the universe [41, 42]. Carroll and friends [43] approach this program in a non-covariant fashion, leaving a clear gap in the literature. In general covariant theories of quantum gravity, the Hamiltonian annihilates the universal quantum state. Time drops out of the Schrödinger equation, meaning all information of the physical universe lives at future infinity, where time stands still. A notion consistent with the experimental realization of the Page-Wootters mechanism in continuous time [40].

In this paper, I report a minimal bootstrapping of quantum universality. No relic classical baggage

is assumed. To the author’s knowledge, this is the first attempt to write down the quantum state of the universe that is not some semiclassical approximation like the Hartle-Hawking no-boundary proposal [44, 45], Vilenkin’s tunneling proposal [46, 47], or numerous minisuperspace models [48, 49], which all assume the WdW equation as a starting point [50].

Modulo continuity, no assumptions of spacetime, matter, or energy are made and put in by hand. Instead, the One fundamental assumption is as follows.

#### Everett’s principle of quantum universality

The wavefunction of the universe exists and evolves unitarily, **always** obeying the Schrödinger equation:

$$\hat{H}\Psi = i\partial_t\Psi. \quad (1)$$

A corollary of quantum universality is this: nature has no fundamental scale—the *form* of the laws of physics remains unchanged on all scales.

From this elementary assumption, everything that follows is pure deduction. This research program is a formal nomology or deductive-nomological model that aims to derive relativistic principles from quantum universality, write down a relativistic cosmological Schrödinger equation, and explore its consequences.

## II. QUANTUM UNIVERSALITY

Quantum universality implies there is no Heisenberg cut between quantum and classical systems. The Everettian view is austere. The universe, on all scales, is quantum. If nature has no fundamental scale, the mass term  $M$  in the Schrödinger equation must multiply out, implying the

#### Everett-Wheeler-DeWitt equation

$$\hat{H}\Psi = 0. \quad (2)$$

Neither time nor mass-energy is fundamental. This further implies the wavefunction of the universe is scale-invariant:  $\Psi \sim a^n$  where  $a$  is the cosmological scale factor, and  $n$  is an arbitrary integer. One can now solve for the potential  $V$  and write down the following one-dimensional EWD equation:

$$\left(-\frac{\partial^2}{\partial a^2} + \frac{n(n-1)}{a^2}\right)\Psi = 0. \quad (3)$$

The above equation is not quite consistent with the ax-

ions of QM since  $\Psi$  can not be normalized. But observe the following. The potential  $V$  is the scalar curvature  $R$  of an  $n$ -sphere with radius  $a$ , the kinetic term is the scalar curvature of an  $n$ -hyperbola, and the right-hand side of the equation is the scalar curvature of Euclidean  $n$ -space. From this minimal conformal bootstrap, the fundamental space forms are derived!

GR assumes the equivalence between energy and curvature. Here, it is a consequence. A scalar curvature singularity is at  $V(a = 0)$ . Since the Hamiltonian annihilates the universal quantum state, the scalar curvature singularity cancels with the kinetic term, yielding a zero-total energy universe with global spatial flatness, thus providing a path to solving the cosmological singularity and flatness problems. In GR, the dark energy is proportional to the constant scalar curvature of dS space:

$$R = \frac{d(d-1)}{L^2} = \frac{2d}{d-2}\Lambda \quad (4)$$

where  $d$  is the spacetime dimension and  $L$  is the radius. However, there is no unitary evolution of  $\Psi$ . A common problem for the WdW equation and its approximations. The timeless equation occupies a liminal space between geometry and quantum mechanics. Under a change of variables, eq. (3) is the reduced radial equation of a zero-energy free particle, i.e., the simplest one-dimensional time-independent Schrödinger equation in elementary QM. There is a direct correspondence between  $V(a)$  and the effective centrifugal potential  $V_{\text{eff}}(r)$  with  $n \leftrightarrow -l$ , where  $l$  is the centrifugal term. A consequence that echoes Mach's principle and violates decoupling.

The physical picture is that of duality. The universe, as an (almost-quantum) whole, can be seen as a Lemaître point-like particle, or Cosmon, with charge  $M$  and position  $a$  from the origin, sourcing a repulsive potential that accelerates the Cosmon away from the origin and *decays to zero at infinity*. Seen from the geometric view, it describes a global charge  $M$  sourcing the scalar curvatures of an  $n$ -sphere and an  $n$ -hyperboloid that annihilates to flat space. The change of variables establishes a duality between *position* and *scale*. It follows from eq. (3) that the preferred basis is the scale basis. The equation is timeless, so one can interpret the Hamiltonian as describing a celestial sphere at future infinity (a kind of celestial holography [54, 55]) dual to a theory in one timelike (and lightlike) dimension higher. The equation is coordinate-independent, meaning general covariance is a consequence of quantum universality. The number of spatial dimensions  $n + 1$  is also unfixed.

Can we bootstrap our way to a unitary instantiation of the timeless equation and explain *why* we observe only three spatial dimensions? Yes, we can—with symmetry. There is a trivial  $\mathbb{Z}_2$  action  $a \rightarrow -a$  and a non-trivial  $\mathbb{Z}_2$  action  $n \rightarrow -n$ . The former is a symmetry of the system; the latter is not. Suppose we demand the latter is a symmetry of the system. The action  $n \rightarrow -n$  gives

the potential  $R_{n+1} = \frac{n(n+1)}{a^2}$  and is interpreted as the scalar curvature of an  $(n+1)$ -sphere. The potential then becomes,

$$V \sim \begin{pmatrix} R_n & 0 \\ 0 & R_{n+1} \end{pmatrix} \quad (5)$$

implying  $\Psi$  is promoted to a pair of complex numbers, i.e., a spinor. A complex 2-spinor solution, together with the consequence of conformal flatness, implies an underlying  $SU(n = 2)$  symmetry, forcing the self-consistent equations of motion to take the unique form of a

relativistic cosmological wave equation

$$(D^2 + \hat{V})\Psi = 0. \quad (6)$$

The  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetry is a fundamental symmetry of the boundary theory and will continuously reappear in the story.

The Dirac-spin operator

$$D = i(\sigma^1 \partial_1 + \sigma^2 \partial_2 + \sigma^3 \partial_3), \quad (7)$$

where the  $\sigma^i$  are the Pauli matrices. In the cosmological setting, the preferred basis is the scale basis:

$$\partial_1 = \partial_2 = \partial_3 = \partial_a.$$

In the limit  $\hat{V} \rightarrow 0$ , we have  $D^2\Psi = D\Psi = 0$ , i.e., the Dirac equation. For large-scale factors, solutions to  $\hat{H}\Psi = 0$  are effectively harmonic spinors, and the underlying space is a Calabi-Yau (spin) manifold.

The Dirac equation is a *conformal* 2-spinor relativistic wave equation for the base manifold  $M = \mathbb{R}^3$ . The physical interpretation is a (massless) spin-1/2 field over  $\mathbb{R}^3$  with a wave function  $\psi : \mathbb{R}^3 \rightarrow \mathbb{C}^2$ , where the components specifies the probability amplitude for the field configuration at each point in  $\mathbb{R}^3$  to be in the spin-up or spin-down state. The physical interpretation of the cosmological wave equation is a spin-1/2 field over  $\mathbb{R}^3$  where the components are the probability amplitudes of the field configuration at each point in  $\mathbb{R}^3$  to be in the expanding or contracting state. Two deductions are immediate.

First, probability takes a strictly frequentist meaning: in a spatially infinite universe, everything that can happen does; some things happen more than others. Consider the state of the observable universe at  $t_{\text{now}}$ . At some arbitrarily large distance beyond our causal horizon is a copy of our observable universe at time  $t_{\text{now}} + \delta t$ , in our future. *In the God's eye view, everything that can happen happens everywhere all at once.* Time is not fundamental in a universe with no fundamental scale. This is the origin of probability in QM.

Second, the potential scalar curvature  $V$  must be quantized. To drive the contracting state, an attractive or negative (quantum) scalar curvature component must emerge in  $\hat{V}$ . In solving the cosmological wave equation, we learn how to quantize geometry. As will soon become apparent, general relativity is a logical consequence of QM at its most minimal.

Let us take a moment to absorb the observable consequences. Starting from the simplest formulation of QM, we have deduced 1) there is an equivalence between energy and curvature, 2) the universe, as a whole, is flat with three spatial directions that are either 3) expanding or contracting at an accelerated rate, driven by a quantum scalar curvature that scales like  $a^{-2}$ .

We have also learned of a deep geometric connection between fermions and spatial directions. Space and fermions have emerged from a conformal bootstrap of quantum universality. And where there are spinors, there is spacetime. A map always exists between spinors of order  $n$  and tensors of order  $2n$ . We deduced  $n = 2$ , so it stands to reason that a path must exist to derive the dynamic field content of our world from the first principle.

### A. Geometry of quantum information

Before quantization, the components of  $V$  are the scalar curvatures of the 2-sphere and 3-sphere, respectively. The Hopf fibration uniquely relates these spheres: one can always think of the 3-sphere as a  $U(1)$ -fiber over the 2-sphere [51]. The fiber bundle structure of the qubit is the Hopf fibration, and the symmetry group of the qubit is  $SU(2)$ . Quantum information naturally follows from quantum universality. The mapping from the unit 3-sphere in a two-dimensional complex Hilbert space  $\mathbb{C}^2$  (otherwise known as the complex projective line  $\mathbb{CP}^1$  or the complex plane  $\mathbb{C}$  with a point at infinity) to the Bloch (or Riemann) sphere is the Hopf fibration. There is a direct correspondence between each spinor in  $\mathbb{C}^2$  and a point on the Bloch sphere. Schematically, we write the fibration as

$$\begin{array}{c} S^1 \hookrightarrow S^3 \\ \downarrow \\ S^2 \end{array} \quad (8)$$

i.e., embedding a 1-sphere in the 3-sphere 'wraps' the 3-sphere around the 2-sphere. The  $S^3$  lives on the surface of  $S^2$  by the identification of phase circles in  $S^3$  with points on  $S^2$ . Every possible state of a two-level system (i.e., qubit) lives on the surface of  $S^2 \cong \mathbb{CP}^1$ . The system under question (the entire universe) is not a two-level system. The components of  $\Psi$  are not simply elements of  $\mathbb{CP}^1$ . There is more structure that captures the intrinsic quantum nature of the  $S^2$ . Nevertheless, the geometry of quantum information is a foundational starting point in solving eq. (6) and emerging time and quantum fields from  $\Psi$ . We know the components of the spinor are not

real-valued power laws. They are complex and have a global  $U(1)$  phase symmetry. This means the potential will take a slightly different form from eq. (5), encoding the *quantum*  $S^3$  as a  $U(1)$ -fiber over the *quantum*  $S^2$ .

The Hopf fibration has a symmetry group: the action of the unitary group  $U(2)$  on  $\mathbb{C}^2$  leaves the  $S^3$  invariant. It thus descends to an action on the  $S^2$  by ordinary rotations, carrying fibers into fibers as it commutes with the  $U(1)$  action [51]. An even less known fact is that  $\tilde{M} = U(2)$  is a conformal compactification of Minkowski space with a "lightcone and 2-sphere at infinity" [52]. What is more, by the exceptional spin isomorphism  $Spin(3,1) \cong SL(2,\mathbb{C})$ , one may identify the points  $x_\mu$  in Minkowski space with the infinitesimal generators of  $U(2)$ ,

$$X = \sigma^\mu x_\mu = Ix_0 + \sigma^1 x_1 + \sigma^2 x_2 + \sigma^3 x_3. \quad (9)$$

Here the Lorentz metric norm is  $\|x_\mu\| = \det(X)$ . From this identification one finds that  $x_\mu$  is lightlike precisely if there is a complex 2-spinor  $\mathbf{2}$ , modulo a rescaling, such that  $X = \mathbf{2} \otimes \bar{\mathbf{2}}$ . This identifies the celestial sphere (the space of lines in the light cone that pass through a point in Minkowski space) with the Riemann sphere, i.e.,  $S^2 \cong \mathbb{CP}^1$ . The symmetry group of the simplest non-trivial fiber bundle is compactified Minkowski space! *Timelike and lightlike directions are derived from the Hopf bundle symmetry of the qubit.* A map always exists between rays of lightcone and points on the surface of the Bloch sphere. Seen from another perspective, the fundamental representations of the Lorentz group contain a pair of complex 2-spinors  $\mathbf{2}$  and  $\bar{\mathbf{2}}$ , i.e., two fermionic directions. Using the Littlewood-Richardson rule, the tensor product  $\mathbf{2} \otimes \bar{\mathbf{2}} = \mathbf{3} \oplus \mathbf{1}$  is the adjoint representation and can be identified with either Minkowski space or a spin-1 field. This is the motivation for a supersymmetry between spatial and fermionic directions.

### B. Quantum geometry

The "inner mechanism" of holography appears to be the simplest non-trivial (the exceptional) fibration between the spheres, i.e., where the base space  $S^2$  is one dimension lower than the total space  $S^3$ . But what, exactly, is a quantum sphere? What does it mean for geometry to be quantum? More precisely, what is the form of the quantum metric tensor? What new mathematical tools does one need to bootstrap a unique solution eq. (6)? Is a unique vacuum solution the key to building up an intuition of quantum geometry and writing down a complete theory of quantum gravity?

The  $\mathbb{Z}_2$  symmetry  $n \rightarrow -n$  leaves future infinity interchangeable with the cosmological singularity by taking  $\Psi(a)$  to  $\Psi(1/a)$ . This duality symmetry implies a compactification with a transition map that identifies zero and infinity as the same point, i.e., the inclusion of a point at infinity with a non-trivial topology that "glues" the singularity to future infinity. The quantum state  $\Psi$

belongs to some conformally flat and compact spin manifold in  $\mathbb{C}^2$ . The simplest such manifold is the Clifford torus. So perhaps our target space has the topology of a 2-torus embedded in  $\mathbb{C}^2$  with a non-trivial conformal spin structure. After some thought, the only such construction is as follows.




FIG. 1. The conformal Clifford torus  $\mathcal{T}^2$  with one dimension suppressed. The scale factor  $a$  goes around the 'line at infinity' and is real, while the imaginary axis  $\beta$  goes through the line at infinity. The left-hand side is defined by the pair  $(a, \beta) \in \mathbb{C}$ , whereas the right-hand side is defined by  $(1/a, 1/\beta) \in \mathbb{C}$ . The entire object lives in  $\mathbb{C}^2$  with the identification  $(a, \beta) \sim_\pi (1/a, 1/\beta)$  where  $\sim_\pi$  denotes  $(a, \beta)$  can be rotated into  $(1/a, 1/\beta)$  by a rotation of  $\pi$ . Above the imaginary axis, the scale factor is positive; below, it is negative. A rotation from  $0_L$  to  $0_R$  takes  $2\pi$  radians and sends  $a \rightarrow -a$ . Thus, the space is, by construction, a spin manifold.  $\beta \in i\mathbb{R}^+$  from  $0_L$  to  $\infty$  and  $1/\beta \in i\mathbb{R}^-$  from  $\infty$  to  $0_R$ . Both zeros are identified. The imaginary axis that runs through  $\infty$  is the thermal circle,  $S_\beta^1$ . Thus, each quadrant has the sign sequence  $(++, +-, --, -+)$ , which is independent of the choice of clockwise or counterclockwise rotation and is nothing but the  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetry.

The following deductions are immediate. The physical interpretation of  $\beta$  is the inverse temperature. The phase angle  $\phi = \phi(\beta)$  is a scalar temperature field. Topologically,  $\mathcal{T}^2 = S_a^1 \times S_\beta^1$  where  $S_\beta^1$  is the thermal circle. The thermal circle implies the de Sitter or Hawking temperature:

$$T \sim \frac{H}{2\pi} \quad (10)$$

where  $H$  is the Hubble constant. The values of the quantum state  $\Psi$  (the amplitudes) are pairs of points on  $\mathcal{T}^2$ , meaning the configuration space of all possible amplitudes is conformally flat and thermal.

We can geometrically bootstrap a unique vacuum solution to the cosmological wave equation from  $\mathcal{T}^2$ . Use

the  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetry and assemble the EWD equation and its dual,  $\hat{H}_n \psi = 0$  and  $\hat{H}_{n+1} \psi = 0$ , into a single pre-spinorial equation:

$$\begin{pmatrix} -\partial_a^2 + R_n & 0 \\ 0 & -\partial_a^2 + R_{n+1} \end{pmatrix} \begin{pmatrix} \psi_n \\ \psi_{n+1} \end{pmatrix} = 0. \quad (11)$$

We determined the space of states  $\mathcal{T}^2$  has the  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetry, so it follows  $\Psi$  too is invariant under this action. The most general solution to the above is

$$\psi_n \sim a^n + (-a)^n, \quad \psi_{n+1} \sim a^{-n} + (-a)^{-n}. \quad (12)$$

When  $n$  is odd, we have the trivial solution  $\psi_n = \psi_{n+1} = 0$ . The components are not complex-valued, that is if  $n$  is an integer. Identifying the phase angle  $\phi = n\pi$  with  $0 \leq n < 2$ , we obtain a unique

wave function of the universe

$$\Psi(a, \phi) = \begin{pmatrix} (1 + e^{i\phi})a^{\phi/\pi} \\ (1 + e^{-i\phi})a^{-\phi/\pi} \end{pmatrix}. \quad (13)$$

The quantum scalar curvature operator is then

$$\hat{V}(a, \phi) = \frac{\phi}{\pi a^2} \begin{pmatrix} \frac{\phi}{\pi} - 1 & 0 \\ 0 & \frac{\phi}{\pi} + 1 \end{pmatrix}. \quad (14)$$

The cosmological wave equation is unlike any other in the foundations of physics. It was solved without initial conditions. The initial data is unified with the first principle—the *form* of the laws of physics is the same on all scales, meaning the conformal zero that is the cosmological singularity is identified with future conformal infinity at  $\phi = \pi$ . It is of profound mathematical beauty that  $\Psi$  vanishes at  $\phi = \pi$  by Euler's identity,

$$\Psi(a, \pi) = \begin{pmatrix} (1 + e^{i\pi})a \\ (1 + e^{-i\pi})a^{-1} \end{pmatrix} = 0. \quad (15)$$

The quantum vacuum state  $\Psi$  is pure, meaning the entropy is zero:

$$Z = 1 = e^S \Rightarrow S = 0. \quad (16)$$

The contracting state is a perfect mirror image of the expanding state. There is only One state in the cosmological Hilbert space. *At  $t = 0$ , the universe is spatially flat and in a thermal state of zero entropy and zero energy! At  $t > 0$ , the dual dynamical theory with a spin-2 field is required.*

Conceptually, the next great challenge is moving between the timeless God's-eye view and the dynamic ant's-eye view. On a technical level, this reduces to the origin of the Born rule, the unitary evolution of the universal quantum state  $\Psi$ , and the nature of the quantum metric tensor.

### 1. The algebra of physical information

Let us begin our journey with symmetry. The global phase symmetry  $U(1)^2$  of  $\Psi$  is the maximal torus of  $U(2)$ . Multiplication of  $\Psi$  by

$$U(1)^2 = \begin{pmatrix} e^{-i\phi} & 0 \\ 0 & e^{i\phi} \end{pmatrix} \quad (17)$$

takes the components of  $\Psi$  to their complex conjugates  $\bar{\Psi}$  by a phase rotation. Unitary evolution is simply a phase rotation. Integrating over  $\mathcal{T}^2$  (see fig. 1) means integrating  $\Psi$  (with components living in the left-hand quadrants) and  $\bar{\Psi}$  (with components living in the right-hand quadrants). The Born rule follows from the double covering of conformal infinity. As it turns out, the double covering of fig. (1) is a pair of Clifford tori squeezed at a common point where past infinity, future infinity, and spacelike infinity,  $I_{L,R}^- = I_{L,R}^+ = I_{L,R}^0 = \Psi = 0$ , with a differentiable structure similar to the  $U(2)$  compactification of Minkowski space, visualized in fig. 8 of [52]. A key difference in the quantum picture of Minkowski space is the cyclic timelike direction of  $\mathcal{T}^2$  is imaginary and hence isomorphic to inverse temperature. The bond between time, causality, quantum information, and geometry is deeper still.

Another way of calculating  $\Psi$  is purely algebraically. Every point in  $\mathbb{R}^3$  has the following matrix representation:

$$Q = \sigma^1 x_1 + \sigma^2 x_2 + \sigma^3 x_3 \quad (18)$$

where the  $\sigma^i$  are the Pauli matrices. The eigenvalues are  $\pm r$  where  $r$  is the radius of curvature of the 2-sphere and  $\text{tr}(Q) = r + (-r) = 0$ . Identifying  $r \leftrightarrow a$  and using the spectral theorem to compute the function of a matrix, the left and right-handed states are computed as

$$\psi_L = \text{tr } Q^{\frac{\phi}{\pi}} = (1 + e^{i\phi})a^{\frac{\phi}{\pi}} \quad (19)$$

and

$$\psi_R = \text{tr } Q^{-\frac{\phi}{\pi}} = (1 + e^{-i\phi})a^{\frac{\phi}{\pi}} \quad (20)$$

where  $0 \leq \phi < 2\pi$ . One interprets  $Q$  as a qubit embedding of  $\mathbb{R}^3$  with an equivalence relation  $a \sim \lambda a$  for all real numbers  $\lambda \neq 0$  that identifies the 2-sphere (of any arbitrary size) as the Bloch sphere. One can always find a  $\lambda$  such that  $\lambda a$  has unit norm and identify the 2-sphere as the Bloch sphere. The physical interpretation is this: every point on the Bloch sphere is a light ray of the celestial sphere with characteristic wavelength  $\lambda$ . Thus, every point of  $\mathcal{T}^2$  is both a qubit and a celestial sphere. From the ant's-eye view, the cosmological Hilbert space is the configuration space of all possible celestial spheres that can surround the observer. From the God's-eye view, it is a *qubit continuum* (or Q-continuum) where every point of  $\mathcal{T}^2$  (except at  $\Psi = 0$ ) is a qubit.

### 2. The form of the quantum metric tensor

Recall what we learned from the geometry of quantum information: Minkowski space has a compactification  $\bar{M} = U(2)$  and maps to the celestial sphere as a pair of 2-spinors by the exceptional spin isomorphism  $Spin(3,1) \cong SL(2,\mathbb{C})$ . The quantum celestial (Riemann) sphere has a quantum metric tensor. Knowing the quantum scalar curvature operator  $\hat{V}(a, \phi)$  is enough to deduce the *form* of the quantum metric operator. The metric tensor  $g_{ij}$  for the  $n$ -sphere is an  $n \times n$  matrix. Since the topological dimension  $n$  is now a fractional quantity, the metric tensor is promoted to a continuous matrix, i.e., an integral kernel operator  $\hat{K}_g$  that acts on a 2-spinor:

$$(\hat{K}_g \psi)(i) \sim \int K_g(i, j) \psi(j) dj. \quad (21)$$

Recall that in QFT, Dirac 4-spinors are *reducible* representations that can always be decomposed into a pair of *irreducible* Weyl 2-spinors. The story here is a timeless boundary, a theory of a pair of irreducible 2-spinors with neither gravity nor observers, somehow dual to a dynamical theory of 4-spinors with gravity and observers. The form and role of this operator will become more precise when we derive the bulk dynamical theory. To do this, we must first understand the unitary evolution of the boundary theory.

### C. Cosmological Hilbert space of states

The simplest inner-product space over  $\mathcal{T}^2$  is reflection positive and chiral with a conformal horizon  $\mathcal{R} = \frac{\lambda_\infty}{\lambda_0}$  that places a fundamental limit on the number of qubits that are, in principle, observable. The cosmological Hilbert space of states  $\mathcal{H}(\mathcal{T}^2)$  is defined as:

$$\begin{aligned} \langle L|L \rangle &:= \int_0^{\lambda_\infty} \int_0^{2\pi} d\phi da \, \psi_L \bar{\psi}_L \\ &= 4\pi \int_0^{\lambda_\infty} da (a^4 - 1) \left( \frac{1}{\ln a} \cdot \frac{\pi^2 + 8 \ln^2 a}{\pi^2 + 4 \ln^2 a} \right) \\ &< \infty \end{aligned}$$

and

$$\begin{aligned} \langle R|R \rangle &:= \int_{\lambda_0}^{\infty} \int_0^{2\pi} d\phi da \, \psi_R \bar{\psi}_R \\ &= 4\pi \int_{\lambda_0}^{\infty} da \left( 1 - \frac{1}{a^4} \right) \left( \frac{1}{\ln a} \cdot \frac{\pi^2 + 8 \ln^2 a}{\pi^2 + 4 \ln^2 a} \right) \\ &< \infty, \end{aligned}$$

where the "cut-offs"  $\lambda_0$  and  $\lambda_\infty$  are the Compton wavelengths of the masses  $m_0$  and  $m_\infty$ , respectively. It is clear  $\lambda_0$  is the Planck length and  $m_0$  is the Planck mass, the mass of the smallest possible black hole. Both  $m_0$  and  $m_\infty$  are therefore built from fundamental constants. In

gravity, the charge is mass. There is a global  $U(1)^2$  phase symmetry, meaning the vacuum has a unique global charge  $m$ . A massive global charge does not break the conformal symmetry. The cosmological wave equation is 2nd-order with dimensions of the Schrödinger equation and vanishing total energy. The charge  $m$  is conformal. One can write down any arbitrary mass matrix  $M = \text{diag}(m_0, m_\infty)$  and multiply eq. (6) by  $M^{-1}$  on both sides such that  $\mathcal{R}$  is conserved. Only dimensionless constants are fundamental—the laws of physics do not depend on the choice of unit system in a theory with no fundamental scale.

Integrating over all possible phases (all possible paths), one obtains the *world densities*  $\rho_L(a)$  and  $\rho_R(a)$ , which are perfect mirror images of each other. Why define the universal wave function as an element of a normed Hilbert space? Because there is no supernatural observer that exists outside of the universe bestowed with the decision to choose the initial conditions of the universe. All initial conditions and, hence, all paths consistent with the symmetries of the system are allowed. There is no Ptolemaic center in the Hilbert space of states—no first cause. This is the origin of the “many worlds” and why quantum mechanics has the structure that it does. Quantum mechanics is the ultimate realization of the Copernican principle.

#### D. The Bang

The charge  $m_0 = M_{pl}$  can not be fundamental as it radiates away with Hawking temperature  $T \sim g/2\pi$ , where  $g$  is the surface gravity of the horizon at  $r = \lambda_0 = L_{pl}$ . Another way to see why this is true is from  $\hat{V}$  (see eq. 14). The Planck mass  $m_0$  sources the left-hand potential  $V_L$  which has a decaying attractive phase for  $0 < \phi < \pi$  and is zero at  $\phi = \pi$ , meaning  $V_L$  decays into the right-hand potential  $V_R$ . Thus, one concludes  $m_0$  is not fundamental and must be a composite of  $m_\infty$ . The right-hand potential  $V_R$  is always repulsive, with  $m_\infty$  sourcing positive quantum curvature that energetically favors an inflationary-like event and then slowly rolls like  $a^{-2}$  to zero at future infinity.

The ultra-light gravitational charge determines the largest scale the observer shall ever see. Future infinity has a horizon at  $\lambda_\infty$ , explaining why the charge  $m_\infty$  is small. (A calculation of this number will appear in future work.) Nature has no fundamental scale, but the observer does. To ensure eq. (6) remains dimensionless send  $a \rightarrow \frac{a}{\lambda_0}$  and  $a \rightarrow \frac{a}{\lambda_\infty}$  in  $\psi_L$  and  $\psi_R$ , respectively. A remarkable duality then follows:

$$\psi_L(\lambda_\infty, \phi) \bar{\psi}_L(\lambda_\infty, \phi) = \psi_R(\lambda_0, \phi) \bar{\psi}_R(\lambda_0, \phi). \quad (22)$$

The UV horizon at  $\lambda_0$  and the IR horizon at  $\lambda_\infty$  are dual. UV/IR mixing is required for unitarity! Unitary evolution is a phase rotation of  $\psi_L$  from  $a = 0_L$  through  $a = \lambda_\infty^L$  to  $a = \lambda_0^R$  (still labeled as  $\psi_L$ ) to  $a = 0_R$ . The UV and IR scales place a fundamental limit on what is,

in principle, observable. From the ant’s-eye view, the observer can access at most  $\mathcal{R}$  qubits (modulo some constant). Observers are finite-dimensional Hilbert spaces embedded in  $\mathcal{H}(\mathcal{T}^2)$ .

The wave function of the universe is pure, meaning the entropy, as witnessed from the God’s-eye perspective at  $t = 0$ , is zero. For  $t > 0$ , the entropy, as witnessed by bulk observers, is increasing. Recall the conformal Clifford torus (fig. 1) has one dimension suppressed, the spinor components  $\psi_L, \psi_R \in L^2(S_a^2 \times S_\beta^1)$ . One can express the world densities  $\rho_L(a)$  and  $\rho_R(a)$  as a function of the surface area of the 2-sphere with the scale transformation  $a \rightarrow \sqrt{\pi}a$ :

$$\rho_L(A) = 4\pi^2 \int_0^{2\pi} \cos^2\left(\frac{\phi}{2}\right) \left(\frac{s^2 A}{\lambda_0^2}\right)^{\phi/\pi} d\phi \quad (23)$$

and

$$\rho_R(A) = 4\pi^2 \int_0^{2\pi} \cos^2\left(\frac{\phi}{2}\right) \left(\frac{s^2 A}{\lambda_\infty^2}\right)^{-\phi/\pi} d\phi. \quad (24)$$

such that  $A = 4\pi a^2$  and the spin  $s = \pm 1/2$ . One can further re-express the above as functions of the Bekenstein-Hawking entropy (the entropy from the ant’s-eye perspective) by the identification:

$$S_\alpha = \frac{s^2 A}{\lambda_\alpha^2} = \frac{A}{4\lambda_\alpha^2} \quad (25)$$

with the conformal indices  $\alpha = 0, \infty$ . ( $\alpha = \infty$  is the  $\mathcal{R} \rightarrow 1/\mathcal{R}$  dual.) The world densities are thus functions of the black hole entropy. The charge  $m_0$  is a quantum black hole! If it seems odd that this can be done, it should not. The black hole entropy is a geometric consequence of integrating over a boundary with an  $S^2 \times S_\beta^1$  topology in Hawking and Gibbon’s original paper [91]. I provide the de Sitter limit from the boundary in appendix C.

These deductions lead to a remarkable prediction. The vacuum has a conserved charge  $m_\infty$ , meaning the mass density  $\rho_\infty = m_\infty/\lambda_0^3$  of empty space is a conserved quantity. But if  $m_0$  is a composite of  $m_\infty$ , the black hole mass function must violate decoupling and scale like  $a^3$  since mass density dilutes like  $a^{-3}$  in an expanding Euclidean 3-space. This means two competing mechanisms set up the Bang: 1) the decay of  $m_0$  into  $m_\infty$  and 2) the inflation of  $m_0$  into larger black holes as space accelerates in all directions. The ‘moment before time’ is a perfect lattice of quantum black holes obeying the Pauli exclusion principle. From the uncertainty principle, the quantum black holes do not all decay into  $m_\infty$  at the same rate. There will always be a fraction that swells to astrophysical sizes, seeding structure formation in the early universe. The spectrum of these black holes serves as the initial conditions of the bulk field equations. To rewind the universe back to  $t = 0$  and hit play, we must derive the local dynamical equations of motion.

### E. The dynamic bulk

For the remainder of the paper, I will use the mostly minus metric signature  $(+ - - -)$  and natural units  $c = G = \hbar = 1$ . Greek indices  $\mu, \nu = 0, 1, 2, 3$  index spacetime, lowercase Latin indices  $i, j = 1, 2, 3$  index space, and upper case  $I, J = 0, 1$  index 2-spinors in the fundamental representation. I use a 4-spinor notation where  $\psi^\mu$  denotes a column 4-spinor and  $\psi_\mu$  the conjugate transpose. One lower and upper repeated index implies summation.

The boundary equations of motion, eq. (6), are defined by the Clifford algebra  $\mathcal{C}\ell_3(\mathbb{R})$ . It follows from the exceptional spin isomorphism that the bulk equations of motion are defined by the Dirac algebra  $\mathcal{C}\ell_{3,1}(\mathbb{C})$ , with the simplest holographic dictionary given by

$$D \leftrightarrow \not{D}, \quad \hat{V} \leftrightarrow \hat{V}_{\mu\nu} \quad (26)$$

where  $\not{D} = i\gamma^\mu \partial_\mu$  is the Dirac operator and the potential  $V_{\mu\nu}$  is a second-rank tensor. The square of the Dirac operator  $\not{D}^2 = \partial^\mu \partial_\mu I_4$ , and the bulk Hamiltonian becomes

$$\hat{H}_{\mu\nu} = -\nabla^2 I_4 + V_{\mu\nu}. \quad (27)$$

One can then write down the

dynamical cosmological wave equation

$$\hat{H}_{\mu\nu} \psi^\nu = E^2 \psi^\nu \quad (28)$$

$$(-\nabla^2 I_4 + V_{\mu\nu}) \psi^\nu = -\partial_t^2 \psi^\nu \quad (29)$$

$$\partial^\mu \partial_\mu \psi^\nu + V_{\mu\nu} \psi^\nu = 0. \quad (30)$$

The bulk equations of motion are four coupled relativistic wave equations second-order in time. Fields live in representations of the Lorentz group. The potential  $V_{\mu\nu}$  decomposes into the following classical irreducible representations of the Lorentz group: an antisymmetric tensor  $F_{\mu\nu} = -F_{\nu\mu}$ , a symmetric ( $H_{\mu\nu} = H_{\nu\mu}$ ) and traceless ( $H^\mu_\mu = 0$ ) tensor  $H_{\mu\nu}$  (not to be confused with the Hamiltonian), and a scalar trace component:

$$V_{\mu\nu} = F_{\mu\nu} + H_{\mu\nu} + X^\mu_\mu. \quad (31)$$

Under the appropriate local phase (gauge) invariance conditions,  $H_{\mu\nu}$  is a massless spin-2 field—the graviton! If we set the scalar trace component  $X^\mu_\mu = 1/\lambda_C^2$  where  $\lambda_C$  is the Compton wavelength and introduce some dimensionful coupling constants, the bulk equations of motion become:

$$\partial^\mu \partial_\mu \psi^\nu + g_{\text{EM}} F_{\mu\nu} \psi^\nu + g_G H_{\mu\nu} \psi^\nu + m^2 \psi^\nu = 0. \quad (32)$$

In the limit  $g_{\text{EM}} \rightarrow g_G \rightarrow 0$  each component of  $\psi^\nu$  reduce to solutions of the Klein-Gordon equation. There is no imaginary number  $i$  in the above equation, meaning  $\psi^\nu$  can either be real-valued and transform like a 4-vector or complex-valued and transform like a 4-spinor.

The timeless boundary severely restricts the dynamic bulk. A pair of 2-spinors defines the boundary theory, so the above wave equation must decompose into a pair of 2-spinor wave equations in the reducible or fundamental representation.

Recall that the Hopf bundle has  $U(2)$ , a compact representation of Minkowski space, as its symmetry group. Locally  $U(2)$  is diffeomorphic to the direct product  $G = SU(2) \times U(1)$ . Up to a local diffeomorphism, the compact gauge group  $G$ , together with the emergence of the antisymmetric tensor  $F = F_{\mu\nu}$ , implies the existence of Yang-Mills equations, with a gauge connection  $A$  on  $G$ . The connection  $A$  is locally a one-form on spacetime, and the curvature or field strength tensor is the two-form:  $F = dA + A \wedge A$ . It follows the antisymmetric component of  $V_{\mu\nu}$  can be expanded into the electroweak field strength tensors (pre spontaneous symmetry breaking) with  $U(1)$  and  $SU(2)$  gauge connections  $A_\mu$  and  $\{W_\mu^i\}_{i=1,2,3}$ , respectively.

Generally speaking, mixing spinor and tensor indices always results in a conflict of notation. Keep in mind when we work with 4-spinors, we do so in the reducible representation, meaning when we move to the fundamental representation (a pair of 2-spinors with the Pauli algebra representation of the field strength tensors), we no longer have the issue of mixing tensor and spinor indices. The limitation is in formalism only—not the physics.

I show in appendix [A] if one restricts themselves to the electromagnetic sector with an agreed-upon index notation, it is not hard to show  $F_{\mu\nu} \psi^\nu \rightarrow e^{i\phi} F_{\mu\nu} \psi^\nu$  under a local  $U(1)$  gauge transformation with the covariant derivative.

### F. Quantum gravity

In semiclassical gravity, the simplest metric is Minkowski, with some quantum fluctuations:

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad (33)$$

where  $h_{\mu\nu}$  ( $\equiv H_{\mu\nu}$ ) is a symmetric and traceless rank-2 tensor. From eq. (21), it was deduced that the quantum metric tensor is an integral kernel operator; hence, so too is the spin-2 field as it appears in the bulk equations of motion. A natural choice is then:

$$H_{\mu\nu} \rightarrow H = \int d\psi_\mu h(\psi^\mu, \psi_\nu) \quad (34)$$

such that the kernel is a complex symmetric

$$h(\psi^\mu, \psi_\nu) = \overline{h(\psi_\nu, \psi^\mu)} := h(\psi^\nu, \psi_\mu), \quad (35)$$

traceless trace-class operator

$$\text{tr } H = \int d\psi_\mu h(\psi^\mu, \psi_\mu) = 0. \quad (36)$$



The classical irreducible representation of the Lorentz group,  $H_{\mu\nu}$ , is promoted to a unique class of Hilbert-Schmidt integral operators where

$$\int_{\Omega} \int_{\Omega} d\psi_{\mu} |h(\psi^{\mu}, \psi_{\nu})|^2 d\psi^{\nu} < \infty \quad (\mu \neq \nu) \quad (37)$$

with the domain  $\Omega$  determined from the boundary theory. Gravity is exceptional in that it acts on the quantum state as an integral operator, whereas the other forces act on the quantum state as differential operators. The classical irrep.  $H_{\mu\nu}$  of the Lorentz group is fully quantized and makes eq. (28) a highly non-linear partial differential equation, where  $H\psi^{\nu} = U(\psi^{\nu})$  is a quantum gravitational potential and self-interaction term. I explicitly calculate the (manifestly gauge-invariant) spin-2 field operator acting on the quantum state in the fundamental representation in appendix B 2.

The 4-spinor  $\psi^{\nu}$  in eq. (28) decomposes into a pair of 2-spinors  $\chi_L$  and  $\chi_R$ , respectively, that satisfies two coupled non-linear relativistic wave equations

$$(-\sigma^{\mu}\sigma^{\nu}\partial_{\mu}\partial_{\nu}I_2 + V)\chi_L = 0 \quad (38)$$

$$(-\bar{\sigma}^{\mu}\bar{\sigma}^{\nu}\partial_{\mu}\partial_{\nu}I_2 + V^{\dagger})\chi_R = 0 \quad (39)$$

where  $\sigma^{\mu} = (I_2, \sigma^i)$ ,  $\sigma^{\nu} = (-I_2, \sigma^j)$ ,  $\bar{\sigma}^{\mu} = (I_2, -\sigma^i)$ , and  $\bar{\sigma}^{\nu} = (-I_2, -\sigma^j)$ , i.e., the generators of  $U(2) \times \mathbb{Z}_2 \times \mathbb{Z}_2$ , and the potential  $V = V_{IJ}$  contains the same information as  $V_{\mu\nu}$  by recasting the field strength tensors in the Pauli matrix representation with the Clifford algebra  $\mathcal{Cl}_{3,0}(\mathbb{R})$ , otherwise known the Pauli algebra. Recall what we learned from the geometry of quantum information: Minkowski space has a compact presentation  $U(2)$ , and the fundamental representation of the Lorentz group contains two fermionic directions. Unpacking the above, we have,

$$(\partial^{\mu}\partial_{\mu}I_2 + F + H + m^2)\chi_L = 0 \quad (40)$$

$$(\partial^{\mu}\partial_{\mu}I_2 + F^{\dagger} + H^{\dagger} + m^2)\chi_R = 0 \quad (41)$$

with the dimensionful coupling constants  $g_{EM}$  and  $g_G$  absorbed in  $F$ ,  $H$ , and their conjugate transposes, respectively. In the limit  $g_{EM} \rightarrow g_G \rightarrow 0$ , the above reduces to the Klein-Gordon equation. Thus, one concludes equation (28) is the adjoint representation where the above is the fundamental representation. I explicitly write down the bulk field equations in the fundamental representation in appendix [B].

In the scale basis and in the limit  $V_{\mu\nu} \rightarrow 0$ , the identification  $a \leftrightarrow \sigma$  (proper space) and  $t \leftrightarrow \tau$  (proper time) of the string worldsheet, the fundamental representation reduces to the equations of motion of the bosonic string in conformal gauge. The implied supersymmetry here is of a more fundamental nature than string theory without quantum covariance. All bosonic fields (including Minkowski space as a spin-1 field) are dual to a pair

of spinors on the timeless boundary. The fundamental representation *is* the boundary at future infinity. This explains why the LHC has not seen superpartners: they represent redundant unphysical degrees of freedom that only appear in perturbative constructions without quantum covariance.

The first step in demonstrating a holographic duality between the timeless boundary and the dynamic bulk is to match the energy spectrum of the bulk and boundary theories. The boundary theory is based on the energy  $E$ , which is zero, whereas the bulk is based on the squared energy  $E^2$ , which is non-zero and has a negative energy solution. (Cosmologically, this implies a CPT-symmetric universe consistent with the proposal of Turok & Boyle [29].) The conformal invariance of the boundary is locally broken from the ant's-eye perspective of the bulk observer. So, how can it be that these two theories are secretly equivalent? How is a dynamical non-linear differential equation secretly a non-dynamical linear one?

### III. CAUSALITY & THE NATURE OF THE OBSERVER

The 'many worlds' in Everettian QM are causally disconnected when a classical global time coordinate is inserted by hand. It is clear that if time is not fundamental, the classical notions of causality break down. When time drops out of the Schrödinger equation, the Hilbert space becomes causally indefinite. The conformal Clifford torus has no causal ordering—light cones can be superimposed. Remarkably, in the first of its kind, the quantum switch experiment has demonstrated *causal indefiniteness* [63, 64]. Since there is no global time coordinate, there must exist a *causal reference frame* [65–68] for each event that establishes an observer-dependent time to describe the evolution of quantum subsystems.

#### 1. Holographic transform as the quantum metric operator

The Dirac field can be written as a sum of chiral Weyl fields. The boundary theory has no timelike direction, so its solutions obviously can not be a linear combination of non-linear bulk states. There must exist an integral transformation, like a Fourier transform, that maps bulk states to boundary states:

$$\Psi(a, \phi) = \int_{-\infty}^{\infty} g_{IJ} \chi_L^J dt + \int_{-\infty}^{\infty} g_{IJ}^{\dagger} \bar{\chi}_R^J dt, \quad (42)$$

where the quantum metric tensor  $g_{IJ}$  is a  $2 \times 2$  Hermitian matrix of kernels, and the fields are in the scale basis  $\chi^J = \chi^J(a, t)$  for simplicity. The quantum metric tensor establishes a causal reference frame for bulk observers that, when integrated over all time, acts as an atlas for locating bulk events on the boundary. The key to understanding the above transformation is the mechanism

responsible for the *illusion* that local conformal symmetry is broken and the connection between mass and time. The above setup can also be seen as a field-theoretic generalization of the Page-Wooters mechanism [40].

A central principle underlying GR is the observer's freedom to choose their coordinate frame. No privileged observer has the authority to fix a preferred coordinate frame for all observers. Weyl [69] and Dicke [70] concluded the principle of general covariance, that the *form* of the laws of physics do not depend on the choice of coordinates, should also apply to the choice of units. A fundamental symmetry *not* present in Einstein's GR. When Alice chooses her clock and decrees, "This is one second!" the unit "one second" is defined globally. Neither Bob nor Chester are free to choose a clock with any natural frequency other than one second. The notion of a *universal democracy* is violated.

Indeed, as the laws of physics are currently formulated, the electron is defined to have the same mass or, equivalently, the same Compton frequency at every spacetime point, independent of the local configuration of the spacetime. But if the form of the laws of physics does not depend on the choice of units, the notion of absolute mass is meaningless—it is only the dimensionless ratios of masses that are well-defined. The vacuum is charged with  $m_\infty$ , so it is the ratio  $m_e/m_\infty$  at every spacetime point that is physically meaningful.

The boundary theory is dimensionless. The observer can choose any reference scale to attach to their local coordinate frame. The reference scale can be defined as a massive scalar field

$$\varphi(a, t) = e^{-i\omega\tau} \Phi(a) \quad (43)$$

where  $\omega = mc^2/\hbar$  is the Compton frequency of the mass reference  $m$  at rest, the proper time  $\tau = t/\gamma$  where  $t$  and  $\gamma$  are the laboratory time and Lorentz factor, respectively; and

$$\Phi(a) = \Phi_0(a) + i\Phi_1(a) \quad (44)$$

is a plane wave solution to the KG equation that assigns a complex scalar value to each scale of the universe. One can then make the following ansatz:

$$g_{IJ} = e^{-i\omega\tau} g_{IJ}[\Phi(a), \bar{\Phi}(\beta)] \quad (45)$$

$$= e^{\omega\beta} g_{IJ}[\Phi(a), \bar{\Phi}(\beta)] \quad (46)$$

where it is forced upon us to identify  $\beta = -i\tau$  for eq. (42) to converge. The oscillations of the wave packet of  $m$  accumulate a phase  $\omega\tau$  just as a clock does following the same trajectory [71]. The observer is free to choose any reference mass  $m_\infty < m < m_0$  as a Compton clock such that  $\mathcal{R} = m_0/m_\infty$  is a conserved quantity all observers agree on. One can further make the ansatz that the global phase of the boundary theory

$$\phi(\omega/\omega_\infty) \sim i \int_{-\infty}^{\infty} e^{(\omega/\omega_\infty)\beta} \bar{\Phi}(\beta) d\beta \quad (47)$$

is essentially a Fourier transform of complex-valued massive scalar fields along the thermal circle at conformal infinity to a real-valued dimensionless scalar field in the frequency domain.

Gauge transformations of the spinor fields  $\chi \rightarrow e^{i\vartheta(a,\tau)} \chi$  are equivalently conformal transformations of the metric field,

$$g_{IJ} \rightarrow e^{i\vartheta(a,\tau)} g_{IJ} \quad (48)$$

where the rescaling of the metric is the accumulation of the local phase  $\vartheta(a, \tau)$  that corresponds to the global phase shift  $\phi \rightarrow \phi'$  on the boundary:

$$\Psi(a, \phi') = i \int_{-\infty}^{\infty} e^{\omega\beta + i\vartheta(a,\beta)} g_{IJ} \chi_L^J d\beta \quad (49)$$

$$+ i \int_{-\infty}^{\infty} e^{\omega\beta - i\vartheta(a,\beta)} g_{IJ}^\dagger \bar{\chi}_R^J d\beta. \quad (50)$$

Local phase transformations in the bulk are global phase transformations on the boundary. The phase  $\phi$  of the universal wave function is coupled to the potential (quantum scalar) curvature (eq. 14) and encoded in the quantum metric tensor  $g_{IJ}$ . The cosmological Hilbert space, the space of all possible configurations of the celestial sphere, is defined as an integral over  $\phi$  and is thus a path integral over all possible histories (for all possible initial conditions). Moreover, the global phase is *relative* and therefore, in principle, observable!

Consider the following thought experiment. Suppose Alice collects photons from the surface of last scattering (the Cosmic Microwave Background or CMB) from our solar system while Bob does the same in the Zeta Reticuli system. Analogous to the Aharonov-Bohm effect, they pick up an observer-dependent local phase  $\vartheta_A(l, b, \tau)$  and  $\vartheta_B(l, b, \tau)$  in galactic coordinates as they rotate around Sagittarius A\* at the center of the galaxy. This means when Alice and Bob compare the photons from their celestial spheres, there will be observer-dependent correlations in the surface of last scattering proportional to the phase change:

$$\Delta\phi = \phi_B - \phi_A. \quad (51)$$

Before the measurements, Alice and Bob's past light cones were superimposed. When the photons interact with their instruments, they become entangled, not in space, but in time. When they compare the amplitudes  $a_{\ell m}$  of the angular oscillations of the CMB temperature, they find their CMB maps do not agree: from their frame of reference, they observe correlations between  $a_{\ell m}$  and their motion around Sagittarius A\*. Alice and Bob conclude their photons do not originate from the same surface of last scattering, their past light cones have "collapsed" into two adjacent worlds! Alice and Bob exist at the center of two causally indefinite overlapping universes a distance:

$$d(\Psi_A, \Psi_B) = \|\Psi(a, \phi_A) - \Psi(a, \phi_B)\| \quad (52)$$

apart in the configuration space of all possible celestial spheres. *Bob has traversed the multiverse to return to Alice!*

In quantum field theory, we are taught the gauge degrees of freedom are redundant. The internal symmetries are an artifact of our description of nature and not physical. But this can not be so in a holographic theory of time. The gauge degrees of freedom connect the coordinate-dependent phase  $\varphi$  in the bulk to a coordinate-independent phase  $\phi$  on the boundary that encodes the initial data of an arbitrary Bang. No unphysical axes exist in the configuration space of the boundary theory. Quantum mechanics is the Copernican principle of the configuration space: there is no preferred axis (a Ptolemaic center) in the cosmological Hilbert space. A gauge transformation (when appropriately gauge fixed with a casual reference frame) acts as a shear along these extra dimensions.

## 2. Is the sky quantum?

Is there any evidence of observer-dependent correlations in cosmic microwave background (CMB)? Indeed, there are. The most puzzling observations are the large-angular-scale anomalies in the CMB temperature. Among the most compelling are the lack of large-angle CMB temperature correlations, hemispherical power asymmetry, octopole planarity and alignment with the quadrupole, the point-parity anomaly, and variations in cosmological parameters over the sky. With no common systematics, WMAP and *Planck* are in remarkable agreement that these are real features of the sky. Individually, the significance of these anomalies ranges from  $2\sigma - 4\sigma$ . Since many of these anomalies are statistically independent, collectively, they are highly significant and exceed  $5\sigma$  [72]!

The most bewildering anomaly is the octopole planarity and alignment with the quadrupole.  $\Lambda$ CDM asserts the  $a_{\ell m}$  are statistically independent random variables. The data says otherwise:  $a_{2m}$  and  $a_{3m}$  show anomalous correlations [73]. More surprising still, the great circle defined by the shared plane of the quadrupole and octopole is nearly perpendicular to the ecliptic—the orbital plane of the Earth around the Sun. This is the infamous ‘axis of evil.’

Both WMAP and *Planck* orbited in the ecliptic and observed nearly identical octopoles and quadrupoles even though their sampling strategies and noise maps are “ecliptic aware.” The fact that the quadrupole and octopole are planar and perpendicular to the ecliptic is fundamentally unlike anything expected from a solar system foreground. Within the framework of  $\Lambda$ CDM there should not be any correlations between the motion of these probes and the light emitted from the surface of last scattering 380,000 years after the Bang. It is not an overstatement to say this is the most profound anomaly in all of science. Either the Copernican principle is wrong, and

there exists a classical supernatural observer outside of the universe that selected the initial conditions for the Earth to be at the center of the universe, or the sky is quantum. The Copernican principle is promoted to the Hilbert space of states. The multiverse is traversable.

Indeed, our thought experiment need not be done at such great distances nor be confined to the realms of theory. WMAP and *Planck* orbit in the Earth-Sun L2 point in the ecliptic. The closest celestial body with the greatest deviation from the ecliptic is Pallas, the third-largest asteroid in the solar system, with an orbital inclination of 34.8 degrees. A Palladian mission to map the CMB should definitively tell us whether or not there are observer-dependent features of the quantum sky. If successful, this mission would represent a key milestone in further understanding the holographic transform, the quantum metric tensor, and the nature of causal indefiniteness. The prerequisites to exploring the multiverse.

## 3. Implications of a traversable multiverse

The local rescaling of the quantum metric  $g_{IJ} \rightarrow e^{i\vartheta(a,\tau)} g_{IJ}$  is the accumulation of a local phase that rotates the observer’s causal reference frame into an adjacent world with adjacent initial conditions. From the constraints on the spin-2 integral operator, the simplest spin-2 field configuration is

$$h(\chi^I, \chi_J) = \begin{pmatrix} 0 & \chi_0 \bar{\chi}_1 \\ \chi_1 \bar{\chi}_0 & 0 \end{pmatrix}, \quad (53)$$

with the quantum gravitational potential given by a pair of left-handed and right-handed spinor potentials:

$$H\chi^J = \int d\chi_I h(\chi^I, \chi_J) \chi^J = \frac{3g_G}{8\sqrt{2}\mathcal{R}^3} |\chi_0|^2 |\chi_1|^2. \quad (54)$$

See appendix B 2 for the detailed derivation. The gravitational potential is the potential of a harmonic oscillator in the spinor degrees of freedom! The above is manifestly gauge-invariant. It is also gauge-independent:  $H\chi^J \rightarrow H\chi^J$  under a local rescaling of the metric. This means the gravitational degrees of freedom are observer-independent for observers confined to the same quantum gravitational potential. Alice and Bob always agree on the Gravitational Wave Background (GWB). The above, however, is not a unique spin-2 field configuration. Other spin-2 field configurations exist, the landscape of which is unknown.

If quantum theory is truly universal and if UV/IR mixing is a general feature of the quantum universe, what are the implications of causal indefiniteness? Can exotic UV/IR couplings that violate EFT reasoning enable the quantum metric tensor to be technologically exploited? Looking at equations (38) - (42), it is tantalizingly possible to engineer the quantum metric tensor with some highly tuned matter-antimatter reaction. The solution space of the bulk field equations has the energy quartet  $(\pm E, \pm E)$ , which is nothing but the  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetry.

Herein lies a great clue: the boundary theory violates decoupling, so bulk solutions must exist where the UV mixes with the IR—exotic scenarios where quantum distortions of spacetime may be observable from a two-fermion-two-antifermion interaction. The future of particle physics may, therefore, not be in building larger circular colliders, but smaller compact colliders that bring matter and antimatter together deep in the IR at low accelerations. Frequency gravimeters (a kind of Compton clock) are uniquely equipped to detect distortions in spacetime [74].

Is the final frontier of fundamental physics the use of cosmological observations to fine-tune particle physics experiments? Should we be successful in the Palladian mission to map the CMB, find the sky is quantum, and detect quantum fluctuations of spacetime from matter-antimatter annihilation events, what are the implications? This begs the profound question: can we manipulate the quantum metric with the precision we manipulate electrons and warp spacetime into shapes of our design? And if so, will humankind one day, in other worlds out there, find their way to us? Will we one day find our way to them? If quantum universality is true and the multiverse exists, there must exist worlds where future humans master the quantum nature of spacetime and travel to "past-adjacent" light cones—a form of time travel in the multiverse. But if this were possible, then where are they?

For most of human history, we have been a deeply divided species. So much so that we have developed nuclear arsenals that may annihilate us all. Unfortunately, it seems we are quickly approaching this precipice point. But if the deepest truth of quantum mechanics is that we are all One, connected in profound and mysterious ways that we have yet to understand, there is hope for a future beyond our wildest imagination. A future where science and spirituality need not be separate. A civilization of humankind cocreating as One people—in the liminal spaces, between worlds—past, present, and future.

## ACKNOWLEDGMENTS

Thank you, Niayesh Afshordi, for the insightful conversations, words of encouragement, and advocacy. Thank you, Will Kinney, Cliff Burgess, Thomas Van Riet, for the conversations contributing to this work. Thank you, David Peak, for the many correspondences on this work over the years, the helpful feedback, and, most importantly, the words of encouragement. Thank you, Neil Turok, for many of your lectures and ideas that have deeply inspired me. Thank you, Daniel Davis, for your continued support. Thank you, Sabine Hossenfelder, for writing your pellucid book, *Lost in Math: How Beauty*

*Leads Physics Astray*, which course-corrected this research program to what it is today. Lastly, thank you Jodi Benowitz and Ethan Benowitz for always being there when I needed you.

## Appendix A: Gauging the bulk equations of motion

### 1. Notation

I use the mostly minus metric signature  $(+---)$  with the standard Einstein notation and natural units  $\hbar = c = G = 1$ . Greek indices index spacetime, lowercase Latin indices index space, and uppercase Latin indices index 2-spinors. One lower and upper repeated index implies summation.

Recall that  $\psi^\nu$  can transform as either a 4-vector or a 4-spinor. Mixing spinor and tensor indices is messy, but it can be done. I use the following convention: for a covariant 2-form multiplied by a 4-spinor,

$$F_{\mu\nu}\psi^\nu \longrightarrow \psi^\nu = \begin{pmatrix} \psi^0 \\ \psi^1 \\ \psi^2 \\ \psi^3 \end{pmatrix} \quad \psi_\nu = (\bar{\psi}_0 \quad \bar{\psi}_1 \quad \bar{\psi}_2 \quad \bar{\psi}_3)$$

where lowering the index on the 4-spinor denotes the conjugate transpose. (The same index convention is used for 2-spinors.) For a contravariant 2-form multiplied by a 4-spinor, the convention is inverted. Note the difference between raising and lowering indices of a 4-spinor and 4-vector. Generally, this is a problem as conflicts of notation will arise. However, for what we are trying to calculate, so long as we are consistent in the notation, all is well.

When propagating the product rule, one runs into the following conflict:

$$\partial_\mu A_\nu \psi^\nu \stackrel{?}{=} A_\nu \partial_\mu \psi^\nu + \psi^\nu \partial_\mu A_\nu \quad (\text{A1})$$

$$\equiv A_\nu \partial_\mu \psi^\nu + (\partial_\mu A_\nu) \psi^\nu \quad (\text{A2})$$

The second term on the first line conflicts with our 4-spinor notation. Left multiplication should have a lowered index denoting the transpose for a row 4-spinor. However, we do not want to conjugate the elements of the 4-spinor or raise the index on the 4-vector for consistency with Einstein notation. Instead, we can use parentheses to denote the product rule has already been applied with right-multiplication of  $\psi^\nu$ . To make the calculations easier on the eyes, I will adopt the following convention,

$$\psi^\nu \partial_\mu A_\nu \equiv (\partial_\mu A_\nu) \psi^\nu \quad (\text{A3})$$

where left-multiplication of  $\psi^\nu$  followed by two lowered indices denotes the product rule has already been applied and to move  $\psi^\nu$  to the right for contraction of the repeated index.

## 2. Adjoint representation

Let's ignore the  $H_{\mu\nu}$  term in eq. (32) and focus on the electromagnetic field strength tensor:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu, \quad (\text{A4})$$

where the four-potential  $A_\mu = (A_0, -A_1, -A_2, -A_3)$ . The field strength tensor is antisymmetric, so  $F_{\mu\mu} = 0$ , meaning there are only six independent degrees of freedom. Computing each component yields:

$$F_{01} = -\frac{\partial A_1}{\partial x_0} - \frac{\partial A_0}{\partial x_1} = E_1 \quad (\text{A5})$$

$$F_{02} = -\frac{\partial A_2}{\partial x_0} - \frac{\partial A_0}{\partial x_2} = E_2 \quad (\text{A6})$$

$$F_{03} = -\frac{\partial A_3}{\partial x_0} - \frac{\partial A_0}{\partial x_3} = E_3 \quad (\text{A7})$$

$$F_{12} = -\frac{\partial A_2}{\partial x_1} + \frac{\partial A_1}{\partial x_2} = -B_3 \quad (\text{A8})$$

$$F_{13} = \frac{\partial A_1}{\partial x_3} - \frac{\partial A_3}{\partial x_1} = B_2 \quad (\text{A9})$$

$$F_{23} = -\frac{\partial A_3}{\partial x_2} + \frac{\partial A_2}{\partial x_3} = -B_1 \quad (\text{A10})$$

Taking the scalar trace component  $X_\mu^\mu = m^2$  and inserting  $F_{\mu\nu}$  explicitly in equation eq. (32) gives:

$$\square \begin{pmatrix} \psi^0 \\ \psi^1 \\ \psi^2 \\ \psi^3 \end{pmatrix} + g_{\text{EM}} \begin{pmatrix} 0 & \frac{\partial A_1}{\partial x_0} & \frac{\partial A_2}{\partial x_0} & \frac{\partial A_3}{\partial x_0} \\ -\frac{\partial A_1}{\partial x_0} & 0 & -\frac{\partial A_2}{\partial x_1} & \frac{\partial A_1}{\partial x_2} \\ -\frac{\partial A_2}{\partial x_0} & \frac{\partial A_2}{\partial x_1} & 0 & -\frac{\partial A_3}{\partial x_2} \\ -\frac{\partial A_3}{\partial x_0} & -\frac{\partial A_1}{\partial x_3} & \frac{\partial A_3}{\partial x_2} & 0 \end{pmatrix} \begin{pmatrix} \psi^0 \\ \psi^1 \\ \psi^2 \\ \psi^3 \end{pmatrix} + g_{\text{EM}} \begin{pmatrix} 0 & \frac{\partial A_0}{\partial x_1} & \frac{\partial A_0}{\partial x_2} & \frac{\partial A_0}{\partial x_3} \\ -\frac{\partial A_0}{\partial x_1} & 0 & \frac{\partial A_1}{\partial x_2} & -\frac{\partial A_3}{\partial x_1} \\ -\frac{\partial A_0}{\partial x_2} & -\frac{\partial A_1}{\partial x_2} & 0 & \frac{\partial A_2}{\partial x_1} \\ -\frac{\partial A_0}{\partial x_3} & \frac{\partial A_3}{\partial x_1} & -\frac{\partial A_2}{\partial x_3} & 0 \end{pmatrix} \begin{pmatrix} \psi^0 \\ \psi^1 \\ \psi^2 \\ \psi^3 \end{pmatrix} + m^2 \begin{pmatrix} \psi^0 \\ \psi^1 \\ \psi^2 \\ \psi^3 \end{pmatrix} = 0.$$

Good. Four KG equations coupled to the mass  $m$  are in the limit  $g_{\text{EM}} \rightarrow 0$ . We have two derivatives acting on  $\psi^\nu$  from  $\square$  and one derivative from the 2-form  $F_{\mu\nu}$  with a local  $U(1)$  gauge-connection  $A_\mu$  that acts on the 4-spinor by the product rule:

$$\begin{aligned} F_{\mu\nu}\psi^\nu &= \partial_\mu A_\nu \psi^\nu - \partial_\nu A_\mu \psi^\nu \\ &= A_\nu \partial_\mu \psi^\nu + \psi^\nu \partial_\mu A_\nu - A_\mu \partial_\nu \psi^\nu - \psi^\nu \partial_\nu A_\mu. \end{aligned} \quad (\text{A11})$$

Promoting the derivative to the gauge-covariant derivative,

$$\partial_\mu \rightarrow \partial_\mu - ieA_\mu, \quad \partial_\nu \rightarrow \partial_\nu - ieA_\nu \quad (\text{A12})$$

the field strength tensor is invariant,

$$F_{\mu\nu} = (\partial_\mu - ieA_\mu)A_\nu - (\partial_\nu - ieA_\nu)A_\mu \quad (\text{A13})$$

$$= \partial_\mu A_\nu - ieA_\mu A_\nu - \partial_\nu A_\mu + ieA_\nu A_\mu \quad (\text{A14})$$

$$= \partial_\mu A_\nu - \partial_\nu A_\mu, \quad (\text{A15})$$

where the last step follows from the fact that  $A_\mu A_\nu = A_\nu A_\mu$ , i.e., the vector potential commutes with itself  $[A_\mu, A_\nu] = 0$ . This is true because the 4-position is commutative:  $[x_\mu, x_\nu] = 0$ . (The covariant derivative  $D_\mu$  in non-commutative geometries will always violate Lorentz invariance with a trace anomaly  $F_\mu^\mu \neq 0$ .)

## 3. The $U(1)$ gauge transformation

Let  $\phi = \phi(x, y, z, t)$  be the local phase angle, i.e., a scalar field. A  $U(1)$  gauge transformation,

$$\psi^\nu \rightarrow e^{i\phi} \psi^\nu \quad (\text{A16})$$

with the vector potential transforming as

$$A_\mu \rightarrow A_\mu - \frac{1}{e}(\partial_\mu \phi), \quad A_\nu \rightarrow A_\nu - \frac{1}{e}(\partial_\nu \phi), \quad (\text{A17})$$

and the covariant derivative

$$\partial_\mu \rightarrow \partial_\mu - ieA_\mu - i(\partial_\mu \phi), \quad \partial_\nu \rightarrow \partial_\nu - ieA_\nu - i(\partial_\nu \phi). \quad (\text{A18})$$

Propagating the product rule

$$F_{\mu\nu}e^{i\phi}\psi^\nu = \partial_\mu A_\nu e^{i\phi}\psi^\nu - \partial_\nu A_\mu e^{i\phi}\psi^\nu \quad (\text{A19})$$

$$= A_\nu \partial_\mu e^{i\phi}\psi^\nu + e^{i\phi}\psi^\nu \partial_\mu A_\nu - A_\mu \partial_\nu e^{i\phi}\psi^\nu - e^{i\phi}\psi^\nu \partial_\nu A_\mu \quad (\text{A20})$$

$$= e^{i\phi} A_\nu \partial_\mu \psi^\nu + A_\nu \psi^\nu \partial_\mu e^{i\phi} + e^{i\phi} \psi^\nu \partial_\mu A_\nu - e^{i\phi} A_\mu \partial_\nu \psi^\nu - A_\mu \psi^\nu \partial_\nu e^{i\phi} - e^{i\phi} \psi^\nu \partial_\nu e^{i\phi} \quad (\text{A21})$$

$$= e^{i\phi} (A_\nu \partial_\mu \psi^\nu + \psi^\nu \partial_\mu A_\nu - A_\mu \partial_\nu \psi^\nu - \psi^\nu \partial_\nu A_\mu) + A_\nu \psi^\nu \partial_\mu e^{i\phi} - A_\mu \psi^\nu \partial_\nu e^{i\phi} \quad (\text{A22})$$

$$= e^{i\phi} F_{\mu\nu} \psi^\nu + A_\nu \psi^\nu \partial_\mu e^{i\phi} - A_\mu \psi^\nu \partial_\nu e^{i\phi}, \quad (\text{A23})$$

we know  $F_{\mu\nu}$  is invariant under the covariant derivative, so we isolate the last two terms and see they cancel under the covariant derivative:

$$A_\nu \psi^\nu \partial_\mu e^{i\phi} = A_\nu \psi^\nu (\partial_\mu - ieA_\mu - i(\partial_\mu \phi)) e^{i\phi} \quad (\text{A24})$$

$$= A_\nu (i(\partial_\mu \phi) e^{i\phi} - ieA_\mu e^{i\phi} - i(\partial_\mu \phi) e^{i\phi}) \psi^\nu \quad (\text{from A3}) \quad (\text{A25})$$

$$= -ieA_\nu A_\mu e^{i\phi} \psi^\nu \quad (\text{A26})$$

$$-A_\mu \psi^\nu \partial_\nu e^{i\phi} = -A_\mu \psi^\nu (\partial_\nu - ieA_\nu - i(\partial_\nu \phi)) e^{i\phi} \quad (\text{A27})$$

$$= -A_\mu (i(\partial_\nu \phi) e^{i\phi} - ieA_\nu e^{i\phi} - i(\partial_\nu \phi) e^{i\phi}) \psi^\nu \quad (\text{from A3}) \quad (\text{A28})$$

$$= ieA_\mu A_\nu e^{i\phi} \psi^\nu. \quad (\text{A29})$$

Thus, the field strength tensor acting on a 4-spinor transforms as

$$F_{\mu\nu} \psi^\nu \rightarrow e^{i\phi} F_{\mu\nu} \psi^\nu \quad (\text{A30})$$

under the  $U(1)$  gauge transformation  $\psi^\nu \rightarrow e^{i\phi} \psi^\nu$ .

## Appendix B: The fundamental representation

In this section, I explicitly write down the bulk equations of motion in the fundamental representation. Unsuppressing the indices of equations (40) and (41), we have

$$\partial^\mu \partial_\mu \chi_L^J + g_{\text{EM}} F_{IJ} \chi_L^J + g_G H \chi_L^J + m^2 \chi_L^J = 0 \quad (\text{B1})$$

$$\partial^\mu \partial_\mu \chi_R^J + g_{\text{EM}} F_{IJ}^\dagger \chi_R^J + g_G H^\dagger \chi_R^J + m^2 \chi_R^J = 0. \quad (\text{B2})$$

### 1. Electromagnetic field

Using the standard Pauli matrix formalism

$$F_{\mu\nu} \rightarrow F_{IJ} = \sigma_{IJ}^l E_l + i\sigma_{IJ}^l B_l = \begin{pmatrix} E_3 & E_1 - iE_2 \\ E_1 + iE_2 & -E_3 \end{pmatrix} + i \begin{pmatrix} B_3 & B_1 - iB_2 \\ B_1 + iB_2 & -B_3 \end{pmatrix}, \quad (\text{B3})$$

the electromagnetic 2-form becomes a first-order differential operator:

$$F_{IJ} = \sigma_{IJ}^l (-\partial_0 A_l - \partial_l A_0) + \frac{i}{2} \sigma_{IJ}^l \epsilon_{lmn} (\partial^m A^n - \partial^n A^m), \quad (\text{B4})$$

where  $\epsilon_{lmn}$  is the 3-dim Levi-Civita tensor and  $\sigma_{IJ}^l$  are the Pauli matrices.

## 2. Gravitational field

The boundary theory provides an explicit recipe for quantizing the classical spin-2 field  $H_{\mu\nu}$  by promoting it to a unique class of Hilbert-Schmidt integral operators:

$$H = \int d\chi_I h(\chi^I, \chi_J) \quad (\text{B5})$$

that acts on a 2-spinor  $\chi^J$  with a complex symmetric kernel,

$$h(\chi^I, \chi_J) = \overline{h(\chi_I, \chi^J)} := h(\chi^J, \chi_I) \quad (\text{B6})$$

such that  $H$  is a traceless trace-class operator:

$$\text{tr } H = \int d\chi_I h(\chi^I, \chi_I) = 0 \quad (\text{B7})$$

where it is assumed the integration constant is zero and

$$\int_{\Omega} \int_{\Omega} d\chi_I |h(\chi^I, \chi_J)|^2 d\chi^J < \infty \quad (I \neq J) \quad (\text{B8})$$

with the domain  $\Omega$  defined by the boundary theory. The simplest (non-unique) choice of a kernel that satisfies all of the above constraints is:

$$h(\chi^I, \chi_J) = \begin{pmatrix} 0 & \chi_0 \overline{\chi_1} \\ \chi_1 \overline{\chi_0} & 0 \end{pmatrix}. \quad (\text{B9})$$

Normalizing  $h(\chi^I, \chi_J)$  we have,

$$\int_{\Omega} \int_{\Omega} d\chi_I |h(\chi^I, \chi_J)|^2 d\chi^J = \int_{\psi_R(a=\lambda_0, \phi=\pi)}^{\psi_L(a=\lambda_{\infty}, \phi=0)} \int_{\psi_R(a=\lambda_0, \phi=\pi)}^{\psi_L(a=\lambda_{\infty}, \phi=0)} d\chi_I |h(\chi^I, \chi_J)|^2 d\chi^J \quad (\text{B10})$$

$$= \int_0^{2\mathcal{R}} \int_0^{2\mathcal{R}} d\chi_I |h(\chi^I, \chi_J)|^2 d\chi^J \quad (\text{B11})$$

$$= \int_0^{2\mathcal{R}} \int_0^{2\mathcal{R}} (d\overline{\chi_1} \ d\overline{\chi_0}) \begin{pmatrix} |\chi_0|^2 |\chi_1|^2 & 0 \\ 0 & |\chi_1|^2 |\chi_0|^2 \end{pmatrix} \begin{pmatrix} d\chi_0 \\ d\chi_1 \end{pmatrix} \quad (\text{B12})$$

$$= \int_0^{2\mathcal{R}} \int_0^{2\mathcal{R}} |\chi_0|^2 |\chi_1|^2 d\overline{\chi_1} d\chi_0 + \int_0^{2\mathcal{R}} \int_0^{2\mathcal{R}} |\chi_1|^2 |\chi_0|^2 d\overline{\chi_0} d\chi_1 \quad (\text{B13})$$

$$= \frac{128\mathcal{R}^6}{9} \quad (\text{B14})$$

where  $\mathcal{R} = \frac{\lambda_{\infty}}{\lambda_0}$ . We absorb the normalization constant inherited from the boundary in the dimensionful constant  $g_G$  that determines the strength of the self-interaction term:

$$H\chi^J = \frac{3g_G}{8\sqrt{2}\mathcal{R}^3} \int d\chi_I h(\chi^I, \chi_J) \chi^J = \frac{3g_G}{8\sqrt{2}\mathcal{R}^3} \int (d\overline{\chi_0} \ d\overline{\chi_1}) \begin{pmatrix} 0 & \chi_0 \overline{\chi_1} \\ \chi_1 \overline{\chi_0} & 0 \end{pmatrix} \begin{pmatrix} \chi_0 \\ \chi_1 \end{pmatrix} \quad (\text{B15})$$

$$= |\chi_0|^2 \frac{3g_G}{8\sqrt{2}\mathcal{R}^3} \int \chi_1 d\overline{\chi_1} + |\chi_1|^2 \frac{3g_G}{8\sqrt{2}\mathcal{R}^3} \int \chi_0 d\overline{\chi_0} \quad (\text{B16})$$

$$= \frac{3g_G}{8\sqrt{2}\mathcal{R}^3} |\chi_0|^2 |\chi_1|^2 \quad (\text{B17})$$

where it is assumed all integration constants are zero. To compute the interaction strength, we must first find a way to calculate  $\lambda_{\infty}$  and normalize the boundary theory. The spin-2 integral operator acting on the spinor field

$$H\chi^J = U(\chi_0, \chi_1) \quad (\text{B18})$$

is the potential of a gravitational harmonic oscillator with a  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetry—a fundamental symmetry of the boundary. Not only is the above manifestly gauge invariant, it is gauge-independent: the spin-2 integral operator transforms as  $H\chi^J \rightarrow H\chi^J$  under the  $U(1)$  gauge transformation  $\chi^J \rightarrow e^{i\phi} \chi^J$ . An interesting observation is this: if we set  $\chi_0 = \chi_1$  we have a  $|\chi|^4$  theory, similar to the potential in a complex  $\varphi^4$  theory like the Higgs potential. We can expand around  $h(\chi^I, \chi_J)$  and add more terms such that we always have even powers of the spinor field.

### 3. Bulk field equations

Putting everything together (and absorbing the Hilbert-Schmidt normalization factor in  $g_G$ ), the gauged bulk field equations are

$$D^\mu D_\mu \chi_L^J + g_E \sigma_{IJ}^l E_l \chi_L^J + i g_B \sigma_{IJ}^l B_l \chi_L^J + g_G |\chi_{L0}|^2 |\chi_{L1}|^2 + m^2 \chi_L^J = 0 \quad (\text{B19})$$

$$D^\mu D_\mu \chi_R^J + g_E \sigma_{IJ}^l E_l \chi_R^J - i g_B \sigma_{IJ}^l B_l \chi_R^J + g_G |\chi_{R0}|^2 |\chi_{R1}|^2 + m^2 \chi_R^J = 0 \quad (\text{B20})$$

(with dimensionful coupling constants  $g_E \sim [V \cdot L]^{-1}$ ,  $g_B \sim [A \cdot L]^{-1}$ , and  $g_G \sim [L]^{\frac{5}{2}}$ ), subject to the constraint of eq. (42), the quantum metric operator:

$$\Psi = \int_{-\infty}^{\infty} g_{IJ} \chi_L^J dt + \int_{-\infty}^{\infty} g_{IJ}^\dagger \bar{\chi}_R^J dt \quad (\text{B21})$$

where local gauge transformations in the bulk correspond to global gauge transformations on the boundary. Solving the above system of equations requires judicious gauge fixing with a deeper understanding of the causal reference frame that is the quantum metric. A strategy for solving the above is to Wick rotate and use Hopf coordinates to look for geon solutions [88]. More exotic spin-2 field configurations  $h(\chi^I, \chi_J)$  are also possible so long as there exists a quantum metric operator that locates the field configuration on the boundary at conformal infinity.

### Appendix C: The De Sitter limit and its dual

The calculation of the dS limit is similar in spirit to Jacobson's [89] and Verlinde's [90] thermodynamic derivation of GR. To rigorously calculate the entropy of the bulk as seen from the ant's-eye view, we need analytic control over the bulk field equations. I leave this to future work. However, if one assumes bulk observers can only access

$$Z_{\text{obs}} \sim e^S \sim e^{2\pi\mathcal{R}} \quad (\text{C1})$$

qubits, where  $\mathcal{R} = \frac{\lambda_\infty}{\lambda_0}$  defines the dimensionless radius of the conformal horizon that hides the cosmological singularity and its dual at future infinity, one can find dS space in the following limit. First, fix the dark energy as a constant by setting

$$\Lambda = \frac{1}{\lambda_0 \lambda_\infty}. \quad (\text{C2})$$

We identified the Bekenstein-Hawking entropy (eq. 25) from the universal wave function and now have two equivalent definitions of entropy, as seen from the ant's-eye view, and can calculate the radius of the cosmic event horizon in the semiclassical limit:

$$\begin{aligned} \frac{\pi R_{\text{CEH}}^2}{\lambda_0^2} &= 2\pi \frac{\lambda_\infty}{\lambda_0} \\ R_{\text{CEH}} &= \sqrt{2\lambda_0 \lambda_\infty} \\ &= \sqrt{\frac{2}{\Lambda}} \\ &\approx 14.26 \text{ billion ly.} \end{aligned} \quad (\text{C3})$$

It is now straightforward to calculate the vacuum energy density (in reduced Planck units):

$$\rho_\Lambda = \rho_\infty = \frac{m_\infty c^2}{\lambda_0^3}. \quad (\text{C4})$$

Now consider a flat acceleration surface with  $2\pi k_B$  entropy per unit (Planck) area  $1/\lambda_0^2$ . Now multiply this quantity by the Hawking temperature  $T(g)/\lambda_0$  per unit length of the accelerating surface. The result is the pressure

$$p_0 = \left( \frac{c^2}{G\lambda_0} \right) g, \quad (\text{C5})$$



where the quantity in parentheses is an *area density* with units  $[\text{mass}][\text{area}]^{-1}$ . The physical construction is an accelerating membrane, like the surface of a glowing soap bubble. The above equation is the ideal gas law  $pV = k_B T$  where  $V = \lambda_0^3$  is the unit volume swept out by the membrane. The above is a thermodynamic derivation of  $F = m_0 g$ , Newton's second law. Now consider the dual acceleration surface with  $4\pi k_B$  entropy per unit  $R_{\text{CEH}}^2$  and repeat the above steps to obtain:

$$p_\infty = \left( \frac{\Lambda \hbar}{c \lambda_0} \right) g, \quad (\text{C6})$$

which is just  $F = m_\infty g$  when multiplied by  $\lambda_0^2$  on both sides. The above pressures obey the ideal gas law, so we can construct the stress-energy tensor  $T_{\mu\nu} = \text{diag}(\rho_\infty, p_\infty, p_\infty, p_\infty)$ . In the UV limit,

$$\lim_{g \rightarrow -c^2/\lambda_0} T_{\mu\nu} = \begin{pmatrix} \rho_\Lambda & 0 & 0 & 0 \\ 0 & p & 0 & 0 \\ 0 & 0 & p & 0 \\ 0 & 0 & 0 & p \end{pmatrix} \quad (\text{C7})$$

one obtains the vacuum stress-energy tensor  $T_{\mu\nu}^{(\text{vac})}$ . The sign on  $g$  distinguishes the expanding and contracting states. Indeed, the first fact we learned from the bootstrap is that energy and curvature are equivalent—the fundamental *assumption* of GR. There are not many options to equate  $T_{\mu\nu}^{(\text{vac})}$  to curvature on the left-hand side. The second fact we learned is that the *form* of the laws of physics are independent of the chosen coordinate system—no coordinate system was used to solve eq. (2). The third fact we learned is that the fundamental charge of the vacuum, the gravitational charge  $m_\infty$ , is equivalent to inertial mass. These deductions are none other than the equivalence principle and general covariance! Given these deductions, the simplest equations that follow are none other than Einstein's field equations:

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = \kappa T_{\mu\nu} \quad (\text{C8})$$

where  $R_{\mu\nu}$  is the Ricci curvature tensor,  $R$  is the scalar curvature,  $g_{\mu\nu}$  is the metric tensor, and  $\kappa = 1$  in natural units. Quantum mechanics—when assumed universal—naturally contains GR.

There is then the dual vacuum stress-energy tensor for the contracting state. As before, we can construct a stress-energy tensor  $T_{\mu\nu} = \text{diag}(\rho_0, p_0, p_0, p_0)$  for an ideal gas. In the IR limit,

$$\lim_{g \rightarrow c^2/\lambda_\infty} T_{\mu\nu} = \begin{pmatrix} \rho_0 & 0 & 0 & 0 \\ 0 & -p & 0 & 0 \\ 0 & 0 & -p & 0 \\ 0 & 0 & 0 & -p \end{pmatrix} \quad (\text{C9})$$

one obtains a bizarre substance with the Planck energy density and a positive vacuum energy pressure  $p = \Lambda$  (with the minus sign absorbed) that satisfies the equation of state for dark matter:  $w = p/\rho_0 \approx 0$ . Are

$$G_{\mu\nu} = \kappa T_{\mu\nu}^{(\text{dual-vac})} \quad (\text{C10})$$

the classical field equations for dark matter? Does this explain how the cosmological constant  $\Lambda$  enters local galaxy dynamics through a universal acceleration scale  $a_0 \approx cH_0 \approx c^2 \Lambda^{1/2}$ , and how something as small as a galaxy "knows" about the dS radius associated with the whole of the observable universe? Is the dark matter graviball solutions to the bulk field equations?

- 
- [1] P. A. R. Ade et al. (Planck), Planck 2015 results. XIII. Cosmological parameters, *Astron. Astrophys.* **594** (2016) A13
  - [2] P.J.E. Peebles, Anomalies in physical cosmology, *Annals Phys.* **447** (2022) 159-169.
  - [3] Y. Aharonov and D. Bohm, Significance of electromagnetic potentials in quantum theory, *Phys. Rev.*, **115**(3)

- (1959) 485–491.
- [4] Y. Aharonov and D. Bohm, Further Considerations on Electromagnetic Potentials in the Quantum Theory, *Phys. Rev.*, **123**(4) (1961) 1511–1524.
- [5] A. Tonomura et. al., Evidence for Aharonov-Bohm effect with magnetic field completely shielded from electron wave, *Phys. Rev. Lett.*, **56**(8) (1986) 792–795.

- [6] N. Osakabe et. al., Experimental confirmation of Aharonov–Bohm effect using a toroidal magnetic field confined by a superconductor, *Phys. Rev. A*, **34**(2) (1986) 815–822.
- [7] M. F. Pusey, J. Barrett, and T. Rudolph, On the reality of the quantum state, *Nat. Phys.*, **8**(6) (2012) 475–478.
- [8] M. F. Pusey, J. Barrett, and T. Rudolph, *The quantum state cannot be interpreted statistically*, arXiv:1111.3328v1 (2011).
- [9] M. Leifer, Is the quantum state real? An extended review of  $\psi$ -ontology theorems, *Quanta*, **3**(1) (2014) 67–155.
- [10] Rovelli C. *Quantum Gravity*. Cambridge University Press (2004).
- [11] H. Everett, *The theory of the universal wave function*, In *The many-worlds interpretation of quantum mechanics* Princeton University Press (2015) 1-140.
- [12] M. Jammer. *The Philosophy of Quantum Mechanics: The Interpretations of Quantum Mechanics in Historical Perspective*, Wiley (1974)
- [13] J. Feldbrugge, J.L. Lehnens, and N. Turok, No rescue for the no boundary proposal: Pointers to the future of quantum cosmology, *Phys. Rev. D* **97** (2018) 023509.
- [14] M. Milgrom, *MOND—a pedagogical review*, arXiv:astro-ph/0112069.
- [15] F. Lelli et. al., One law to rule them all: the radial acceleration relation of galaxies, *Astrophys. J.* **836** (2017) 152.
- [16] Kyu-Hyun Chae et. al., On the Presence of a Universal Acceleration Scale in Elliptical Galaxies, *Astrophys. J. Lett.* **903** (2020) L31.
- [17] L. Susskind, String theory and the principles of black hole complementarity, *Phys. Rev. Lett.* **71** (1993) 2367-2368.
- [18] L. Susskind, Strings, black holes and Lorentz contraction, *Phys. Rev. D* **49** (1994) 6606-6611.
- [19] S. Hossenfelder, Minimal Length Scale Scenarios for Quantum Gravity, *Living Rev. Rel.* **16** (2013) 2.
- [20] R. Israel et. al., Stringy Horizons and UV/IR Mixing, *JHEP* **11** (2015) 164.
- [21] R.J., Adler, P. Chen, and D.I Santiago, The Generalized Uncertainty Principle and Black Hole Remnants, *Gen. Relativ. Gravit.* **33** (2001) 2100-2108.
- [22] R.J., Adler and D.I., Santiago, On gravity and the uncertainty principle, *Mod. Phys. Lett.* **14** (1999) 1371.
- [23] J. Feldbrugge, J.L. Lehnens, and N. Turok, Inconsistencies of the New No-Boundary Proposal, *Universe* **4** (2018) 100.
- [24] A. M. Polyakov, De Sitter space and eternity, *Nucl. Phys. B* **797** (2008) 199–217.
- [25] A. M. Polyakov, Infrared instability of the de Sitter space, arXiv:1209.4135
- [26] G. Obied, H. Ooguri, L. Spodyneiko, and C. Vafa, *De Sitter Space and the Swampland*, arXiv:1806.08362.
- [27] N. Agmon et. al., *Lectures on the string landscape and the Swampland*, arXiv:2212.06187
- [28] L. Boyle and N. Turok, Two-Sheeted Universe, Analyticity and the Arrow of Time, arXiv:2109.06204
- [29] L. Boyle, F. Kieran, and N. Turok, CPT-Symmetric Universe, *Phys. Rev. Lett.* **121** (2018) 251301.
- [30] L. Boyle, F. Kieran, and N. Turok, The Big Bang, CPT, and neutrino dark matter, *Annals Phys.* **438** (2022) 168767.
- [31] N. Afshordi, Dark Energy, Black Hole Entropy, and the First Precision Measurement in Quantum Gravity, 1003.4811
- [32] C. Prescod-Weinstein, N. Afshordi, et al., Stellar Black Holes and the Origin of Cosmic Acceleration, *Phys. Rev. D* **80** (2009) 043513
- [33] J.M. Maldacena, The Large N limit of superconformal field theories and supergravity, *Adv. Theor. Math. Phys.* **2** (1998) 231–252.
- [34] L. Susskind, The World as a hologram, *J. Math. Phys.*, **36** (1995) 6377–6396.
- [35] G. 't Hooft, The Holographic principle: Opening lecture, *Subnucl. Ser.* **37** (2001) 72–100.
- [36] A. Laddha, et. al., The Holographic Nature of Null Infinity, *SciPost Phys.* **10** (2021) 041.
- [37] A. Laddha, et. al., Squinting at massive fields from infinity, arXiv:2207.06406
- [38] S. Raju, Failure of the split property in gravity and the information paradox, *Class. Quant. Grav.* **39** (2022) 064002.
- [39] V. Hubeny, The AdS/CFT Correspondence, *Class. Quant. Grav.* **32** (2015) 124010.
- [40] M. Ekaterina et. al., Quantum time: Experimental multi-time correlations, *Phys. Rev. D*, **96** (5) (2017) 102005.
- [41] S.M. Carroll, *Reality as a vector in Hilbert space*, Springer (2022).
- [42] S.M. Carroll, *Mad-dog everettianism: Quantum mechanics at its most minimal*, Springer (2019).
- [43] Carroll, S. M., & Singh, A. (2021). Quantum mereology: Factorizing Hilbert space into subsystems with quasiclassical dynamics. *Phys. Rev. A*, **103** (2), 022213.
- [44] J.B. Hartle and S.W. Hawking, Wave Function of the Universe, *Phys. Rev. D* **28** (1983) 2960–2975.
- [45] J.B. Hartle, S.W. Hawking, and T. Hertog, The Classical Universes of the No-Boundary Quantum State, *Phys. Rev. D* **77** (2008) 123537.
- [46] A. Vilenkin, Creation of Universes from Nothing, *Phys. Lett. B* **117** (1982) 25–28.
- [47] A. Vilenkin, The Birth of Inflationary Universes, *Phys. Rev. D* **27** (1983) 2848.
- [48] Torre, C.G. Midisuperspace Models of Canonical Quantum Gravity, *Int. J. Theor. Phys.* **38** (1999) 1081–1102.
- [49] J. Feldbrugge, J.L. Lehnens, and N. Turok, Lorentzian quantum cosmology, *Phys. Rev. D* **95** (2017) 103508
- [50] B.S. DeWitt, Quantum Theory of Gravity. I. The Canonical Theory, *Phys. Rev.* **160** (1967) 1113.
- [51] H.K. Urbantke, The Hopf fibration—seven times in physics, *J. Geom. Phys.* **46** (2003) 125-150.
- [52] A. Jadczyk, On Conformal Infinity and Compactifications of the Minkowski Space, *Adv. Appl. Clifford Algebras*, **21** (2011) 721–756. arXiv:1008.4703
- [53] F. Wilczek, Physics in 100 years, *Physics Today*. **69** (2016) 32–39
- [54] A.M. Reclariu, Lectures on Celestial Holography, arXiv:2107.02075
- [55] S. Pasterski, M. Pate, and A.M. Raclariu, Celestial Holography, arXiv:2111.11392
- [56] A. Paramekanti, L. Balents and M. P. Fisher, Ring exchange, the exciton bose liquid, and bosonization in two dimensions, *Physical Review B* **66** (2002) 054526.
- [57] R. M. Nandkishore and M. Hermele, Fractons, *Annu. Rev. Condens. Matter Phys.* **10** (2019) 295–313.
- [58] N. Seiberg and S.-H. Shao, Exotic Symmetries, Duality, and Fractons in 2+1-Dimensional Quantum Field Theory, arXiv:2003.10466.
- [59] N. Seiberg and S.-H. Shao, Exotic U(1) Symmetries, Duality, and Fractons in 3+1-Dimensional Quantum Field

- Theory, *SciPost Phys.* **9** (2020) 046.
- [60] Y. You, J. Bibo, T. L. Hughes and F. Pollmann, Fractonic critical point proximate to a higher-order topological insulator: How does uv blend with ir?, arXiv:2101.01724.
  - [61] G. W. Gibbons and S. W. Hawking, Action Integrals and Partition Functions in Quantum Gravity, *Phys. Rev. D* **15** (1977) 2752-2756.
  - [62] C. White, The double copy: gravity from gluons, *Contemp. Phys.* **59** (2018) 109.
  - [63] G. Rubino et al., Experimental verification of an indefinite causal order, *Sci. Adv.* **3** (2017) e1602589.
  - [64] G. Chiribella, et al., Quantum computations without definite causal structure, *Phys. Rev. A* (2017).
  - [65] Li, Y. Indefinite causality. *Nat. Phys.* **13** (2017) 419.
  - [66] P.A. Guérin and Č. Brukner, Observer-dependent locality of quantum events, *New J. Phys.* (2018) **20** 103031.
  - [67] M. Zych et al., Bell's theorem for temporal order. *Nat Commun* (2019) **10**, 3772.
  - [68] E. Castro-Ruiz, F. Giacomini, and Č. Brukner, Dynamics of Quantum Causal Structures, *Physical Review X*, (2018) **8** 1.
  - [69] H. Weyl, Gravitation and electricity, *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)* **1918** (1918) 465.
  - [70] R. H. Dicke, Mach's principle and invariance under transformation of units, *Phys. Rev.*, **125** (1962) 2163.
  - [71] Shau-Yu Lan et al., A Clock Directly Linking Time to a Particle's Mass. *Science*, **339** (2013) 554-557.
  - [72] E. Abdall, et al., Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies, *JHEAp*, **34** (2022) 49–211.
  - [73] A. de Oliveira-Costa, M. Tegmark, M. Zaldarriaga, and A. Hamilton, The Significance of the largest scale CMB fluctuations in WMAP, *Phys. Rev. D* **69** (2004) 063516.
  - [74] R. Howl and I. Fuentes, Quantum frequency interferometry: With applications ranging from gravitational wave detection to dark matter searches, *AVS Quantum Sci.* **5(1)** (2023) 014402.
  - [75] S. Hossenfelder, *Lost in Math: How Beauty Leads Physics Astray*, *Basic Books* (2018).
  - [76] E. Anderson, The Problem of Time in Quantum Gravity, arXiv:1009.2157
  - [77] A. O. Barvinsky, Unitarity approach to quantum cosmology, *Phys. Rept.* **bf 230** (1993) 237.
  - [78] A. Y. Kamenshchik, Time in quantum theory, the Wheeler-DeWitt equation and the Born-Oppenheimer approximation, arXiv:1809.08083v1
  - [79] J. Feldbrugge, J.L. Lehnert, and N. Turok, No smooth beginning for spacetime, *Phys. Rev. Lett.* **119** (2017) 171301.
  - [80] H.O. Ukavwe, The Role of Deductive Reasoning in the Deductive-Nomological Model of Scientific Explanation, *Nasara Journal of Philosophy* (2018) 13-19.
  - [81] Misner, Thorne, and Wheeler (MTW), *Gravitation*, chapters 7, 20, and 25.
  - [82] H. Roos, Independence of local algebras in quantum field theory, *Comm. Math. Phys.* **16** (1970) 238–246.
  - [83] R. Haag, *Local quantum physics: Fields, particles, algebras*, Springer (1992).
  - [84] M. Kamionkowski and A. Riess, The Hubble tension and early dark energy, arXiv:2211.04492.
  - [85] N. Turok and L. Boyle, Gravitational entropy and the flatness, homogeneity and isotropy puzzles, arXiv:2201.07279.
  - [86] N. Turok and L. Boyle, Thermodynamic solution of the homogeneity, isotropy and flatness puzzles (and a clue to the cosmological constant), arXiv:2210.01142.
  - [87] J. M. Bardeen, B. Carter and S. W. Hawking, The Four laws of black hole mechanics, *Commun. Math. Phys.* **31** (1973) 161-170.
  - [88] J. A. Wheeler, Geons. *Phys. Rev.*, **97** (1955) 511–536.
  - [89] T. Jacobson, Thermodynamics of Spacetime: The Einstein Equation of State, *Phys. Rev. Lett.* **75** (1995) 1260-1263.
  - [90] E. P. Verlinde, On the Origin of Gravity and the Laws of Newton, *JHEP* **4** (2011) 29.
  - [91] The Bekenstein-Hawking area entropy law is a geometric consequence of the fact that  $\psi \in L^2(S_a^2 \times S_b^1)$ . The action is computable in Hawking and Gibbons's original paper because the Euclidean section is non-singular: the entropy is evaluated on a region of a spacetime manifold bounded by some surface  $r = r_0 > 2M$  with compact topology  $S^2 \times S^1$  (i.e., a 2-sphere cross periodic time) [61]. The Ricci scalar vanishes in the Schwarzschild metric, so the action is determined only by the Gibbons-Hawking-York boundary term and is thus an integral over the boundary  $S^2 \times S^1$ . The world density  $\rho = |\psi|^2$  is nothing but an integral over the boundary  $S_a^2 \times S_b^1$  and precisely why one can write  $\rho = \rho(S)$  with  $S = \frac{A}{4\lambda^2}$ .