

S1 Fig — all supporting information figures




Figure A: For each model, the 1850–2014 trend in its ensemble members of the AMOC streamfunction strength at 26.5°N and at 35°N. Even though the aerosol forcing is the same within each model, there is still a large scatter of ensemble member slopes.




Figure B: Fraction of all timeseries for a given model that have a significant increase in λ (so both in streamfunction strength and SST Index). Only the models that do have such significant increases are shown, and the full list of these can be found in Table D in S1 Table. Note that the colours for the models are different from other figures.




Figure C: Same as Fig 3 in the Main Text, but with the slices coloured according to the p-value of the corresponding λ . Darker blue is a smaller p-value and so more statistically significant.




Figure D: Fraction of timeseries for a given model that have a significant increase in any of the three CSD indicators. Only the models that do have such significant increases are shown, and the full list of these can be found in Tables D, E, F in S1 Table. Note that the colours for the models are different from other figures.




Figure E: Same as Fig 6 in the Main Text but for the correlations between the p-values of different CSD indicators.




Figure F: Same as Fig 6 in the Main Text for the AR1 instead of λ .




Figure G: Same as Fig 6 in the Main Text for the variance instead of λ .




Figure H: Same as Fig 3 in the Main Text, but with a window size of 50 years.




Figure I: Same as Fig 3 in the Main Text, but with a window size of 70 years.




Figure J: Same as Fig 4 in the Main Text, but with a window size of 50 years.




Figure K: Same as Fig 4 in the Main Text, but with a window size of 70 years.




Figure L: Same as Fig 6 in the Main Text for the ensemble mean values of each of the models.




Figure M: The distribution of the ensemble member p-values for the λ trend of the AMOC streamfunction strengths at 26.5°N and 35°N and the AMOC SST Index, plotted against the ocean component of the model. The models with lower p-values have a range of different ocean components.




Figure N: Mean p-value for all λ timeseries of a given model. Note that the colours and shading of the models are different from other figures. Note that the models with highest indication of CSD are those with the *lowest* p-values.




Figure O: Mean p-value for all CSD indicator timeseries of a given model. Note that the colours and shading of the models are different from other figures.