

1.4 Matrix Group

June 12, 2026

Matrix Group

Definition 0.1. In $\mathbb{C}^n$, $\bar{e}_i = (0, \dots, 1^{\text{th}}, 0, \dots)$ is a vector space basis element.

Definition 0.2. An *endomorphism* α of $\mathbb{C}^n$ is defined as

$$\alpha \bar{e}_i = \sum_{j=1}^n a_{ji} \bar{e}_j$$

where $\alpha = (a_{ij}) \in M_n(\mathbb{C})$. (We can think of $\alpha \bar{e}_i$ as the i^{th} column vector.)

We use α for a matrix as well as for the endomorphism.

We define multiplication of two endomorphisms α, β with corresponding matrices $(a_{ij}), (b_{ij})$ and construct (c_{ij}) where

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}.$$

Lemma 0.1. $M_n(\mathbb{C}) \cong \mathbb{C}^{n^2}$ as vector spaces.

Proof. Construct the function

$$\begin{aligned} f: M_n(\mathbb{C}) &\rightarrow \mathbb{C}^{n^2}, \\ (a_{ij}) &\mapsto (b_{ij}), \end{aligned}$$

where $b_{i+(j-1)n} = a_{ij}$.

Observe as j goes from 1 to n , $(j-1)n$ goes from 0 to $(n-1)n$ and as i goes from 1 to n as well, we get $i + (j-1)n$ go from 1 to n^2 .

Observe f is surjective by construction and if $a_{ij} \neq a_{i'j'}$ then $b_{i+(j-1)n} \neq b_{i'+(j'-1)n}$ hence f is bijective.

Observe f is linear hence $M_n(\mathbb{C}) \cong \mathbb{C}^{n^2}$. □

Remark 0.1. Observe we can equip $M_n(\mathbb{C})$ with a topology by ensuring the given map above to be a homeomorphism. By constructing

$$\mathcal{T}_{M_n(\mathbb{C})} = \{f^{-1}(U) \mid U \in \mathcal{T}_{\mathbb{C}^{n^2}}\}.$$

Lemma 0.2. Let T be any topological space,

$$\phi: T \rightarrow M_n(\mathbb{C}).$$

ϕ is continuous if and only if $\pi_{ij} \circ \phi$ is continuous.

Proof. Only if condition is trivial.

If condition:

$$\pi_{ij} \circ \phi: T \rightarrow \mathbb{C} \text{ is continuous.}$$

Now $\prod_{1 \leq i, j \leq n} \pi_{ij} \circ \phi: T \rightarrow \mathbb{C}^{n^2}$ is continuous by product topology (or box topology as $\mathbb{C}^{n^2}$ is finite Cartesian product). $\square$

Corollary 0.1. The product of two matrices is continuous.

Proof. Consider the map

$$q: M_n(\mathbb{C}) \times M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$$

defined by matrix multiplication, i.e.,

$$q(\sigma, \tau) = \sigma\tau.$$

We can factor q as follows:

$$\begin{array}{ccc} M_n(\mathbb{C}) \times M_n(\mathbb{C}) & \xrightarrow{q} & M_n(\mathbb{C}) \\ \downarrow f \times f & & \downarrow f \\ \mathbb{C}^{n^2} \times \mathbb{C}^{n^2} & \xrightarrow{\rho} & \mathbb{C}^{n^2} \end{array}$$

where $f: M_n(\mathbb{C}) \rightarrow \mathbb{C}^{n^2}$ is the isomorphism identifying matrices with their entries, and ρ is a polynomial map defined by

$$\rho((b_{ij}), (b'_{ij})) = (c_{ij}),$$

with

$$c_{ij} = \sum_{k=1}^n b_{ik} b'_{kj}.$$

Since ρ is a polynomial map, it is continuous. The map f is a homeomorphism, and thus $f \times f$ is also continuous. Therefore, the composition $q = f \circ \rho \circ (f \times f)$ is continuous.

For any open set U in $M_n(\mathbb{C})$, we can express $U = f^{-1}(O)$ for some open set O in $\mathbb{C}^{n^2}$. Since q is continuous, the preimage $q^{-1}(U)$ is open in $M_n(\mathbb{C}) \times M_n(\mathbb{C})$. This proves the continuity of the matrix product. $\square$

—

$$q^{-1}(U) = q^{-1}(f^{-1}(O)) = (f \circ q)^{-1}(O)$$

which is open as $f \circ q$ is continuous.

$\square$

****Notation:**** tA is the transpose of a matrix and $\overline{A}$ is the complex conjugate of a matrix.

Lemma 0.3. Transpose and complex conjugate are homeomorphisms.

Now in $(\mathbb{C}^{n^2}, \|\cdot\|_1)$, let $v = (v_k)_{k=1}^{n^2}$ where $|v_k| \leq 1$ for $1 \leq k \leq n^2$. We define

$$\|v\|_1 = \max_{1 \leq k \leq n^2} |v_k|.$$

Observe $f : M_n(\mathbb{C}) \rightarrow \mathbb{C}^{n^2}$ as a homeomorphism. $f(UC_n)$ is closed and bounded, and compact. This implies UC_n is compact.

□

Remark 0.10. The choice of the norm does not matter as all norms in finite dimension vector space are equivalent (add the proof to that in the appendix).

—