

Measurable function.

* If $(X, \mathcal{M})$ and $(Y, \mathcal{N})$ are measurable spaces a mapping $f: X \rightarrow Y$ is called $(\mathcal{M}, \mathcal{N})$ -measurable or measurable if $f^{-1}(E) \in \mathcal{M}$ for all $E \in \mathcal{N}$.

Remark

Composition of measurable mappings are measurable.

Proposition

If $\mathcal{N}$ is generated by $\mathcal{E}$, then $f: X \rightarrow Y$ is $(\mathcal{M}, \mathcal{N})$ -measurable iff $f^{-1}(E) \in \mathcal{M}$ for all $E \in \mathcal{E}$.

Proof: Only if condition is trivial. For if condition, observe $\{E \subseteq Y \mid f^{-1}(E) \in \mathcal{M}\}$ is a σ -algebra, that contains $\mathcal{E}$ by hypothesis; it therefore contains $\mathcal{N}$. □

Corollary

If X and Y are metric (or topological spaces),

every continuous mapping is (B_X, B_Y) measurable.

* If $(X, \mathcal{M})$ is a measurable space, a real or complex value f^n on X is called measurable if f is $(\mathcal{M}, B_{\mathbb{R} \text{ or } \mathbb{C}})$ measurable.