

Graphical Models

David A. Levin

DAIICT

January 2020

Graphical Models

- ▶ A **graphical model** specifies a factorization which a joint density must obey.
- ▶ Example: If X_1, \dots, X_n are independent, then the joint density completely factors into marginals:

$$f(x_1, \dots, x_n) = \prod_{i=1}^n f_{X_i}(x_i). \quad (1)$$

- ▶ This is too strong a constraint to require of general applicable probability models.
- ▶ A graphical model specifies a class of joint densities which obey a factorization which still affords useful simplification but preserves some generality.

Graphical Models

- ▶ A **graphical model** specifies a factorization which a joint density must obey.
- ▶ Example: If X_1, \dots, X_n are independent, then the joint density completely factors into marginals:

$$f(x_1, \dots, x_n) = \prod_{i=1}^n f_{X_i}(x_i). \quad (1)$$

- ▶ This is too strong a constraint to require of general applicable probability models.
- ▶ A graphical model specifies a class of joint densities which obey a factorization which still affords useful simplification but preserves some generality.

Graphical Models

- ▶ A **graphical model** specifies a factorization which a joint density must obey.
- ▶ Example: If X_1, \dots, X_n are independent, then the joint density completely factors into marginals:

$$f(x_1, \dots, x_n) = \prod_{i=1}^n f_{X_i}(x_i). \quad (1)$$

- ▶ This is too strong a constraint to require of general applicable probability models.
- ▶ A graphical model specifies a class of joint densities which obey a factorization which still affords useful simplification but preserves some generality.


Graphical Models

- ▶ A **graphical model** specifies a factorization which a joint density must obey.
- ▶ Example: If X_1, \dots, X_n are independent, then the joint density completely factors into marginals:

$$f(x_1, \dots, x_n) = \prod_{i=1}^n f_{X_i}(x_i). \quad (1)$$

- ▶ This is too strong a constraint to require of general applicable probability models.
- ▶ A graphical model specifies a class of joint densities which obey a factorization which still affords useful simplification but preserves some generality.

- ▶ nodes represent variables
- ▶ directed arrows represent dependencies
- ▶ Any variable with an arrow pointing to x_i is called a **parent** of x_i ; we denote by pa_i all the parents of x_i .




- ▶ A joint distribution respects the graphical model encoded by a directed graph if

$$p(x_1, \dots, x_n) = \prod_{i=1}^n p(x_i \mid \text{pa}_i).$$

- ▶ In the example graph, a joint distribution obeys the graphical model if

$$p(x_1, x_2, x_3, x_4) = p(x_1)p(x_2 \mid x_1, x_4)p(x_3 \mid x_2, x_4)p(x_4).$$

- ▶ nodes represent variables
- ▶ directed arrows represent dependencies
- ▶ Any variable with an arrow pointing to x_i is called a **parent** of x_i ; we denote by pa_i all the parents of x_i .




- ▶ A joint distribution respects the graphical model encoded by a directed graph if

$$p(x_1, \dots, x_n) = \prod_{i=1}^n p(x_i \mid \text{pa}_i).$$

- ▶ In the example graph, a joint distribution obeys the graphical model if

$$p(x_1, x_2, x_3, x_4) = p(x_1)p(x_2 \mid x_1, x_4)p(x_3 \mid x_2, x_4)p(x_4).$$

- ▶ nodes represent variables
- ▶ directed arrows represent dependencies
- ▶ Any variable with an arrow pointing to x_i is called a **parent** of x_i ; we denote by pa_i all the parents of x_i .




- ▶ A joint distribution respects the graphical model encoded by a directed graph if

$$p(x_1, \dots, x_n) = \prod_{i=1}^n p(x_i \mid \text{pa}_i).$$

- ▶ In the example graph, a joint distribution obeys the graphical model if

$$p(x_1, x_2, x_3, x_4) = p(x_1)p(x_2 \mid x_1, x_4)p(x_3 \mid x_2, x_4)p(x_4).$$

- ▶ nodes represent variables
- ▶ directed arrows represent dependencies
- ▶ Any variable with an arrow pointing to x_i is called a **parent** of x_i ; we denote by pa_i all the parents of x_i .




- ▶ A joint distribution respects the graphical model encoded by a directed graph if

$$p(x_1, \dots, x_n) = \prod_{i=1}^n p(x_i \mid \text{pa}_i).$$

- ▶ In the example graph, a joint distribution obeys the graphical model if

$$p(x_1, x_2, x_3, x_4) = p(x_1)p(x_2 \mid x_1, x_4)p(x_3 \mid x_2, x_4)p(x_4).$$

- ▶ nodes represent variables
- ▶ directed arrows represent dependencies
- ▶ Any variable with an arrow pointing to x_i is called a **parent** of x_i ; we denote by pa_i all the parents of x_i .



- ▶ A joint distribution respects the graphical model encoded by a directed graph if

$$p(x_1, \dots, x_n) = \prod_{i=1}^n p(x_i \mid \text{pa}_i).$$

- ▶ In the example graph, a joint distribution obeys the graphical model if

$$p(x_1, x_2, x_3, x_4) = p(x_1)p(x_2 \mid x_1, x_4)p(x_3 \mid x_2, x_4)p(x_4).$$

- ▶ The graphical model encodes *conditional independence* statements.
- ▶ A set of variables X and Y are conditionally independent given Z if

$$\mathbb{P}(X \in A, Y \in B \mid Z) = \mathbb{P}(X \in A \mid Z) \mathbb{P}(Y \in B \mid Z).$$

- ▶ In terms of pdfs/pmfs,

$$p(x, y \mid z) = p(x \mid z) p(y \mid z).$$

- ▶ Or equivalently,

$$p(x \mid y, z) = p(x \mid z).$$

- ▶ The graphical model encodes *conditional independence* statements.
- ▶ A set of variables X and Y are conditionally independent given Z if

$$\mathbb{P}(X \in A, Y \in B \mid Z) = \mathbb{P}(X \in A \mid Z) \mathbb{P}(Y \in B \mid Z).$$

- ▶ In terms of pdfs/pmfs,

$$p(x, y \mid z) = p(x \mid z) p(y \mid z).$$

- ▶ Or equivalently,

$$p(x \mid y, z) = p(x \mid z).$$

- ▶ The graphical model encodes *conditional independence* statements.
- ▶ A set of variables X and Y are conditionally independent given Z if

$$\mathbb{P}(X \in A, Y \in B \mid Z) = \mathbb{P}(X \in A \mid Z) \mathbb{P}(Y \in B \mid Z).$$

- ▶ In terms of pdfs/pmfs,

$$p(x, y \mid z) = p(x \mid z) p(y \mid z).$$

- ▶ Or equivalently,

$$p(x \mid y, z) = p(x \mid z).$$

- ▶ The graphical model encodes *conditional independence* statements.
- ▶ A set of variables X and Y are conditionally independent given Z if

$$\mathbb{P}(X \in A, Y \in B \mid Z) = \mathbb{P}(X \in A \mid Z) \mathbb{P}(Y \in B \mid Z).$$


- ▶ In terms of pdfs/pmfs,

$$p(x, y \mid z) = p(x \mid z) p(y \mid z).$$

- ▶ Or equivalently,

$$p(x \mid y, z) = p(x \mid z).$$

Example 1




- ▶ Note that

$$p(x, y | z) = \frac{p(x, y, z)}{\sum_{x, y} p(x, y, z)} = \frac{p(x | z) p(y | z) p(z)}{\sum_x p(x | z) \sum_y p(y | z) p(z)} = p(x | z) p(y | z)$$

- ▶ Any time X and Y are separated by a **tail-to-tail** vertex Z , they are conditionally independent.
- ▶ Note that X and Y are not independent unconditionally, in general.
- ▶ Exercise: Show by example that X and Y are not necessarily independent.

Example 1




- ▶ Note that

$$p(x, y | z) = \frac{p(x, y, z)}{\sum_{x, y} p(x, y, z)} = \frac{p(x | z) p(y | z) p(z)}{\sum_x p(x | z) \sum_y p(y | z) p(z)} = p(x | z) p(y | z)$$

- ▶ Any time X and Y are separated by a **tail-to-tail** vertex Z , they are conditionally independent.
- ▶ Note that X and Y are not independent unconditionally, in general.
- ▶ Exercise: Show by example that X and Y are not necessarily independent.

Example 1




- ▶ Note that

$$p(x, y | z) = \frac{p(x, y, z)}{\sum_{x, y} p(x, y, z)} = \frac{p(x | z) p(y | z) p(z)}{\sum_x p(x | z) \sum_y p(y | z) p(z)} = p(x | z) p(y | z)$$

- ▶ Any time X and Y are separated by a **tail-to-tail** vertex Z , they are conditionally independent.
- ▶ Note that X and Y are not independent unconditionally, in general.
- ▶ Exercise: Show by example that X and Y are not necessarily independent.

Example 1




- ▶ Note that

$$p(x, y | z) = \frac{p(x, y, z)}{\sum_{x, y} p(x, y, z)} = \frac{p(x | z) p(y | z) p(z)}{\sum_x p(x | z) \sum_y p(y | z) p(z)} = p(x | z) p(y | z)$$

- ▶ Any time X and Y are separated by a **tail-to-tail** vertex Z , they are conditionally independent.
- ▶ Note that X and Y are not independent unconditionally, in general.
- ▶ Exercise: Show by example that X and Y are not necessarily independent.

Example 1




- ▶ Note that

$$p(x, y | z) = \frac{p(x, y, z)}{\sum_{x, y} p(x, y, z)} = \frac{p(x | z) p(y | z) p(z)}{\sum_x p(x | z) \sum_y p(y | z) p(z)} = p(x | z) p(y | z)$$

- ▶ Any time X and Y are separated by a **tail-to-tail** vertex Z , they are conditionally independent.
- ▶ Note that X and Y are not independent unconditionally, in general.
- ▶ Exercise: Show by example that X and Y are not necessarily independent.


Example 2



$$\begin{aligned} p(x, y | z) &= \frac{p(x, y, z)}{p(z)} = \frac{p(x) p(z | x) p(y | z)}{p(z)} \\ &= \frac{p(x | z) p(z) p(y | z)}{p(z)} = p(x | z) p(y | z) \end{aligned}$$

- ▶ Thus, if x and y are connected by a **head-to-tail** vertex z , then they are conditionally independent.


Example 2



$$\begin{aligned} p(x, y | z) &= \frac{p(x, y, z)}{p(z)} = \frac{p(x)p(z | x)p(y | z)}{p(z)} \\ &= \frac{p(x | z)p(z)p(y | z)}{p(z)} = p(x | z)p(y | z) \end{aligned}$$

- ▶ Thus, if x and y are connected by a **head-to-tail** vertex z , then they are conditionally independent.

Example 2



$$\begin{aligned} p(x, y | z) &= \frac{p(x, y, z)}{p(z)} = \frac{p(x)p(z | x)p(y | z)}{p(z)} \\ &= \frac{p(x | z)p(z)p(y | z)}{p(z)} = p(x | z)p(y | z) \end{aligned}$$


- ▶ Thus, if x and y are connected by a **head-to-tail** vertex z , then they are conditionally independent.



$$\begin{aligned}
 p(x, y) &= \sum_z p(x, y, z) = \sum_z p(x) p(y) p(z | x, y) \\
 &= p(x) p(y) \sum_z p(z | x, y) = p(x) p(y).
 \end{aligned}$$

so X and Y are independent given \emptyset .

- ▶ Exercise: Give an example to show that X and Y are not conditionally independent given Z .
- ▶ Even more is true:




X and Y are not independent given W (in general).



$$\begin{aligned}
 p(x, y) &= \sum_z p(x, y, z) = \sum_z p(x) p(y) p(z \mid x, y) \\
 &= p(x) p(y) \sum_z p(z \mid x, y) = p(x) p(y).
 \end{aligned}$$

so X and Y are independent given \emptyset .

- ▶ Exercise: Give an example to show that X and Y are not conditionally independent given Z .
- ▶ Even more is true:



X and Y are not independent given W (in general).




$$\begin{aligned}
 p(x, y) &= \sum_z p(x, y, z) = \sum_z p(x) p(y) p(z \mid x, y) \\
 &= p(x) p(y) \sum_z p(z \mid x, y) = p(x) p(y).
 \end{aligned}$$

so X and Y are independent given \emptyset .

- ▶ Exercise: Give an example to show that X and Y are not conditionally independent given Z .

- ▶ Even more is true:




X and Y are not independent given W (in general).



$$\begin{aligned}
 p(x, y) &= \sum_z p(x, y, z) = \sum_z p(x) p(y) p(z \mid x, y) \\
 &= p(x) p(y) \sum_z p(z \mid x, y) = p(x) p(y).
 \end{aligned}$$

so X and Y are independent given \emptyset .

- ▶ Exercise: Give an example to show that X and Y are not conditionally independent given Z .
- ▶ Even more is true:



X and Y are not independent given W (in general).

- ▶ Let A, B, C be sets of vertices. The set C **blocks** A to B if every path from a vertex $a \in A$ to a vertex $b \in B$ contains either
 - ▶ a vertex $c \in C$ so that c is head-to-tail or tail-to-tail, or
 - ▶ a vertex v which is head-to-head, and which neither belongs to C or has a decendent belonging to C .



Theorem 1 (*d*-separation).

If C blocks A and B in the di-graph, then the variables A and the variables B are conditionally independent given C .

- ▶ Let A, B, C be sets of vertices. The set C **blocks** A to B if every path from a vertex $a \in A$ to a vertex $b \in B$ contains either
 - ▶ a vertex $c \in C$ so that c is head-to-tail or tail-to-tail, or
 - ▶ a vertex v which is head-to-head, and which neither belongs to C or has a decendent belonging to C .



Theorem 1 (*d*-separation).

If C blocks A and B in the di-graph, then the variables A and the variables B are conditionally independent given C .

- ▶ Let A, B, C be sets of vertices. The set C **blocks** A to B if every path from a vertex $a \in A$ to a vertex $b \in B$ contains either
 - ▶ a vertex $c \in C$ so that c is head-to-tail or tail-to-tail, or
 - ▶ a vertex v which is head-to-head, and which neither belongs to C or has a decendent belonging to C .



Theorem 1 (*d*-separation).

If C blocks A and B in the di-graph, then the variables A and the variables B are conditionally independent given C .

- ▶ Let A, B, C be sets of vertices. The set C **blocks** A to B if every path from a vertex $a \in A$ to a vertex $b \in B$ contains either
 - ▶ a vertex $c \in C$ so that c is head-to-tail or tail-to-tail, or
 - ▶ a vertex v which is head-to-head, and which neither belongs to C or has a decendent belonging to C .
- ▶


Theorem 1 (*d*-separation).

If C blocks A and B in the di-graph, then the variables A and the variables B are conditionally independent given C .

- ▶ Let A, B, C be sets of vertices. The set C **blocks** A to B if every path from a vertex $a \in A$ to a vertex $b \in B$ contains either
 - ▶ a vertex $c \in C$ so that c is head-to-tail or tail-to-tail, or
 - ▶ a vertex v which is head-to-head, and which neither belongs to C or has a decendent belonging to C .
- ▶

Theorem 1 (d -separation).

If C blocks A and B in the di-graph, then the variables A and the variables B are conditionally independent given C .




- ▶ The above is a **Hidden Markov Model**.
- ▶ We can use d -separation to establish useful conditional independence relations:



$$(X_1, \dots, X_k) \perp (X_{k+1}, \dots, X_n) \mid Z_k$$



$$X_1, \dots, X_{k-1} \perp X_k \mid Z_k$$




- ▶ The above is a **Hidden Markov Model**.
- ▶ We can use d -separation to establish useful conditional independence relations:



$$(X_1, \dots, X_k) \perp (X_{k+1}, \dots, X_n) \mid Z_k$$



$$X_1, \dots, X_{k-1} \perp X_k \mid Z_k$$



- ▶ The above is a **Hidden Markov Model**.
- ▶ We can use d -separation to establish useful conditional independence relations:



$$(X_1, \dots, X_k) \perp (X_{k+1}, \dots, X_n) \mid Z_k$$



$$X_1, \dots, X_{k-1} \perp X_k \mid Z_k$$

