

MATH324 (Statistics) – Lecture Notes
McGill University

Masoud Asgharian

Winter 2019

Contents

1	Lecture 0	2
2	Lecture 1	2
3	Lecture 2	2
3.1	Markov's Inequality	2
3.2	Chebyshev's Inequality	3
4	Lecture 3	4
5	Lecture 4	4
6	Lecture 5	4
7	Lecture 6	4
8	Lecture 7	4
9	Lecture 8	4
10	Lecture 9	4
11	Lecture 10	4
12	Lecture 11	4
13	Lecture 12	4

1 Lecture 0

2 Lecture 1

3 Lecture 2

3.1 Markov's Inequality

Let X be a random variable and h be a **non-negative** function; ie:

$$h : R \rightarrow R^+ \cup \{0\} = [0, \infty)$$

Suppose $E(h(x)) < \infty$, then for some $\lambda > 0$, we have:

$$P(h(x) \geq \lambda) \leq \frac{E[h(x)]}{\lambda} \quad (1)$$

Proof. Suppose X is a continuous random variable:

$$\begin{aligned} E[h(x)] &= \int_x h(x) d_x(x) dx \\ &= \left(\int_{x:h(x) \geq \lambda} h(x) f_x(x) dx + \int_{x:h(x) < \lambda} h(x) f_x(x) dx \right) \\ &\geq \int_{x:h(x) \geq \lambda} h(x) f_x(x) dx \quad \text{since } h \geq 0 \\ &\geq \lambda \int_{x:h(x) \geq \lambda} f_x(x) dx = \lambda P(h(x) \geq \lambda) \\ \implies P(h(x) \geq \lambda) &\leq \frac{E(h(x))}{\lambda} \end{aligned}$$

The proof for the discrete case is similar. □

Now consider $h(x) = (x - \mu)^2$, then:

$$\begin{aligned} P(|x - \mu| \geq \lambda) &= P((x - \mu)^2 \geq \lambda^2) \\ &\leq \frac{E[(x - \mu)^2]}{\lambda^2} \quad \text{if } E[(x - \mu)^2] < \infty \end{aligned}$$

Let $\mu = E(X)$, then $E[(x - \mu)^2] = \text{Var}(X)$ denoted by σ_x^2 . We therefore have:

$$P(|x - \mu_x| \geq \lambda) \leq \frac{\sigma_x^2}{\lambda^2} \quad \text{where } \mu_x = E(x) \quad (2)$$

Now consider $\lambda = K\sigma_x$ where K is a known number. Then:

$$P(|x - \mu_x| \geq K\sigma_x) \geq \frac{\sigma_x^2}{K^2\sigma_x^2} = \frac{1}{K^2} \quad (3)$$

3.2 Chebyshev's Inequality

Chebyshev's Inequality is a special case of Markov's Inequality.

- 4 Lecture 3
- 5 Lecture 4
- 6 Lecture 5
- 7 Lecture 6
- 8 Lecture 7
- 9 Lecture 8
- 10 Lecture 9
- 11 Lecture 10
- 12 Lecture 11
- 13 Lecture 12