

Data Consistency Conditions in Image Reconstruction from Projections

Rolf Clackdoyle


Hubert Curien Laboratory, CNRS UMR 5516, St. Etienne

Supported by ANR grant “DROITE” (ANR-12-BS01-0018)

2D parallel projections:

2D parallel projections:

$p(\phi, \cdot)$ - parallel projection at angle ϕ



2D parallel projections:

$$p(\phi, r) = \int f(r\alpha + s\beta) ds$$


$$\phi \in [-\pi/2, \pi/2]$$

$$\alpha = (\cos \phi, \sin \phi)$$

$$\beta = (-\sin \phi, \cos \phi)$$

$p(\phi, \cdot)$ - parallel projection at angle ϕ

For short.... $p = \mathbf{R}f$



2D parallel projections:

$$p(\phi, r) = \int f(r\alpha + s\beta) ds$$


$$\phi \in [-\pi/2, \pi/2]$$

$$\alpha = (\cos \phi, \sin \phi)$$

$$\beta = (-\sin \phi, \cos \phi)$$

$p(\phi, \cdot)$ - parallel projection at angle ϕ

For short.... $p = \mathbf{R}f$



We are interested in **truncated** parallel projections.

Data Consistency Conditions (DCC)

Helgason-Ludwig DCC

$$\text{Let } M_0(\phi) = \int p(\phi, r) \, dr.$$

Helgason-Ludwig DCC

$$\text{Let } M_0(\phi) = \int p(\phi, r) \, dr.$$

If $p = \mathbf{R}f$ then $M_0(\phi) = \text{constant}$

Helgason-Ludwig DCC

$$\text{Let } M_0(\phi) = \int p(\phi, r) \, dr.$$

If $p = \mathbf{R}f$ then $M_0(\phi) = \text{constant}$

$$\text{Let } M_n(\phi) = \int p(\phi, r) \, r^n \, dr.$$

Helgason-Ludwig DCC

$$\text{Let } M_0(\phi) = \int p(\phi, r) \, dr.$$


If $p = \mathbf{R}f$ then $M_0(\phi) = \text{constant}$

$$\text{Let } M_n(\phi) = \int p(\phi, r) \, r^n \, dr.$$

If $p = \mathbf{R}f$ then $M_n(\phi) = a_0 U^n + a_1 U^{n-1} V + \dots a_n V^n$
where $U = \cos \phi$ and $V = \sin \phi$


New DCC

Take some line outside the target object,
w.l.o.g. the line $y = y_0$ with $y_0 > 0$.



New DCC


Take some line outside the target object,
w.l.o.g. the line $y = y_0$ with $y_0 > 0$.



Let $B_0(x) = \int_0^\pi p(\phi, r^*) \frac{1}{\cos \phi} d\phi$, where $r^* = x \cos \phi + y_0 \sin \phi$.

New DCC


Take some line outside the target object,
w.l.o.g. the line $y = y_0$ with $y_0 > 0$.



Let $B_0(x) = \int_0^\pi p(\phi, r^*) \frac{1}{\cos \phi} d\phi$, where $r^* = x \cos \phi + y_0 \sin \phi$.

If $p = \mathbf{R}f$ then $B_0(x) = \text{constant}$

Take some line outside the target object,
w.l.o.g. the line $y = y_0$ with $y_0 > 0$.




Let $B_0(x) = \int_0^\pi p(\phi, r^*) \frac{1}{\cos \phi} d\phi$, where $r^* = x \cos \phi + y_0 \sin \phi$.

If $p = \mathbf{R}f$ then $B_0(x) = \text{constant}$

Let $B_n(x) = \int_0^\pi p(\phi, r^*) \frac{\tan^n \phi}{\cos \phi} d\phi$, where $r^* = x \cos \phi + y_0 \sin \phi$.

New DCC

Take some line outside the target object,
w.l.o.g. the line $y = y_0$ with $y_0 > 0$.



Let $B_0(x) = \int_0^\pi p(\phi, r^*) \frac{1}{\cos \phi} d\phi$, where $r^* = x \cos \phi + y_0 \sin \phi$.

If $p = \mathbf{R}f$ then $B_0(x) = \text{constant}$


Let $B_n(x) = \int_0^\pi p(\phi, r^*) \frac{\tan^n \phi}{\cos \phi} d\phi$, where $r^* = x \cos \phi + y_0 \sin \phi$.

If $p = \mathbf{R}f$ then $B_n(x) = b_0 + b_1 x + b_2 x^2 + \dots + b_n x^n$

Comparing Helgason-Ludwig DCC and the new DCC


$$M_n(\phi) = \int p(\phi, r) r^n dr$$

Comparing Helgason-Ludwig DCC and the new DCC

$$M_n(\phi) = \int p(\phi, r) r^n dr$$



no truncation allowed

Comparing Helgason-Ludwig DCC and the new DCC

$$M_n(\phi) = \int p(\phi, r) r^n dr$$


no truncation allowed
(limited-angle OK)


Comparing Helgason-Ludwig DCC and the new DCC

$$M_n(\phi) = \int p(\phi, r) r^n dr \quad B_n(x) = \int_0^\pi p(\phi, r^*) \frac{\tan^n \phi}{\cos \phi} d\phi$$


$$\text{where } r^* = x \cos \phi + y_0 \sin \phi$$

no truncation allowed
(limited-angle OK)

Comparing Helgason-Ludwig DCC and the new DCC


$$M_n(\phi) = \int p(\phi, r) r^n dr \quad B_n(x) = \int_0^\pi p(\phi, r^*) \frac{\tan^n \phi}{\cos \phi} d\phi$$


$$\text{where } r^* = x \cos \phi + y_0 \sin \phi$$

no truncation allowed
(limited-angle OK)

no limited-angle allowed

Comparing Helgason-Ludwig DCC and the new DCC


$$M_n(\phi) = \int p(\phi, r) r^n dr \quad B_n(x) = \int_0^\pi p(\phi, r^*) \frac{\tan^n \phi}{\cos \phi} d\phi$$


$$\text{where } r^* = x \cos \phi + y_0 \sin \phi$$



no truncation allowed
(limited-angle OK)

no limited-angle allowed
(truncation OK)



Simulations





Simulations





Simulations






Simulations





Simulations






Simulations





Simulations






Simulations





Simulations






Simulations






Simulations






Simulations



Simulations





Simulations




$$T = 18, \quad (t_0, t_1) = (2, 17), \quad A = 7$$


Dynamic phantom and parallel geometry






Dynamic phantom and parallel geometry




Simulations





Simulations








Simulations



Simulations



Search for the (non-linear) parameters (t_0, t_1) and A such that the DCC are “best satisfied” (are polynomials of the correct degree), when the moving component is subtracted from the data.

(The alternative, not using DCC, is to solve for (t_0, t_1) , A , and $f(x, y)$ - which combines a small non-linear problem with a huge linear problem.)




Search for the (non-linear) parameters (t_0, t_1) and A such that the DCC are “best satisfied” (are polynomials of the correct degree), when the moving component is subtracted from the data.

(The alternative, not using DCC, is to solve for (t_0, t_1) , A , and $f(x, y)$ - which combines a small non-linear problem with a huge linear problem.)


The non-linear estimation found $(\tilde{t}_0, \tilde{t}_1) = (1.98, 16.99)$ and $\tilde{A} = 7.03$

(The true values were $(t_0^*, t_1^*) = (2, 17)$ and $A^* = 7$)




Uncorrected






**Inconsistent component
subtracted**






**Inconsistent component
subtracted**



Simulations



Simulations



Data Consistency Conditions - Summary

Data Consistency Conditions - Summary

- Data Consistency Conditions (DCC) express redundancies between projections

Data Consistency Conditions - Summary

- Data Consistency Conditions (DCC) express redundancies between projections
- Checking the DCC is a way to see if the data follows the forward model

Data Consistency Conditions - Summary

- Data Consistency Conditions (DCC) express redundancies between projections
- Checking the DCC is a way to see if the data follows the forward model
- The advantage of the DCC is that they can be checked without having to perform reconstruction (i.e. much quicker)

Data Consistency Conditions - Summary

- Data Consistency Conditions (DCC) express redundancies between projections
- Checking the DCC is a way to see if the data follows the forward model
- The advantage of the DCC is that they can be checked without having to perform reconstruction (i.e. much quicker)
- New DCC for truncated projections have been described (2014)

Data Consistency Conditions - Summary

- Data Consistency Conditions (DCC) express redundancies between projections
- Checking the DCC is a way to see if the data follows the forward model
- The advantage of the DCC is that they can be checked without having to perform reconstruction (i.e. much quicker)
- New DCC for truncated projections have been described (2014)
- DCC were successfully used to estimate non-linear motion parameters

Thanks for your attention!

