

Direct Comparisons **Total Heart Failure Adverse Drug Reactions**
P–score(Comparison: other vs Placebo) **OR** **95%–CI**

Osimertinib + antiangiogenesis	0	0.14		13.96	[1.59; 122.88]
Icotinib	0	0.17		17.55	[0.47; 652.53]
Antiangiogenesis	0	0.19		9.99	[0.78; 128.33]
Gefitinib + PARP inhibitor	0	0.29		7.53	[0.17; 342.91]
Erlotinib + MMP inhibitor	0	0.33		5.57	[0.17; 182.38]
Erlotinib + IGF–1R inhibitor	0	0.33		4.03	[0.53; 30.41]
Osimertinib + chemotherapy	0	0.34		3.65	[0.81; 16.57]
Erlotinib + HER3 inhibitor	0	0.35		4.63	[0.16; 130.03]
Erlotinib + antiangiogenesis	0	0.42		2.64	[0.54; 12.93]
Gefitinib	1	0.46		2.48	[0.32; 19.53]
Chemotherapy	0	0.50		2.08	[0.32; 13.31]
Gefitinib + chemotherapy	0	0.53		1.81	[0.08; 39.02]
Dacomitinib	0	0.54		1.81	[0.13; 25.28]
Erlotinib + MET inhibitor	0	0.54		1.79	[0.30; 10.54]
Erlotinib	3	0.54		1.81	[0.47; 6.97]
Osimertinib	2	0.58		1.56	[0.42; 5.80]
Lazertinib	0	0.59		1.44	[0.20; 10.60]
Afatinib	1	0.65		1.23	[0.32; 4.79]
Erlotinib + chemotherapy	0	0.65		1.08	[0.09; 12.64]
Almonertinib	0	0.67		0.83	[0.02; 37.45]
Lazertinib + Amivantamab	0	0.73		0.80	[0.11; 5.70]
Erlotinib + HGF inhibitor	0	0.86		0.20	[0.00; 8.60]
Gefitinib/Erlotinib	0	0.88		0.31	[0.03; 2.77]

0.01 0.1 1 10 100
 Favours Control Favours Experimental