

Study or Subgroup	Events	Total	Weight (common)	Weight (random)	Proportion [95% CI]	Serious Vascular Adverse Drug Reactions	
drug = Afatinib							
Mountzios,2021	1	59	0.2%	0.8%	0.02 [0.00; 0.09]		
Takata,2023	1	103	0.7%	1.3%	0.01 [0.00; 0.05]		
Tu,2022	8	541	2.3%	1.7%	0.01 [0.01; 0.03]		
NCT02131259,2016	10	1602	16.2%	1.9%	0.01 [0.00; 0.01]		
NCT02047903,2018	7	152	0.2%	0.7%	0.05 [0.02; 0.09]		
2009-014563-39,2016	35	1154	2.5%	1.7%	0.03 [0.02; 0.04]		
NCT02044380,2016	1	14	0.0%	0.1%	0.07 [0.00; 0.34]		
Goldberg,2020	3	85	0.2%	0.6%	0.04 [0.01; 0.10]		
Goss,2021	17	392	0.6%	1.2%	0.04 [0.03; 0.07]		
Park,2016	9	160	0.2%	0.7%	0.06 [0.03; 0.10]		
Kato,2015	7	229	0.5%	1.1%	0.03 [0.01; 0.06]		
Miller,2012	15	390	0.7%	1.3%	0.04 [0.02; 0.06]		
Total (common effect, 95% CI)		4881	24.2%	.	0.01 [0.01; 0.02]		
Total (random effect, 95% CI)				13.2%	0.03 [0.02; 0.04]		
Heterogeneity: Tau ² = 0.0002; Chi ² = 53.18, df = 11 (P < 0.0001); I ² = 79.3%							
drug = Osimertinib							
Lorenzi,2022	12	126	0.1%	0.4%	0.10 [0.05; 0.16]		
Park,2020(LM)	1	40	0.1%	0.4%	0.03 [0.00; 0.13]		
NCT02474355,2019	80	3014	7.3%	1.9%	0.03 [0.02; 0.03]		
NCT02094261,2015	10	210	0.3%	0.9%	0.05 [0.02; 0.09]		
Malik,2022	1	60	0.2%	0.8%	0.02 [0.00; 0.09]		
Kiura,2018(80mg, Extension)	3	201	0.9%	1.4%	0.01 [0.00; 0.04]		
Kiura,2018(80mg, First Line)	2	60	0.1%	0.5%	0.03 [0.00; 0.12]		
Provencio,2021	2	155	0.8%	1.3%	0.01 [0.00; 0.05]		
Lv,2023a	1	40	0.1%	0.4%	0.03 [0.00; 0.13]		
NCT02157883,2015	1	38	0.1%	0.4%	0.03 [0.00; 0.14]		
Tsubata,2022	1	33	0.1%	0.3%	0.03 [0.00; 0.16]		
2014-000531-17,2018	14	210	0.2%	0.7%	0.07 [0.04; 0.11]		
NCT03434418,2022	1	17	0.0%	0.1%	0.06 [0.00; 0.29]		
Kenmotsu,2022	3	60	0.1%	0.4%	0.05 [0.01; 0.14]		
Soo,2022	3	77	0.1%	0.5%	0.04 [0.01; 0.11]		
John,2023	5	337	1.4%	1.6%	0.01 [0.00; 0.03]		
Ohe,2019	10	279	0.5%	1.2%	0.04 [0.02; 0.06]		
Akamatsu,2018	10	279	0.5%	1.2%	0.04 [0.02; 0.06]		
García Campelo,2024	18	428	0.7%	1.3%	0.04 [0.03; 0.07]		
NCT04035486,2023	4	275	1.2%	1.5%	0.01 [0.00; 0.04]		
Lu,2024	3	143	0.4%	1.1%	0.02 [0.00; 0.06]		
Total (common effect, 95% CI)		6082	15.2%	.	0.03 [0.02; 0.03]		
Total (random effect, 95% CI)				18.3%	0.03 [0.02; 0.03]		
Heterogeneity: Tau ² = < 0.0001; Chi ² = 30.26, df = 20 (P = 0.0657); I ² = 33.9%							
drug = Gefitinib							
Douillard,2014	3	107	0.2%	0.8%	0.03 [0.01; 0.08]		
Schuetz,2018	1	222	3.1%	1.8%	0.00 [0.00; 0.02]		
Cappuzzo,2016	1	58	0.2%	0.7%	0.02 [0.00; 0.09]		
Noronha,2020	42	170	0.1%	0.3%	0.25 [0.18; 0.32]		
Hosomi,2020	3	171	0.6%	1.3%	0.02 [0.00; 0.05]		
Zhao,2021	4	154	0.4%	1.0%	0.03 [0.01; 0.07]		
Park,2016	6	159	0.3%	0.9%	0.04 [0.01; 0.08]		
Yang,2016	1	118	0.9%	1.4%	0.01 [0.00; 0.05]		
Goss,2013	5	249	0.8%	1.4%	0.02 [0.01; 0.05]		
Wu,2017	2	607	11.6%	1.9%	0.00 [0.00; 0.01]		
Zhang,2012	2	147	0.7%	1.3%	0.01 [0.00; 0.05]		
NCT04248829,2022	7	197	0.4%	1.0%	0.04 [0.01; 0.07]		
Total (common effect, 95% CI)		2359	19.2%	.	0.01 [0.00; 0.01]		
Total (random effect, 95% CI)				13.7%	0.02 [0.01; 0.04]		
Heterogeneity: Tau ² = 0.0003; Chi ² = 72.09, df = 11 (P < 0.0001); I ² = 84.7%							
drug = Erlotinib							
Markóczy,2018	1	62	0.2%	0.8%	0.02 [0.00; 0.09]		
NCT01183858,2013(300mg)	2	158	0.8%	1.4%	0.01 [0.00; 0.04]		
Park,2015	4	207	0.7%	1.3%	0.02 [0.01; 0.05]		
Brueckl,2018	2	385	4.7%	1.8%	0.01 [0.00; 0.02]		
Kato,2018	12	77	0.0%	0.2%	0.16 [0.08; 0.26]		
Spigel,2017	10	244	0.4%	1.0%	0.04 [0.02; 0.07]		
Scagliotti,2015a	24	517	0.7%	1.3%	0.05 [0.03; 0.07]		
Piccirillo,2022	5	79	0.1%	0.4%	0.06 [0.02; 0.14]		
Nakagawa,2022	3	225	1.1%	1.5%	0.01 [0.00; 0.04]		
Spigel,2018	1	64	0.3%	0.8%	0.02 [0.00; 0.08]		
Deplanque,2016	4	74	0.1%	0.4%	0.05 [0.01; 0.13]		
Lilenbaum,2008	1	52	0.2%	0.6%	0.02 [0.00; 0.10]		
Saito,2019	2	114	0.4%	1.1%	0.02 [0.00; 0.06]		
Scagliotti,2012	17	477	0.9%	1.4%	0.04 [0.02; 0.06]		
Wu,2015	1	110	0.8%	1.3%	0.01 [0.00; 0.05]		
Natale,2011	20	614	1.2%	1.5%	0.03 [0.02; 0.05]		
Cicenas,2016	4	322	1.6%	1.6%	0.01 [0.00; 0.03]		
Scagliotti,2020	1	70	0.3%	0.9%	0.01 [0.00; 0.08]		
Groen,2013	4	64	0.1%	0.3%	0.06 [0.02; 0.15]		
Herbst,2011	19	313	0.3%	1.0%	0.06 [0.04; 0.09]		
Stinchcombe,2019	6	45	0.0%	0.1%	0.13 [0.05; 0.27]		
Goss,2021	14	395	0.7%	1.3%	0.04 [0.02; 0.06]		
Sequist,2011	5	83	0.1%	0.4%	0.06 [0.02; 0.14]		
Zhou,2021	6	153	0.3%	0.8%	0.04 [0.01; 0.08]		
Scagliotti,2015b	2	290	2.7%	1.8%	0.01 [0.00; 0.02]		
Ramalingam,2012	4	94	0.1%	0.6%	0.04 [0.01; 0.11]		
Kelly,2015	7	611	3.4%	1.8%	0.01 [0.00; 0.02]		
Ciuleanu,2012	6	196	0.4%	1.1%	0.03 [0.01; 0.07]		
NCT02186301,2017	1	50	0.2%	0.6%	0.02 [0.00; 0.11]		
Cappuzzo,2010	3	433	3.9%	1.8%	0.01 [0.00; 0.02]		
Ramalingam,2014	7	436	1.7%	1.6%	0.02 [0.01; 0.03]		
Lara,2016	1	32	0.1%	0.3%	0.03 [0.00; 0.16]		
NCT01211483,2013	3	71	0.1%	0.5%	0.04 [0.01; 0.12]		
Hirsch,2011	6	69	0.1%	0.3%	0.09 [0.03; 0.18]		
Gridelli,2012	5	372	1.8%	1.6%	0.01 [0.00; 0.03]		
Khozin,2014	3	84	0.2%	0.6%	0.04 [0.01; 0.10]		
NCT01887886,2015	1	5	0.0%	0.0%	0.20 [0.01; 0.72]		
NCT00977470,2015	1	38	0.1%	0.4%	0.03 [0.00; 0.14]		
NCT00606502,2010	1	101	0.6%	1.3%	0.01 [0.00; 0.05]		
Total (common effect, 95% CI)		7786	31.3%	.	0.02 [0.01; 0.02]		
Total (random effect, 95% CI)				37.6%	0.02 [0.02; 0.03]		
Heterogeneity: Tau ² = 0.0001; Chi ² = 101.53, df = 38 (P < 0.0001); I ² = 62.6%							
drug = Befotertinib							
Lu,2022(50mg)	12	176	0.2%	0.6%	0.07 [0.04; 0.12]		
Lu,2023	15	182	0.2%	0.6%	0.08 [0.05; 0.13]		
Total (common effect, 95% CI)		358	0.3%	.	0.07 [0.05; 0.10]		
Total (random effect, 95% CI)				1.2%	0.07 [0.05; 0.10]		
Heterogeneity: Tau ² = 0; Chi ² = 0.26, df = 1 (P = 0.6094); I ² = 0%							
drug = Nazartinib							
Tan,2020(150mg, Phase I Part)	1	73	0.3%	1.0%	0.01 [0.00; 0.07]		
Tan,2020(150mg, Phase II Part)	1	45	0.1%	0.5%	0.02 [0.00; 0.12]		
Tan,2020(225mg, Phase I Part)	1	28	0.1%	0.2%	0.04 [0.00; 0.18]		
Tan,2020(350mg, Phase I Part)	1	11	0.0%	0.0%	0.09 [0.00; 0.41]		
Total (common effect, 95% CI)		157	0.5%	.	0.02 [0.00; 0.04]		
Total (random effect, 95% CI)				1.8%	0.02 [0.00; 0.04]		
Heterogeneity: Tau ² = 0; Chi ² = 1.09, df = 3 (P = 0.7798); I ² = 0%							
drug = Furmonertinib							
Shi,2021	3	220	1.0%	1.5%	0.01 [0.00; 0.04]		
Shi,2020(80mg)	1	48	0.1%	0.6%	0.02 [0.00; 0.11]		
Total (common effect, 95% CI)		268	1.2%	.	0.01 [0.00; 0.03]		
Total (random effect, 95% CI)				2.0%	0.01 [0.00; 0.03]		
Heterogeneity: Tau ² = 0; Chi ² = 0.11, df = 1 (P = 0.7441); I ² = 0%							
drug = Almonertinib							
Yang,2020(110mg, Escalation Cohort 2)	2	6	0.0%	0.0%	0.33 [0.04; 0.78]		
Yang,2020(110mg, Phase II Extension)	16	244	0.2%	0.8%	0.07 [0.04; 0.10]		
Lu,2022	10	214	0.3%	0.9%	0.05 [0.02; 0.08]		
Total (common effect, 95% CI)		464	0.6%	.	0.06 [0.04; 0.08]		
Total (random effect, 95% CI)				1.7%	0.06 [0.03; 0.09]		
Heterogeneity: Tau ² = 0.0002; Chi ² = 2.85, df = 2 (P = 0.2400); I ² = 29.9%							
drug = Lazertinib							
Cho,2022(120mg, phase I)	3	25	0.0%	0.1%	0.12 [0.03; 0.31]		
Cho,2022(160mg, phase I)	1	23	0.0%	0.2%	0.04 [0.00; 0.22]		
Cho,2022(240mg, phase I)	1	24	0.0%	0.2%	0.04 [0.00; 0.21]		
Cho,2022(240mg, phase II)	3	78	0.1%	0.5%	0.04 [0.01; 0.11]		
NCT04075396,2021(240mg)	3	15	0.0%	0.0%	0.20 [0.04; 0.48]		
NCT04075396,2021(320mg)	1	13	0.0%	0.1%	0.08 [0.00; 0.36]		
García Campelo,2024	13	213	0.2%	0.8%	0.06 [0.03; 0.10]		
NCT04248829,2022	8	196	0.3%	0.9%	0.04 [0.02; 0.08]		
Total (common effect, 95% CI)		587	0.8%	.	0.05 [0.03; 0.07]		
Total (random effect, 95% CI)				2.7%	0.05 [0.03; 0.07]		
Heterogeneity: Tau ² = 0; Chi ² = 4.62, df = 7 (P = 0.7059); I ² = 0%							
drug = Mobocertinib							
Hida,2021(120mg, Phase 1 Part)	1	4	0.0%	0.0%	0.25 [0.01; 0.81]		
Hida,2021(160mg, Phase II Part)	1	33	0.1%	0.3%	0.03 [0.00; 0.16]		
Total (common effect, 95% CI)		37	0.1%	.	0.03 [0.00; 0.09]		
Total (random effect, 95% CI)				0.3%	0.04 [0.00; 0.10]		
Heterogeneity: Tau ² = 0.0003; Chi ² = 1.01, df = 1 (P = 0.3148); I ² = 1%							
drug = Dacomitinib							
Lacouture,2015(45mg)	1	25</					