

Partial Differentiation

Mathematical Methods in the Physical Sciences


Steve Mazza

Naval Postgraduate School
Monterey, CA



SE3030, Winter/2014
Quantitative Methods of Systems Engineering

Lorenz Attractor



Definition

Derivatives of a multi-variable function where all variables are held fixed during differentiation except the variable of interest.

To denote this we generically write $\frac{\partial}{\partial r}$, which means the partial derivative with respect to r . We more frequently see $\frac{\partial z}{\partial r}$, which means the partial derivative of z with respect to r . In equations of more than two variables we may see $\left(\frac{\partial z}{\partial r}\right)_x$, which denotes the partial derivative of z with respect to r , holding x constant.

4.1.12

$$z = x^2 + 2y^2, x = r\cos\theta, y = r\sin\theta$$

$$z = x^2 + 2y^2$$

$$z = r^2\cos^2\theta + 2r^2\sin^2\theta$$

$$\left(\frac{\partial z}{\partial y}\right)_\theta = r^2\cos^2\theta + 2r^2\sin^2\theta$$

Power Series in Two Variables

Our standard power series expansions can be re-written as partial differential equations.

Definition

$$f(x, y) = \sum_{n=0}^{\infty} \frac{1}{n!} \left(h \frac{\partial}{\partial x} + k \frac{\partial}{\partial y} \right)^n f(a, b)$$

Total Differentiation in 2 Variables

$$dz = \frac{\partial}{\partial x} dx + \frac{\partial}{\partial y} dy$$

Approximation: For sufficiently small values of Δx and Δy ,

- $\Delta z = \Delta f = f_x(x, y)\Delta x + f_y(x, y)\Delta y$, and
- $f(x + \Delta x, y + \Delta y) \equiv f(x, y) + f_x(x, y)\Delta x + f_y(x, y)\Delta y$.

Approximations Using Differentials

Example 4, Boas, p. 197

The relative error in length measurement is $\pm 5\%$ and the relative error in radius measurement is $\pm 10\%$. We want to find the largest value that $|dR/R|$ can have.

$$R = \frac{kl}{r^2}$$

$$\ln R = \ln k + \ln l - 2 \ln r$$

$$\begin{aligned}\frac{dR}{R} &= \left| \frac{dl}{l} \right| - 2 \left| \frac{dr}{r} \right| \\ &= 0.05 + 2(0.10) \\ &= 0.25\end{aligned}$$

Chain Rule

The chain rule is a formula for computing the derivative of the composition of two or more functions.

In General

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx}$$

Find dy/dx if $y = \ln \sin 2x$

$$\begin{aligned}\frac{dy}{dx} &= \frac{1}{\sin 2x} \cdot \frac{d}{dx}(\sin 2x) \\ &= \frac{1}{\sin 2x} \cdot \cos 2x \cdot \frac{d}{dx}(2x) \\ &= 2 \cot 2x\end{aligned}$$

Implicit Differentiation

We can differentiate with respect to a given term in some cases where we cannot solve for that term with respect to another.

Given $x + e^x = t$, find dx/dt

We realize that x is a function of t even though we cannot solve x for t directly.

$$\begin{aligned}x + e^x &= t \\ \frac{dx}{dt} + e^x \frac{dx}{dt} &= 1 \\ \frac{dx}{dt} &= \frac{1}{1 + e^x}\end{aligned}$$

This example can be found in Boas, p 202.


Chain Rule (Redux)

We can extend our earlier examples of the Chain Rule where $z = f(x, y)$ where x and y were functions of t by considering the case where x and y are two variables, s and t .

Applications

Helps us locate

- Hills
- Valleys
- Saddle Points



FIN