



STELLAR ASTROPHYSICS

Eddington Luminosity

Oriol Abril

Teacher
Jordi Isern

December 2017

Assume a cloud around a star of mass M and luminosity L .

1. Find a mass-luminosity condition for ejecting the cloud

As seen in class, the radiation force per unit mass can be written as $f_r = \frac{\kappa}{c} F_r$, where κ is the opacity, c the speed of light and F_r is the flux of the star. In order to eject the outer layers, this force should be greater than the gravitational force felt per unit mass by a particle of the envelope.

In the case of spherical symmetry, if the mass of said envelope is neglected, and only the inner mass is taken into account as the whole star, then $f_G = G \frac{M}{R^2}$ (independently of the inner mass distribution) where G is the universal gravitation constant, M the mass of the star and R the radius of the star.

Therefore, substituting the flux by luminosity over surface, the condition is:

$$\begin{aligned} f_r > f_G &\Rightarrow \frac{\kappa}{c} \frac{L}{4\pi R^2} > G \frac{M}{R^2} \\ \frac{M}{L} &< \frac{\kappa}{4\pi G c} \end{aligned} \quad (1)$$

2. Find the terminal velocity of the cloud if it starts at rest at a distance R

The energy per unit mass of the envelope will be the energy due to the radiation pressure and the energy due to the gravitational potential of the inner star:

$$u = u_r + u_G = \frac{\kappa}{c} \frac{L}{4\pi R} - G \frac{M}{R} \quad (2)$$

Note that $f = -\nabla u$.

The terminal velocity will be achieved when all the potential energy is converted to kinetic energy:

$$\begin{aligned} t &= \frac{1}{2} v_t^2 \\ t = u &\rightarrow \frac{1}{2} v_t^2 = \frac{\kappa}{c} \frac{L}{4\pi R} - G \frac{M}{R} \\ v_t^2 &= \frac{2GM}{R} \left(\frac{\kappa L}{4\pi c G M} - 1 \right) \end{aligned} \quad (3)$$

where all energies continue being expressed as energy per unit mass of the envelope and v_t^2 is the terminal velocity. Note that as the terminal velocity is only defined when the condition in Eq. 1 is fulfilled, Eq. 3 will always have a real solution.

3. Assume that the opacity per unit mass is κ . Find the maximum luminosity the object can have without ejecting the outer layers (assume pure hydrogen completely ionized). This luminosity is called the Eddington luminosity

The maximum luminosity an object can have before ejecting its outer layers has already been found in Eq. 1. Particularly, in the case of completely ionized hydrogen, $\kappa = \sigma_T/m_H$ being σ_T the Thompson scattering cross section and m_H the mass of the hydrogen atom, thus, any luminosity fulfilling the following condition will imply the ejection of the envelope:

$$L > \frac{4\pi G M c}{\kappa} = \frac{4\pi G M m_H c}{\sigma_T}$$

Therefore, the maximum luminosity before ejecting the envelope is:

$$L_{EDD} = \frac{4\pi G M m_H c}{\sigma_T}$$

which simplifies the expression of the terminal velocity in cases where $L > L_{EDD}$ to:

$$v_t^2 = \frac{2GM}{R} \left(\frac{L}{L_{EDD}} - 1 \right)$$

In the case of the Sun, its Eddington luminosity is:

$$\begin{aligned} L_{\odot} &= 4 \times 10^{33} \text{ erg/s} ; M_{\odot} = 2 \times 10^{33} \text{ g} ; G = 6.674 \times 10^{-8} \text{ cm}^3 \text{g}^{-1} \text{s}^{-2} \\ m_H &= 1.6733 \times 10^{-24} \text{ g} ; c = 2.99792458 \times 10^{10} \text{ cm/s} ; \sigma_T = 6.65 \times 10^{-25} \text{ cm}^2 \\ L_{EDD} &= 1.265 \times 10^{38} \text{ erg/s} = 3.16 \times 10^5 L_{\odot} \end{aligned}$$