

Univariate Real Analysis

Muchang Bahng

Spring 2025

Contents

1	The Real Numbers	5
1.1	Dedekind Completeness	6
1.2	Cauchy Completeness	9
1.3	Least Upper Bound Completeness	11
1.4	Nested Intervals Completeness	13
1.5	Properties of the Real Line	14
1.6	Exponentials, Roots, and Logarithms	16
1.7	Extended Reals	21
1.8	Hyperreals	22
1.9	Some Algebraic Inequalities	22
1.10	Exercises	24
2	Euclidean Topology	52
2.1	Exercises	55
3	Sequences in Euclidean Space	75
3.1	Convergence Theorems from Completeness and Order	76
3.2	Arithmetic	77
3.3	Limsup and Liminf	80
3.4	Convergence Tests for Real Series	83
3.5	Ratio and Root Tests	87
3.6	Euler's Number and Trigonometric Functions	89
3.7	Exercises	91
4	Limits and Continuity of Functions	101
4.1	Limits of Functions	101
4.2	Asymptotic Behavior of Functions	106
4.2.1	Approximations of Functions	109
4.3	Continuous Functions	111
4.3.1	Intermediate and Extreme Value Theorem	113
4.3.2	Inverse Function Theorem	114
4.4	Uniform Continuity	115
4.5	Lipshitz and Holder Continuity	119
4.6	Discontinuity	120
4.7	Exercises	123
5	Differentiation of Single-Variable Functions	125
5.1	Rules of Differentiation	130
5.2	Theorems of Differentiable Functions	132
5.3	Extrema and Concavity	135
5.4	Theorems of Continuously Differentiable Functions	137

5.5	Higher Order Derivatives	139
5.5.1	Taylor's Formula	141
5.6	Exercises	145
6	Riemann Integration	149
6.1	Axiomatic Properties of the Integral	151
6.2	Conditions for Riemann Integrability	153
6.3	Further Properties of the Riemann Integral	158
6.4	The Fundamental Theorem of Calculus	160
6.5	Change of Variable	163
6.6	Integration by Parts	164
6.7	Improper Integrals	164
6.8	Integration over Paths and Rectifiable Curves	167
7	Sequences of Functions	172
7.1	Uniform Convergence	174
7.2	Limits of Uniformly Convergent Functions	178
7.3	Equicontinuous Families	181
7.4	The Stone-Weierstrass Theorem	185
7.5	Approximation of the Identity	186

Let's first talk about why we need analysis in general in the first place. Algebra allows us to define certain algebraic structures, which are essentially sets with operations. These operations are defined to have a finite number of arguments. For example, let's take a look at the negation $x \mapsto -x$ and the addition $x, y \mapsto x + y$ operations in a group G . We can compose these operations up to any finite length n , removing the parentheses due to associativity, but note that the “sum” below is not a single operation. It is a composition of $n - 1$ operations.

$$x_1 + x_2 + \dots + x_n \in G \quad (1)$$

$$-(-(\dots(-x))) \in G \quad (2)$$

This is still well defined due to closure, but what if we wanted to do this an infinite number of times?

$$x_1 + x_2 + \dots =? \quad (3)$$

$$\dots(-(-x)) =? \quad (4)$$

For someone who has learned about sequences and series in high school, this may not be a big jump in logic, but it is. The objects above are not even well-defined and trying to define them with algebraic tools is manifested in the famous Zeno's paradox. So we simply need to add more tools in order to define these new mathematical objects, which we call *series*. To define series, we need to first define sequences. Can we do this with algebra? Yes,¹ since we can simply model it as a function.

Definition 0.1 (Sequence)

A sequence is a function $f : \mathbb{N} \rightarrow X$. We usually denote a sequence by writing out the first few terms of the sequence, followed by an ellipsis.

$$a_1 = f(1), a_2 = f(2), \dots \quad (5)$$

or as an indexed set over the naturals $\{a_i\}_{i \in \mathbb{N}}$.

Therefore, we can consider series as a sequence of finite sums, each element which is well-defined.

$$x_1, x_1 + x_2, x_1 + x_2 + x_3, \dots \quad (6)$$

For any $n \in \mathbb{N}$, we can get the value of $a_n = \sum_{i=1}^n a_i$, but can we say something about the limiting behavior of a_n ? That is, maybe we can just slap a value x onto this series such that it doesn't “break” any of the rules we have in the finite sense. Unfortunately, it is not possible to define such values for all series, but it is possible for some of them, which we call *convergent series*. To rigorously determine which ones are convergent and which ones are not, we need the tools of topology. Defining the concept of sequences that model infinitely composed operations is what allows us to define differentiation and integration.

Great, we've motivated the need for analysis, but before jumping straight into real analysis, let's talk about what analysis in general works with. It studies functions of the form $f : X \rightarrow Y$, and minimally both X, Y must be *Banach spaces*, i.e. complete normed vector spaces over some field \mathbb{F} . Almost all flavors of analysis, including real (\mathbb{R}), complex (\mathbb{C}), multivariate (\mathbb{R}^n), p-adic, and functional (infinite-dimensional Banach spaces) analysis require *at least* a Banach space structure. Why are Banach spaces so great? Well if we were to define convergence in X or Y , then it only makes sense to talk about convergence with respect to a topology. So X, Y must at least be topological spaces. It would also be bad if we were to take a sequence in X and find out that it converges to some element outside of X . Therefore, we want a notion of *completeness* in the sense that all sequences that “get closer,” i.e. Cauchy sequences, actually converge in X . Unfortunately, while convergence of sequences is preserved under homeomorphisms (and is thus a topological property), convergence of Cauchy sequences is not.² Furthermore, the notion of uniform convergence is a metric space

¹In fact, we don't even need algebra, just set theory.

²Consider the sequence $a_n = 1/(n+1)$ in $(0, 1)$ and the map $f(x) = 1/x$ to the set $(1, +\infty)$. a_n is Cauchy but $f(a_n)$ is not.

property, not a topological one. Therefore, the concept of distances is crucial to the construction of analysis. As for the norm, I'm still not sure why we need this.³

But in college courses such as real and complex analysis, why do we say we work over the *fields* \mathbb{R} and \mathbb{C} rather than the Banach spaces \mathbb{R} and \mathbb{C} ? This is because of the following theorem.

Theorem 0.1 ()

Every field \mathbb{F} is a 1-dimensional vector space over itself.

Therefore, when we talk about the *field* \mathbb{R} , we are really treating it as a vector space \mathbb{R} over the field \mathbb{R} .⁴ Every other structure beyond this is a “bonus” property that gives us extra tools to prove stronger properties. The most notable is the total ordering on \mathbb{R} , which allows us to define upper/lower bounds and other real-analysis specific theorems like the intermediate value theorem or the mean value theorem. Other structures include the inner product or the measure.

Now that we've taken in the big picture, for each type of analysis, we should construct the underlying relevant Banach space. At the very least, we can with the tools of set theory and algebra define the rationals \mathbb{Q} as an ordered field over the quotient space $\mathbb{Z} \times \mathbb{Z} / \sim$. Furthermore, \mathbb{Q} itself is a normed vector space (over \mathbb{Q})⁵ and the only thing we need now is completeness.

1. If the norm on \mathbb{Q} is defined as the normal absolute value (Euclidean norm), completing it gives \mathbb{R} as an ordered field which also has a compatible order as that of \mathbb{Q} . We study functions mapping to and from \mathbb{R} with *single-variable real analysis*.
2. If we take the *p-adic* norm, then completing it with respect to this gives the *p-adic numbers*, which also forms a field but loses the ordering. We deal with functions over the p-adics with *p-adic analysis*.
3. We can construct \mathbb{C} by taking \mathbb{R}^2 and endowing it with a bit more structure. We get *complex analysis*.
4. We can construct \mathbb{R}^n and \mathbb{C}^n by easily defining its vector space structure and then endowing it with a norm, and showing that it is complete with respect to the norm-induced metric. This is known as *multivariate analysis*.
5. With all these defined, we can define Banach function spaces like L^p and perform analysis on operators $f : L^p \rightarrow L^q$. This is *functional analysis*.

³Aspinwall and Ng told me this, but I'm not sure why. The Frechet derivative seems like it can be purely defined with a metric.

⁴Thanks to Prof. Lenny Ng for clarifying this.

⁵Note that while we define the norm and metric to usually map to \mathbb{R}^+ , \mathbb{R} isn't even defined yet and so to avoid circular definitions, we define the norm on the rationals to have codomain \mathbb{Q} .

1 The Real Numbers

By constructing \mathbb{Q} and its topology in my algebra and topology notes, we can talk about convergence. The first question to ask (if you were the first person inventing the reals) is “how do I know that there exists some other numbers at all?” The first clue is trying to find the side length of a square with area 2. As we see, this number is not rational.

Theorem 1.1 ($\sqrt{2}$ is Not Rational)

There exists no $x \in \mathbb{Q}$ s.t. $x^2 = 2$.

Proof. Assume such a number $x = p/q$ exists, where $\gcd(p, q) = 1$. Then, by the field axioms of \mathbb{Q} , we can deduce that

$$2 = \frac{p^2}{q^2} \implies 2q^2 = p^2 \quad (7)$$

This implies that $2 \mid p$, so we have $p = 2p_0$, and we can write $2q^2 = 4p_0^2$. Dividing both sides by 2, we get $q^2 = 2p_0^2$, which implies that $2 \mid q$. This contradicts the fact that p and q are coprime.

We can “imagine” that a square with area 2 certainly exists, but the fact that its side length is undefined is certainly unsettling. I don’t know about you, but I would try to “invent” $\sqrt{2}$. We can do this in 4 distinct ways, though some may be more similar than others.

1. *Dedekind Completeness.* I would try and generate new elements by taking a specific *cut*—a partition into two sets such that elements of one set is always greater than that of the other—and seeing which elements lie right in between these cuts. We will often see that we can always find a cut for every rational, but there are additional cuts for each irrational number. This is the idea behind *Dedekind completeness*.
2. *Cauchy Completeness.* I write out the decimal expansion one by one, which gives our first exposure to sequences.

$$1, 1.4, 1.41, 1.414, \dots \quad (8)$$

It is clear that on \mathbb{Q} , this sequence does not converge. Our intuition tells that if the terms get closer and closer to each other, they must be getting closer and closer to *something*, though that something is not in \mathbb{Q} . This motivates the definition for *Cauchy completeness*.

3. *Nested Interval Completeness.* I would write out maybe some nested intervals so that $\sqrt{2}$ *must* lie within each interval.

$$[1, 2] \supset [1.4, 1.5] \supset [1.41, 1.42] \supset \dots \quad (9)$$

This motivates the definition of *nested-interval completeness*.

4. *Least Upper Bound Completeness.* I would define the set of all rationals such that $x^2 < 2$, and try to define $\sqrt{2}$ as the max or supremum of this set. We will quickly find that neither the max nor the supremum exists in \mathbb{Q} , and this motivates the definition for *least upper bound completeness*.

All four of these methods points at the same intuition that there should not be any “gaps” or “missing points” in the set that we will construct to be \mathbb{R} , which is the general notion of *completeness*. This contrasts with the rational numbers, whose corresponding number line has a “gap” at each irrational value. Even though constructing the reals with one method is sufficient, knowing the different flavors in which completeness manifests is very useful. It allows us to view properties of \mathbb{R} through different lens.

The main division between these four properties is that the first two are methods to directly *construct* the reals from \mathbb{Q} , while the latter two are more *axiomatic properties* that we use to verify completeness for a given set. In the next two sections, we will take the rationals and add on extra elements using Dedekind cuts and Cauchy sequences. However, it isn’t as conventional (though possible) to construct them as the single point contained in a sequence of nested intervals nor as a supremum of all upper-bounded sets. In fact, an

alternative way to construct the reals is to define it axiomatically—as a totally ordered field with either the LUB property or the nested interval property.⁶

Therefore, in the following sections, we will

1. first define the relevant notion of completeness,
2. show that the rationals are not complete
3. and then construct the completed version of the rationals as a version of the reals.

Once we have done this for all three versions, we will unify them by proving they are all equivalent.

There is a second essential property of the reals that is not talked about as often is the Archimidean principle.

Definition 1.1 (Archimidean Principle)

An ordered ring $(X, +, \cdot, \leq)$ that embeds the naturals \mathbb{N}^a is said to obey the **Archimidean principle** if given any $x, y \in X$ s.t. $x, y > 0$, there exists an $n \in \mathbb{N}$ s.t. $\iota(n) \cdot x > y$. Usually, we don't care about the canonical injection and write $nx > y$.

^aas in, there exists an ordered ring homomorphism $\iota : \mathbb{N} \rightarrow X$

Lemma 1.1 (Rationals are Archimidean)

\mathbb{Q} satisfies the Archimidean principle.

Proof. Take any $x = p_1/q_1, y = p_2/q_2 \in \mathbb{Q}$. Then, take $n = q_1 p_2$, and we get

$$nx = q_1 p_2 \frac{p_1}{q_1} = p_1 p_2 > \frac{p_2}{q_2} = y \iff p_1 > \frac{1}{q_2} \quad (10)$$

which must be true since $p_1 \geq 1 \geq \frac{1}{q_2}$.

Usually, we just construct \mathbb{R} right out of \mathbb{Q} , and it turns out that the Archimidean principle just trivially follows. However, if we construct \mathbb{R} axiomatically (without the rationals), it needs to be stated. In this axiomatic formulation, we will find that certain types of completeness—like Dedekind completeness—actually *implies* the Archimidean principle, while others—like Cauchy and nested-intervals completeness—does not. Therefore, in a sense, Dedekind completeness is the “strongest” form of completeness.

1.1 Dedekind Completeness

This is an explicit construction from the rationals.

Definition 1.2 (Dedekind Cut)

A **Dedekind cut** is a partition of the rationals $\mathbb{Q} = A \sqcup A'$ satisfying the three properties.^a

1. $A \neq \emptyset$ and $A \neq \mathbb{Q}$.^b
2. $x < y$ for all $x \in A, y \in A'$.
3. The maximum element of A does not exist in \mathbb{Q} .

The minimum of A' may exist in \mathbb{Q} , and if it does, the cut is said to be **generated** by $\min A'$.

^aThis can really be defined for any totally ordered set.

^bBy relaxing this property, we can actually complete \mathbb{Q} to the extended real number line.

⁶In fact, you also need the Archimidean principle, but we'll talk about this later.

Note that in \mathbb{Q} , there will be two types of cuts:

1. ones that are generated by rational numbers, such as

$$A = \{x \in \mathbb{Q} \mid x < 2/3\}, A' = \{x \in \mathbb{Q} \mid x \geq 2/3\} \quad (11)$$

2. and the ones that are not

$$A = \{x \in \mathbb{Q} \mid x^2 < 2\}, A' = \{x \in \mathbb{Q} \mid x^2 \geq 2\} \quad (12)$$

We can intuitively see that the set of all Dedekind cuts (A, A') will “extend” the rationals into a bigger set. We can then define some operations and an order to construct this into an ordered field, and finally it will have the property that we call “completeness.”

Definition 1.3 (Dedekind Completeness)

A totally ordered algebraic field \mathbb{F} is **Dedekind-complete** if every Dedekind cut of \mathbb{F} is generated by an element of \mathbb{F} .

Lemma 1.2 (Rationals are Not Dedekind-Complete)

\mathbb{Q} is not Dedekind-complete.

Proof. Take a look at the cut

$$A = \{x \in \mathbb{Q} \mid x^2 < 2\}, A' = \{x \in \mathbb{Q} \mid x^2 \geq 2\} \quad (13)$$

We wish to show that A' has no minimum. Assume that it does, and call it p . Then, define

$$q := p - \frac{p^2 - 2}{p + 2} = \frac{2p + 2}{p + 2} \quad (14)$$

We can see that $p > 2 \implies p^2 - 2 > 0 \implies q < p$. But we also see that

$$q^2 - 2 = \frac{2(p^2 - 2)}{(p + 2)^2} \implies q^2 > 2 \quad (15)$$

Therefore, we have found a $q \in A'$ that is smaller than p , a contradiction.

These Dedekind cuts are simply subsets of the power set of \mathbb{Q} , which always exists due to the power set axiom. Therefore, we can simply use the axiom of restricted comprehension (?) to create a well-defined set of Dedekind cuts.

Definition 1.4 (Reals as the Dedekind-Completion of Rationals)

Let \mathbb{R}_D be the set of all Dedekind cuts A^a of \mathbb{Q} .

$$\mathbb{R}_D := \{A \in 2^{\mathbb{Q}} \mid (A, A^c) \text{ is a Dedekind cut}\} \quad (16)$$

By doing this we can intuitively think of a real number as being represented by the set of all smaller rational numbers. Let $A, B \in \mathbb{R}_D$ two Dedekind cuts. Then, we define the following order and operations.

1. *Order.* $A \leq_{\mathbb{R}} B \iff A \subset B$.
2. *Addition.* $A +_{\mathbb{R}} B := \{a +_{\mathbb{Q}} b \mid a \in A, b \in B\}$.
3. *Additive Identity.* $0_{\mathbb{R}} := \{x \in \mathbb{Q} \mid x < 0\}$.

4. *Additive Inverse.* $-B := \{a - b \mid a < 0, b \in (\mathbb{Q} \setminus B)\}$.
5. *Multiplication.* If $A, B \geq 0$, then we define $A \times_{\mathbb{R}} B := \{a \times_{\mathbb{Q}} b \mid a \in A, b \in B, a, b \geq 0\} \cup 0_{\mathbb{R}}$. If A or B is negative, then we use the identity $A \times B = -(A \times_{\mathbb{R}} -B) = -(-A \times_{\mathbb{R}} B) = (-A \times_{\mathbb{R}} -B)$ to convert A, B to both positives and apply the previous definition.
6. *Multiplicative Identity.* $1_{\mathbb{R}} = \{x \in \mathbb{Q} \mid x < 1\}$.
7. *Multiplicative Inverse.* If $B > 0$, $B^{-1} := \{a \times_{\mathbb{Q}} b^{-1} \mid a \in 1_{\mathbb{R}}, b \in (\mathbb{Q} \setminus B)\}$. If B is negative, then we compute $B^{-1} = -((-B)^{-1})$ by first converting to a positive number and then applying the definition above.

We claim that $(\mathbb{R}, +_{\mathbb{R}}, \times_{\mathbb{R}}, \leq_{\mathbb{R}})$ is a Dedekind-complete totally ordered field, and the canonical injection $\iota : \mathbb{Q} \rightarrow \mathbb{R}$ defined

$$\iota(q) = \{x \in \mathbb{Q} \mid x < q\} \quad (17)$$

is an ordered field isomorphism.

^aFor convenience we can uniquely represent (A, A') with just A since $A' = \mathbb{Q} \setminus A$.

Proof.

By the canonical injections $\mathbb{N} \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \rightarrow \mathbb{R}_D$, we can talk about whether this set has the Archimedean property. By construction, Archimedean is trivial since \mathbb{R}_D contains \mathbb{Q} which is Archimedean.

Theorem 1.2 (Dedekind Reals is Archimedean)

\mathbb{R}_D satisfies the Archimedean principle.

Proof. \mathbb{R}_D contains \mathbb{Q} .

Definition 1.5 (Axiomatic Construction of Dedekind-Reals)

\mathbb{R}'_D is a totally ordered field that is Dedekind complete.^a

^aJust for this section, I will denote \mathbb{R}' as reals constructed axiomatically.

Theorem 1.3 (Axiomatic Dedekind Reals is Archimedean)

\mathbb{R}'_D satisfies the Archimedean principle.

Proof. Assume that this property doesn't hold. Then for any fixed x , $nx < y$ for all $n \in \mathbb{N}$. Consider the set

$$A = \bigcup_{n \in \mathbb{N}} (-\infty, nx), \quad B = \mathbb{R} \setminus A \quad (18)$$

A by definition is nonempty, and B is nonempty since it contains y . Then, we can show that $a \in A, b \in B \implies a < b$ using proof by contradiction. Assume that there exists $a' \in A, b' \in B$ s.t. $a' > b'$. Since $a' \in A$, there exists a $n' \in \mathbb{N}$ s.t. $a' \in (-\infty, n'x) \iff a' < n'x$. But by transitivity of order, this means $b' < n'x \iff b' \in (-\infty, n'x) \implies b' \in A$.

Going back to the main proof, we see that A is upper bounded by y , and so by the least upper bound property it has a supremum $z = \sup A$.

1. If $z \in A$, then by the induction principle^a $z + x \in A$, contradicting that z is an upper bound.
2. If $z \notin A$, then by the induction principle^b $z - x \notin A \implies z - x \in B$. Since every element of B upper bounds A and since $x > 0$, this means that $z - x < z$ is a smaller upper bound of A , contradicting that z is a least upper bound.

Therefore, it must be the case that $nx > y$ for some $n \in \mathbb{N}$.

^aNote that \mathbb{N} is defined recursively as $1 \in \mathbb{N}$ and if $n \in \mathbb{N}$, then $n + 1 \in \mathbb{N}$.

^bThe contrapositive of the recursive definition of \mathbb{N} is: if $n \notin \mathbb{N}$, then $n - 1 \notin \mathbb{N}$.

1.2 Cauchy Completeness

In many cases we are not working with ordered fields, and so different types of completeness may be more useful. In actual practice, you tend to use Cauchy completeness which only assumes a metric structure.

Definition 1.6 (Cauchy Sequence)

A sequence $(x_n)_n$ in a metric space (X, d) is a **Cauchy sequence** if $\forall \epsilon > 0, \exists N \in \mathbb{N}$ s.t.

$$n, m \geq N \implies d(x_n, x_m) < \epsilon \quad (19)$$

To motivate this definition, note that in a general topological space X , we can define convergence of a sequence $x_n \rightarrow x$ perfectly fine. However, take some subset $U \subset X$, and let x be a limit point of U . In this case, x_n does not converge in U , but it does converge to something outside of U —namely, $x \in X$. This is similar to $\mathbb{Q} \subset \mathbb{R}$, where x is an irrational point. However, we are trying to *construct* \mathbb{R} , so Cauchy convergence allows us to speak of convergence without actually referring to *what* a sequence is converging to.

Note that it is not sufficient to say that a sequence is Cauchy by claiming that each term becomes arbitrarily close to the preceding term. That is,

$$\lim_{n \rightarrow \infty} d(x_{n+1}, x_n) = 0 \quad (20)$$

Example 1.1 (Adjacent Terms Converging Doesn't Imply Sequence is Cauchy)

For example, look at the sequence

$$a_n = \sqrt{n} \implies a_{n+1} - a_n = \frac{1}{\sqrt{n+1} + \sqrt{n}} < \frac{1}{2\sqrt{n}} \quad (21)$$

However, it is clear that a_n gets arbitrarily large, meaning that a finite interval can contain at most a finite number of terms in $\{a_n\}$.

It is often more convenient to think of the limit of the *diameter* of rest of the sequence. That is, a sequence is Cauchy if

$$\lim_{n \rightarrow \infty} \text{diam}\{x_m\}_{m \geq n} = 0 \quad (22)$$

It is trivial that convergence implies Cauchy convergence, but the other direction is not true. Therefore, we would like to work in a space where these two are equivalent, and this is called completeness. Therefore, we can construct the reals as equivalence classes over Cauchy sequences. Rather than using the order, we take advantage of the metric.

Definition 1.7 (Cauchy Completeness)

A metric space (X, d) is **Cauchy complete** if every Cauchy sequence in that space converges to an element in X .

\mathbb{Q} is not Cauchy-complete. Let a_n be the largest number x up to the n th decimal expansion such that x^2

does not exceed 2. The first few terms are

$$1.4, 1.41, 1.414, \dots \quad (23)$$

In this case, we can see that this is Cauchy since at the n th element and on, the first n decimal places are kept fixed and so the most that the rest of the sequence can deviate by is 10^{-n} .

Definition 1.8 (Reals as the Cauchy-Completion of the Rationals)

Let \mathbb{R}_C be the quotient space of all Cauchy (under the Euclidean metric) sequences (x_n) of rational numbers with the equivalence relation $(x_n) = (y_n)$ iff their difference tends to 0.^a That is, for every rational $\epsilon > 0$, there exists an integer N s.t. for all naturals $n > N$, $|x_n - y_n| < \epsilon$.

1. *Order.* $(x_n) \leq_{\mathbb{R}} (y_n)$ iff $x = y$ or there exists $N \in \mathbb{N}$ s.t. $x_n \leq_{\mathbb{Q}} y_n$ for all $n > N$.
2. *Addition.* $(x_n) + (y_n) := (x_n + y_n)$.
3. *Additive Identity.* $0_{\mathbb{R}} := (0_{\mathbb{Q}})$.
4. *Additive Inverse.* $-(x_n) := (-x_n)$.
5. *Multiplication.* $(x_n) \times_{\mathbb{R}} (y_n) = (x_n \times_{\mathbb{Q}} y_n)$.
6. *Multiplicative Identity.* $1_{\mathbb{R}} := (1)$.
7. *Multiplicative Inverse.* $(x_n)^{-1} := (x_n^{-1})$.

We claim that $(\mathbb{R}, +_{\mathbb{R}}, \times_{\mathbb{R}}, \leq_{\mathbb{R}})$ is a totally ordered field, and the canonical injection $\iota : \mathbb{Q} \rightarrow \mathbb{R}$ defined

$$\iota(q) = (q) \quad (24)$$

is an ordered field isomorphism. Finally, by construction \mathbb{R} is Cauchy-complete.

^aThis equivalence class reflects that the same real number can be approximated in many different sequences. In fact, this shows *by definition* that $1, 1, \dots$ and $0.9, 0.99, 0.999, \dots$ are the same number!

Proof.

Theorem 1.4 (Cauchy Reals is Archimidean)

\mathbb{R}_C satisfies the Archimedean principle.

Proof. \mathbb{R}_C contains \mathbb{Q} , which is Archimidean.

The best thing about Cauchy completeness is that we can just take \mathbb{Q}^n to create \mathbb{R}^n . It becomes quite general. However, note that first, Cauchy completion depends on *which* metric you use to complete it.

Example 1.2 (P-adic Numbers)

Let p be a prime number. For a non-zero rational $x = p^k \cdot \frac{a}{b}$ where $p \nmid a, b$, define the **p-adic norm**^a as

$$|x|_p := p^{-k}, \quad |0|_p := 0 \quad (25)$$

The **p-adic numbers** \mathbb{Q}_p are the Cauchy completion of \mathbb{Q} with respect to the p-adic metric $d_p(x, y) = |x - y|_p$. This set not only does not satisfy the Archimidean principle; it doesn't even have a natural ordering!

^aThis measures divisibility by p : the more p divides x , the smaller $|x|_p$. For example, $|8|_2 = 2^{-3} = 1/8$ and $|3|_2 = 1$.

Definition 1.9 (Axiomatic Construction of Cauchy-Reals)

\mathbb{R}'_D is a totally ordered field that is Cauchy complete and that satisfies the Archimidean principle.

Note that we require the extra Archimidean assumption in the axiomatic construction.

Example 1.3 (Ordered Cauchy-Complete Fields that are Not Archimidean)

Provide examples of ordered, Cauchy-complete fields that are not Archimedean. Hyperreals?

1.3 Least Upper Bound Completeness**Definition 1.10 (Least Upper Bound Property)**

A totally ordered algebraic field \mathbb{F} (must it be a field?) is **least-upper-bound complete**, or is said to satisfy the **least upper bound (LUB) property**, if every nonempty set of \mathbb{F} having an upper bound must have a least upper bound (supremum) in \mathbb{F} .

Theorem 1.5 (LUB is Equivalent to GLB)

A set (\mathbb{F}, \leq) has the least upper bound property iff it has the *greatest lower bound property*, which states that every set bounded below has a greatest lower bound.

Proof. We will prove one direction since the other is the same logic. Let $S \subset X$ be a nonempty set that is bounded below by some $l \in X$. Let $L \subset X$ be the set of all lower bounds of S . Since l exists, it is nonempty. Furthermore, L is bounded above by any element of S . Due to LUB property L has a least upper bound, call it $z = \sup L$. We claim that $z = \inf S$.

1. z is a lower bound of S . Assume that it is not. Then there exists $s \in S$ s.t. $s < z$. But by construction s is an upper bound for L and so z is not the *least* upper bound, a contradiction.
2. z is a *greatest* lower bound. Assume that z is not. Then there exists a $z' \in X$ s.t. $z < z' \leq s$ for all $s \in S$. But since z', z are lower bounds, this means $z, z' \in L$ by definition and $z < z'$ contradicts the fact that z is an upper bound of L .

We are done.

\mathbb{Q} does not satisfy the least upper bound property, but proving this can be tricky for the first time. We state this as a lemma.

Theorem 1.6 (Rationals Doesn't Satisfy LUB Property)

\mathbb{Q} does not satisfy the LUB property.

Proof. Assume it does, and let us denote

$$p := \sup\{x \in \mathbb{Q} \mid x^2 < 2\} \in \mathbb{Q} \quad (26)$$

The key here is to find another rational that we can always “squeeze” in between p and 2. This can be done with the Archimidean principle, which is already satisfied in \mathbb{Q} . Since we have proved that there exists no rational that squares to 2, we only need to consider the two cases.

1. $p^2 < 2$. Take $\epsilon \in \mathbb{Q}$ so small that

$$p^2 + 2p\epsilon + \epsilon^2 = (p + \epsilon)^2 < 2 \quad (27)$$

- To show complete steps, we can see that by density of reals, there exists some rational r s.t. $0 < r < 2 - p^2$. Therefore, we can invoke Archimidean principle to find a $n \in \mathbb{N}$ s.t. $\epsilon = p/n < 2p/n < r$. Therefore, p is not an upper bound, so this cannot be true.
2. $p^2 > 2$. We can again take an $\epsilon \in \mathbb{Q}$ so small that

$$p^2 - 2p\epsilon + \epsilon^2 = (p - \epsilon)^2 > 2 \quad (28)$$

Therefore, p is not least, so this cannot be true.

Definition 1.11 (Axiomatic Construction of Reals with LUB Property)

\mathbb{R}'_I is a totally ordered field that satisfies the least upper bound property.

Note that we don't need to explicitly assume Archimidean principle here. The LUB property is strong enough that it implies Archimidean!

Theorem 1.7 (LUB Property Implies Archimidean)

\mathbb{R}'_I is Archimidean.

Now let's see how our previous constructions of the reals relate to the LUB property.

Theorem 1.8 (Dedekind Completed Reals Satisfies LUB Property)

\mathbb{R}_D satisfies the least upper bound property.

Proof. Let \mathcal{A} be a nonempty subset of \mathbb{R}_D bounded from above by T . Then, $\forall A \in \mathcal{A}$, $A := (A, A^c)$ is a Dedekind cut, and we can define

$$(B, B') := \left(\bigcup_{A \in \mathcal{A}} A, \bigcap_{A \in \mathcal{A}} A^c \right) \quad (29)$$

We claim that this is a Dedekind cut.

1. First, it is nonempty set $\mathcal{A} \neq \emptyset$ and so for each $A \in \mathcal{A}$, $A \subset \mathbb{Q}$ is nonempty. It is also not all of \mathbb{Q} since T is an upper bound of \mathcal{A} , and so $T \geq a \forall a \in A \forall A \in \mathcal{A}$, which implies that

$$T \notin \bigcup_{A \in \mathcal{A}} A \quad (30)$$

2. Now let $x \in B, y \in B'$. Then, $x \in A_0$ for some $A_0 \in \mathcal{A}$, and since y is in the intersection of all the corresponding A^c , it must be in the corresponding A_0^c . Therefore we invoke the Dedekind cut property of (A_0, A_0^c) and see $x < y$.
3. Finally, for the sake of contradiction, let $m \in \mathbb{Q}$ be the maximum of B . Then, $m \in A^*$ for some $A^* \in \mathcal{A}$. But m is an upper bound for the whole B , so this means that $m = \max\{A^*\}$, which contradicts the fact that lower cut cannot have a rational maximum.

Now we claim that B is the supremum. It is an upper bound since

$$A \subset B = \bigcup_{A \in \mathcal{A}} A \quad \forall A \in \mathcal{A} \quad (31)$$

To prove least, we should see that if (S, S^c) is another upper bound of \mathcal{A} , then $A \subset S$ for all $A \in \mathcal{A} \implies B = \bigcup_{A \in \mathcal{A}} A \subset S$, which establishes that B is least.

1.4 Nested Intervals Completeness

The next flavor we present is nested-intervals completeness. This is the least popular way to construct the reals, and it is used more as a post-hoc tool to analyze the reals after you construct it using either of the two previous methods.

Definition 1.12 (Nested Interval Completeness)

Let \mathbb{F} be a totally ordered algebraic field. Let $I_n = [a_n, b_n]$ ($a_n < b_n$) be a sequence of decreasing nested intervals that are

1. closed,
2. bounded,
3. nested, $I_1 \supset I_2 \supset I_3 \supset \dots$
4. and decreasing to 0 in the sense that $\lim_{n \rightarrow \infty} b_n - a_n = 0$.

\mathbb{F} is **nested-interval complete** if the intersection of all of these intervals I_n contains exactly one point.

$$\bigcup_{n=1}^{\infty} I_n \in \mathbb{F} \quad (32)$$

Note that defining nested intervals requires only an ordered field. One may look at this and try to ask if this is a specific instance of the following conjecture: The intersection of a nested sequence of nonempty closed sets in a topological space has exactly 1 point. This claim may not even make sense, actually. If we define nested in terms of proper subsets, then for a finite topological space a sequence cannot exist since we will run out of open sets and so this claim is vacuously true and false. If we allow $S_n = S_{n+1}$ then we can just select $X \supset X \supset \dots$, which is obviously not true. However, a slightly weaker claim is that every proper nested non-empty closed sets has a non-empty intersection is a consequence of compactness.

Theorem 1.9 (Rationals are Not Nested-Interval Complete)

\mathbb{Q} is not nested-interval complete.

Proof. This is a nice proof that uses a class method of algorithmically selecting nested intervals that satisfy a following property. This trick will be used many times in analysis.

For the sake of contradiction, let us assume that \mathbb{Q} satisfies nested intervals completeness. Since \mathbb{Q} is countable, enumerate it as q_1, q_2, \dots

1. Choose any closed interval I_1 of length 1 with rational endpoints that doesn't contain q_1 .
2. Now partition I_1 into two segments of equal length, and choose I_2 to be the segment that doesn't contain q_2 .

At this point, I_n is a closed interval with rational endpoints of length 2^{-n} that cannot contain q_n . Therefore, $\bigcap_n I_n$ cannot contain any $q_n \in \mathbb{Q}$. But this contradicts our assumption of nested interval completeness.

One may ask: what is the relationship between LUB property and nested-intervals completeness? It turns out that they are equivalent.

Theorem 1.10 (LUB is Equivalent to Nested Interval Completeness in Reals)

Listed.

1. \mathbb{R}'_L satisfies nested intervals completeness.
2. \mathbb{R}'_I satisfies least upper bound completeness.

Proof. Listed.

1. Note that $\{a_n\}$ is bounded above by b_1 . Therefore by LUB property it must have a supremum, call it $x = \sup_n \{a_n\}$. Then, we see that $a_n \leq x \leq b_n$ for all n , and so x is in the intersection.

Theorem 1.11 (Cantor's Intersection Theorem)

\mathbb{R} is nested-interval complete.

Proof. We prove this by first proving the claim that given nested, closed, and bounded sets I_n (not even necessarily intervals), then

$$\bigcap_{n=1}^{\infty} I_n \neq \emptyset \quad (33)$$

Suppose this is not true. For every $x \in \mathbb{R}$, there exists a n_x s.t. $x \notin I_{n_x}$ (and all later I_m for $m > n_x$). Let $O_n = I_n^c$ open sets. Then, $\mathbb{R} \subset \bigcup_n O_n$. In particular, $I_1 \subset \bigcup_n O_n$. But I_1 is closed and bounded. So we can extract a finite subcover. $O_{n_1}, O_{n_2}, \dots, O_{n_m}$ (ordered $n_1 < n_2 < \dots < n_m$). Then since O_n are increasing, $I_1 \subset O_{n_m} = I_{n_m}^c$. But $I_{n_m} \subset I_1$, a contradiction.

Now with this, we know that because the limits of the endpoints of the intervals go to 0, then there cannot be more than 2 points in the intersection. Thus there must be 1 unique point.

Definition 1.13 (Axiomatic Construction of Reals with Nested Intervals)

\mathbb{R}_I is a totally ordered field that

1. satisfies the nested intervals completeness, and
2. satisfies the Archimidean principle.

1.5 Properties of the Real Line

Perfect, now all that remains is to unite the two constructions of the reals.

Theorem 1.12 (Dedekind and Cauchy Complete Reals are Isomorphic)

Given that \mathbb{R}_D is the Dedekind-completed version of the rationals and \mathbb{R}_C is the Cauchy-completed version of the rationals, we claim that the two are isomorphic as ordered fields.

$$\mathbb{R}_D \simeq \mathbb{R}_C \quad (34)$$

Proof.

Therefore, it doesn't really matter which one we talk about, and we can refer to *the* real numbers as a single set. Great! Now we can finally feel satisfied about defining metrics, norms, and inner products as mappings to the codomain \mathbb{R} .

Definition 1.14 (Reals (as Construction from Rationals))

The **reals** \mathbb{R} is the totally ordered complete Archimedean field constructed as the completion^a of \mathbb{Q} .

^aeither one

So far, we have taken the completion of the rationals as our main mode of construction. However, we can take an axiomatic approach, and it turns out that there is only one set, up to isomorphism, that satisfies all

these properties.

Theorem 1.13 (Axiomatic Definition of Reals)

The **real numbers**, denoted \mathbb{R} , is any totally ordered complete Archimedean field. \mathbb{R} is unique up to field isomorphism. That is, if two individuals construct two ordered complete Archimedean fields \mathbb{R}_A and \mathbb{R}_B , then

$$\mathbb{R}_A \simeq \mathbb{R}_B \quad (35)$$

Proof. The proof is actually much longer than I expected, so I draw a general outline.^a We want to show how to construct an isomorphism $f : \mathbb{R}_A \rightarrow \mathbb{R}_B$.

1. Realize that there are unique embeddings of \mathbb{N} in \mathbb{R}_A and \mathbb{R}_B that preserve the inductive principle, the order, closure of addition, and closure of multiplication, the additive identity, and the multiplicative identity. Call these ordered doubly-monoid (since it's a monoid w.r.t. $+$ and \times) homomorphisms ι_A, ι_B .
2. Construct an isomorphism $f_1 : \iota_A(\mathbb{N}) \rightarrow \iota_B(\mathbb{N})$ that preserves the inductive principle, order, addition, and multiplication. This is easy to do by just constructing $f_1 = \iota_B \circ \iota_A^{-1}$.
3. Extend f_1 to the ordered ring isomorphism f_2 by explicitly defining what it means to map additive inverses, i.e. negative numbers.
4. Extend f_2 to the ordered field isomorphism f_3 by explicitly defining what it means to map multiplicative inverses, i.e. reciprocals.
5. Extend f_3 to the ordered field isomorphism on the entire domain \mathbb{R}_A and codomain \mathbb{R}_B . There is no additional operations that we need to support, but we should explicitly show that this is both injective and surjective, which completes our proof.

^aFollowed from here.

It seems that the real numbers is *any* set that satisfies the definition above. Therefore, a line \mathbb{L} with $+$ associated with the translation of \mathbb{L} along itself and \cdot associated with the "stretching/compressing" of the line around the additive origin 0 is a valid representation of the reals. \mathbb{R} can also be represented as an uncountable list of numbers with possibly infinite decimal points, known as the decimal number system.

$$\dots, -2.583\dots, \dots, 0, \dots, 1.2343\dots, \dots, \sqrt{2}, \dots \quad (36)$$

The first property we should know is that the reals are uncountable.

Theorem 1.14 (Cantor's Diagonalization)

\mathbb{R} is uncountable.

Proof. We proceed by contradiction. Suppose the real numbers are countable. Then there exists a bijection $f : \mathbb{N} \rightarrow \mathbb{R}$. This means we can list all real numbers in $[0, 1]$ as an infinite sequence.^a

$$f(1) = 0.a_{11}a_{12}a_{13}\dots \quad (37)$$

$$f(2) = 0.a_{21}a_{22}a_{23}\dots \quad (38)$$

$$f(3) = 0.a_{31}a_{32}a_{33}\dots \quad (39)$$

$$\vdots \quad (40)$$

where each a_{ij} is a digit between 0 and 9.

Now construct a new real number $r = 0.r_1r_2r_3 \dots$ where:

$$r_n = \begin{cases} 1 & \text{if } a_{nn} \neq 1 \\ 2 & \text{if } a_{nn} = 1 \end{cases} \quad (41)$$

This number r is different from $f(n)$ for every $n \in \mathbb{N}$, since r differs from $f(n)$ in the n th decimal place. Therefore $r \in [0, 1]$ but $r \notin \text{range}(f)$, contradicting that f is surjective. Thus our assumption that the real numbers are countable must be false.

^aThis must be explicitly proven, but we can take the set of all Cauchy sequences of rationals in their decimal expansion and construct the reals this way.

With this, we can add the inner product, metric, and topology.

1.6 Exponentials, Roots, and Logarithms

Now we will focus on some other operations that become well-defined in the reals. We know that x^n for $n \in \mathbb{N}$ denotes repeated multiplication and x^{-1} denotes the multiplicative inverse. We need to build up on this notation. As a general outline, we will show that x^{-n} is well defined, then x^q , $q \in \mathbb{Q}$ is well-defined, and finally x^r , $r \in \mathbb{R}$ is well-defined. For the naturals, we have defined x^n as the repeated multiplication of n . It is trivial that the canonical injection $\iota_0 : \mathbb{N} \rightarrow \mathbb{R}$ commutes with the exponential map of naturals. We prove that $\iota_1 : \mathbb{Z} \rightarrow \mathbb{R}$ also commutes.

Lemma 1.3 (Integer Exponents)

We have

1. For $x_1, \dots, x_n \in \mathbb{R}$, $(x_1 \dots x_n)^{-1} = x_n^{-1} \dots x_1^{-1}$.
2. For $x \in \mathbb{R}$, $x > 0$, $(x^n)^{-1} = (x^{-1})^n$. This value is denoted x^{-n} .
3. For $x \in \mathbb{R}$ and $w, z \in \mathbb{Z}$, $x^{w+z} = x^w x^z$.
4. For $w, z \in \mathbb{Z}$, $x^{wz} = (x^z)^w = (x^w)^z$.

Proof. Listed.

1. The proof is trivial, but for $n = 2$ and $x_1 = x, x_2 = y$, we see that by associativity, $(x^{-1}x^{-1})(xy) = y^{-1}(x^{-1}x)y = y^{-1}y = 1$ and we know inverses are unique.
2. Set $x_i = x$ using (1).
3. If $w, z > 0$ this is trivial by the associative property. If either or both are negative, say $w < 0 < z$, then we set $w' = -w > 0$ and using (2) we know that

$$x^w x^z = (x^{-1})^{w'} x^z = x^{-w'+z} = x^{w+z} \quad (42)$$

by associativity in the second last equality.

Therefore, we have successfully defined x^z for all $z \in \mathbb{Z}$, and if z is negative, we're allowed to "swap" the -1 and $|z|$ in the exponents. Now we want to extend this into rational exponents, first by proving the existence and uniqueness of n th roots for any real. The proof is a little involved, but the general idea is that we want to use the LUB property to define the n th root as the supremum of a set.

Theorem 1.15 (Existence of Nth Roots)

For any real $x > 0$ and every $n \in \mathbb{N}$ there is one and only one positive real $y \in \mathbb{R}$ s.t. $y^n = x$. This is denoted $x^{1/n}$.

Proof. Let E be the set consisting of all reals $t \in \mathbb{R}$ s.t. $t^n < x$. We show that

1. it is nonempty. Consider $t = x/(1+x)$. Then $0 \leq t < 1 \implies t^n \leq t < x$. Thus $t \in E$ and E is nonempty.
2. it is bounded. Consider any number $s = 1+x$. Then $s^n \geq s > x$, so $s \notin E$, and $s = 1+x$ is an upper bound of E .

Therefore, E is a nonempty set that is upper bounded, so it has a least upper bound, called $y = \sup E$. We claim that $y^n = x$, proving by contradiction. For both cases, we use the fact that the identity $b^n - a^n = (b-a)(b^{n-1} + b^{n-2}a + \dots + a^{n-1})$ gives the inequality

$$b^n - a^n < (b-a)nb^{n-1} \text{ for } 0 < a < b \quad (43)$$

1. Assume $y^n < x$. Then we choose a fixed $0 < h < 1$ s.t.

$$h < \frac{x - y^n}{n(y+1)^{n-1}} \quad (44)$$

Then by putting $a = y, b = y+h$, we have

$$(y+h)^n - y^n < hn(y+h)^{n-1} < hn(y+1)^{n-1} < x - y^n \quad (45)$$

and thus $y^n < (y+h)^n < x$. This means that $y+h \in E$, and so y is not an upper bound.

2. Assume $y^n > x$. Then we set a fixed number

$$k = \frac{y^n - x}{ny^{n-1}} \quad (46)$$

Then $0 < k < y$. If we take any $t \in \mathbb{R}$ s.t. $t \geq y-k$, this implies that $t^n \geq (y-k)^n \implies -t^n \geq -(y-k)^n$, and so

$$y^n - t^n \leq y^n - (y-k)^n < kny^{n-1} = y^n - x \quad (47)$$

Thus $t^n > x$ and $t \notin E$. So it must be the case that $t < y-k$, and so $y-k$ is an upper bound of E , contradicting that y is least.

At this point, rooting has been introduced as sort of an independent map from exponentiation. We show that they have the nice property of commuting.

Lemma 1.4 (Rooting and Exponentiation Commute)

For $p \in \mathbb{Z}, q \in \mathbb{N}$ and $x \in \mathbb{R}$ with $x > 0$, we have

$$(x^p)^{1/q} = (x^{1/q})^p \quad (48)$$

Proof. If $p > 0$, then let $r = (x^p)^{1/q}$. By definition $r^q = x^p$. Let $s = x^{1/q}$. By definition $s^q = x$. Therefore $r^q = (s^q)^p = s^{qp}$ from the lemma on integer exponents. But since roots are well-defined and unique

$$r = (r^q)^{1/q} = (s^{qp})^{1/q} = s^p \implies (x^p)^{1/q} = (x^{1/q})^p \quad (49)$$

If $p = 0$, this is trivially 0, and if $p < 0$ the by the same logic with $p = -p'$ for $p' > 0$ and $y = x^{-1} > 0$. we know

$$(x^p)^{1/q} = ((y^{-1})^{-p'})^{1/q} = (y^{-(-p')})^{1/q} = (y^{p'})^{1/q} \quad (50)$$

$$= (y^{1/q})^{p'} = ((x^{-1})^{1/q})^{p'} = (x^{1/q})^{-p'} = (x^{1/q})^p \quad (51)$$

Theorem 1.16 (Rational Exponential Function)

Given $m, p \in \mathbb{Z}$ and $n, q \in \mathbb{N}$, prove that

$$(b^m)^{1/n} = (b^p)^{1/q} \quad (52)$$

Hence it makes sense to define $b^r = (b^m)^{1/n}$, since every element of the equivalence class r of each rational number maps to the same value.

Proof. Since $m/n = p/q \implies mq = np$,

$$b^{mq} = b^{np} \implies (b^m)^q = (b^p)^n \quad (53)$$

$$\implies b^m = ((b^m)^q)^{1/q} = ((b^p)^n)^{1/q} \quad (54)$$

$$\implies b^m = (b^p)^{1/q \cdot n} \quad (55)$$

$$\implies (b^m)^{1/n} = (b^p)^{1/q} \quad (56)$$

Therefore we can define for any $r \in \mathbb{Q}$

$$x^r = x^{m/n} = (x^m)^{1/n} = (x^{1/n})^m \quad (57)$$

where the final equality holds from the commutativity of rooting and exponentiation.

It turns out that this is a homomorphism.

Corollary 1.1 (Rational Exponential Function is a Homomorphism)

The rational exponential function is a homomorphism. That is, given $r, s \in \mathbb{Q}$ and $x \in \mathbb{R}$,

$$x^{r+s} = x^r \cdot x^s \quad (58)$$

Proof. Let $r = m/n, s = p/q$. Then

$$\begin{aligned} x^{r+s} &= x^{m/n+p/q} = x^{\frac{mq+np}{nq}} && \text{(addition on } \mathbb{Q}) \\ &= (x^{mq+np})^{1/nq} && \text{(exp and roots commute)} \\ &= (x^{mq} \cdot x^{np})^{1/nq} && \text{(int exp lemma)} \\ &= (x^{mq})^{1/nq} (x^{np})^{1/nq} && \text{(int exp lemma)} \\ &= x^{mq/nq} x^{np/nq} && \text{(exp and roots commute)} \\ &= x^{m/n} x^{p/q} && \text{(relation from } \mathbb{Q}) \end{aligned}$$

With rational exponents defined, we can use the least upper bound property to define a consistent extension of a real exponent.

Lemma 1.5 ()

If $r \in \mathbb{Q}$ with $r \geq 0$, then for $x \in \mathbb{R}, x > 1, 1 \leq b^r$.

Proof. Let $r = m/n$. Then $x^r = x^{m/n} = (x^m)^{1/n}$. Since $1 < x$, and $m \geq 0$, we have

$$1 \leq x \leq x^2 \leq \dots \leq x^m \implies 1 \leq b^m \quad (59)$$

Now set $y = x^{m/n}$ and assume that $y < 1$. Then

$$x^m = y^n < y^{n-1} < \dots < y < 1 \quad (60)$$

and so $x^m < 1$, which is a contradiction. So it must be the case that $y > 1$.

Lemma 1.6 (Monotonicity of Rational Exponents)

If $x, y \in \mathbb{R}$, then for any rational $r \in \mathbb{Q}$ with $r < x + y$, there exists a $p, q \in \mathbb{Q}$ s.t. $p < x, q < y$ and $p + q = r$. The converse is true as well.

Proof. $r < x + y \implies r - y < x$. By density of \mathbb{Q} in \mathbb{R} , we can choose $r - y < p < x$. Then $-r + y > -p > x \implies r - r + y > r - p > r - x \implies y > r - p > r - x$, and we set $q = r - p$. We are done. The converse is trivial since given $p, q \in \mathbb{Q}$ with $p < x, q < y$, then by the ordered field properties $p + q < x + y$.

Corollary 1.2 (Real Exponential Function)

Given $x \in \mathbb{R}$, we define

$$B(x) := \{x^q \in \mathbb{R} \mid q \in \mathbb{Q}, q \leq x\} \quad (61)$$

We claim that given $r \in \mathbb{R}$,

$$x^r := \sup B(r) \quad (62)$$

is well-defined and is a homomorphism extension of the rational exponential function. That is,

$$\sup B(x + y) = \sup B(x) \cdot \sup B(y) \quad (63)$$

Proof. To show that $x^r := \sup B(r)$ where $B(r) = \{x^t \in \mathbb{R} \mid t \in \mathbb{Q}, t \leq r\}$,

1. We show it's an upper bound. Assume it wasn't. Then $x^r < x^t$ for some $t \in \mathbb{Q}$ satisfying $t \leq r$. But $t \leq r \implies 0 \leq r - t$, and by the previous lemma, $1 \leq x^{r-t}$. So $1 \leq x^{r-t} = x^r x^{-t} = x^r (x^t)^{-1} \implies x^t \leq x^r$, which is a contradiction.
2. We show that it is least. Assume that it is not. Then $\exists r' \in \mathbb{Q}$ s.t. $x^t \leq x^{r'}$ and $r' < r$. Now let $s \in \mathbb{Q}$ be an element between r' and r , which is guaranteed to exist due to density of rationals in reals. But $s < r$, so by definition $x^s \in B(r)$, but

$$0 < s - r' \implies 1 < b^{s-r'} \quad (64)$$

$$\implies b^{r'} (b^{r'})^{-1} < b^s (b^{-r'}) \quad (65)$$

$$\implies 1 < b^s (b^{r'})^{-1} \quad (66)$$

$$\implies b^{r'} < b^s \quad (67)$$

and so $b^{r'}$ is not an upper bound for $B(r)$. By contradiction, b^r is least.

Since this is defined, the analogous definition for real numbers is consistent with that of the rationals, and it is upper bounded by the Archimedean principle, so such a supremum must exist. Note that t is rational. For the second part, from the previous lemma and the homomorphism properties of the rational exponent,

$$B(x + y) = B'(x + y) := \{b^{p+q} \in \mathbb{R} \mid p, q \in \mathbb{Q}, p \leq x, q \leq y\} \quad (68)$$

$$= \{b^p b^q \in \mathbb{R} \mid p, q \in \mathbb{Q}, p \leq x, q \leq y\} \quad (69)$$

$$(70)$$

Therefore we can treat B and B' as the same set.

1. Prove upper bound $\sup B(x+y) \leq \sup B(x) \sup B(y)$. Given $\alpha \in B'(x+y)$, there exists $p_\alpha, q_\alpha \in \mathbb{Q}$ (with $p_\alpha < x, q_\alpha < y$) s.t. $b^{p_\alpha} b^{q_\alpha} = \alpha$. But

$$b^{p_\alpha} b^{q_\alpha} \leq \sup_{p_\alpha} \{b^{p_\alpha}\} \cdot \sup_{q_\alpha} \{b^{q_\alpha}\} = \sup B(x) \sup B(y) \quad (71)$$

2. To prove least, assume there exists $K \in \mathbb{R}$ s.t. $\sup B'(x+y) \leq K < \sup B(x) \sup B(y)$. Then, since the image of b^x is always positive, we assume $0 < K$. We bound its factors as so: $K < \sup B(x) \sup B(y) \implies K/\sup B(x) < \sup B(y)$. By density of the rationals, there exists a $\beta \in \mathbb{Q}$, s.t.

$$\frac{K}{\sup B(x)} < \beta < \sup B(y) \quad (72)$$

This means $K/\beta < \sup B(x)$ and $\beta < \sup B(y)$. But this means that there exists $\phi, \gamma \in B(x), B(y)$ s.t. $K/\beta < \phi, \beta < \gamma \implies K = (K/\beta) \cdot \beta < \phi\gamma \implies \phi\gamma \in B'(x+y)$ by definition. So K is not an upper bound.

Furthermore, this is an isomorphism, and the inverse is defined. Let's define this analytically.

Theorem 1.17 (Logarithm)

For $b > 1$ and $y > 0$, there is a unique real number x s.t. $b^x = y$. We claim

$$x = \sup\{w \in \mathbb{R} \mid b^w < y\} \quad (73)$$

x is called the **logarithm of y to the base b** .

Proof. We use the inequality $b^n - 1 \leq n(b-1)$ for all $n \in \mathbb{N}$.^a By substituting $b = b^{1/n}$ (valid since $b > 1 \iff b^{1/n} > 1$) so $b-1 \geq n(b^{1/n} - 1)$. Now set some $t > 1$, and by Archimidean principle, we can choose some $n \in \mathbb{N}$ s.t. $n > \frac{b-1}{t-1}$. Then $n(t-1) > b-1$, and with the inequality derived we get

$$n(t-1) > b-1 \geq n(b^{1/n} - 1) \implies t > b^{1/n} \quad (74)$$

This allows us to prove 2 things.

1. If w satisfies $b^w < y$, then $b^{w+(1/n)} < y$ for sufficiently large n . Setting $t = yb^{-w}$ (which is greater than 1 since $b^w < y$) gives $y \cdot b^{-w} > b^{1/n} \implies b^w b^{1/n} < y \implies b^{w+(1/n)} < y$.
2. If w satisfies $b^w > y$, then $b^{w-(1/n)} > y$ for sufficiently large n . Setting $t = b^w y^{-1}$ (which is greater than 1 since $b^w > y$) gives $b^w y^{-1} > b^{1/n} \implies b^{w-(1/n)} > y$.

Now we can prove existence. Let A the set of all w s.t. $b^w < y$. We claim that $x = \sup A$.

1. Assume that $b^x < y$. We know that there exists $n \in \mathbb{N}$ s.t. $b^{x+(1/n)} < y \implies x + (1/n) \in A$, contradicting that x is an upper bound.
2. Assume that $b^x > y$. We know that there exists $n \in \mathbb{N}$ s.t. $b^{x-(1/n)} > y \implies x - (1/n)$ is also an upper bound for A , contradicting that x is least. Therefore $b^x = y$.

We now prove uniqueness. Assume that there are two such x 's, call them x, x' . By total ordering and $x \neq x'$, WLOG let $x > x' \implies x - x' > 0 \implies b^{x-x'} > 1$. By density of rationals, since we can choose $r \in \mathbb{R}$ s.t. $0 < r < x - x'$, we have $1 < b^r < b^{x-x'}$ and so $B(r) \subset B(x - x')$. Since $1 < b^{x-x'} \implies 1 \cdot b^{x'} < b^{x-x'} \cdot b^{x'} = b^x$, we have $b^{x'} < b^x$ and they cannot both be y . So $x = x'$.

^aWe prove by induction. For $n = 1$ $b^1 - 1 \leq 1(b-1)$. Assume that this holds for some n . Then $b^{n+1} - 1 = b^{n+1} - b + b - 1 = b(b^n - 1) + (b-1) \geq bn(b-1) + (b-1) = (bn+1)(b-1) \geq (n+1)(b-1)$, where the last step follows since $b \geq 1 \implies bn \geq n \implies bn+1 \geq n+1$.

1.7 Extended Reals

Often, we deal with numbers that are not finite, and we would like to have a system to incorporate $\pm\infty$ into the real line. Most first courses glaze over this, but it's important to see the construction as well. The problem is that we can't really add in these numbers without breaking a lot of the algebraic properties, but let's see for ourselves. It should be pretty obvious that we want (note the strict inequalities)

$$-\infty < x < +\infty \quad \forall x \in \mathbb{R} \quad (75)$$

To define addition, we can't make $x + \infty$ a finite number since then

$$\infty \leq x + \infty = y \quad (76)$$

which is a contradiction. So $x + \infty = +\infty$. We keep doing this but the main problem comes in with trying to define $\infty - \infty$ or ∞/∞ , which are known as **indeterminate forms**. These are particularly bad since we cannot deduce $x = y$ from $x + \infty = y + \infty$ or from $x \cdot \infty = y \cdot \infty$. The solution to this is to *simply avoid them*⁷ by making these indeterminate terms undefined.

Definition 1.15 (Extended Real Number Line)

The **extended real number line** is the set $\mathbb{R} \cup \{\pm\infty\}$ with the following operations.

1. *Order.* $-\infty \leq x$ and $x \leq +\infty$ for all $x \in \mathbb{R}$.
2. *Addition.*

$$\forall x \in \mathbb{R}, x + \infty = \infty + x = +\infty \quad (77)$$

$$\forall x \in \mathbb{R}, x - \infty = \infty - x = -\infty \quad (78)$$

$$+\infty + \infty = +\infty \quad (79)$$

$$-\infty - \infty = -\infty \quad (80)$$

$$+\infty - \infty, -\infty + \infty \text{ are undefined} \quad (81)$$

3. *Multiplication.*^a

$$\forall x \in \mathbb{R} \setminus \{0\}, x \cdot +\infty = +\infty \cdot x = \begin{cases} +\infty & \text{if } x > 0 \\ -\infty & \text{if } x < 0 \end{cases} \quad (82)$$

$$\forall x \in \mathbb{R} \setminus \{0\}, x \cdot -\infty = -\infty \cdot x = \begin{cases} -\infty & \text{if } x < 0 \\ +\infty & \text{if } x > 0 \end{cases} \quad (83)$$

$$0 \cdot +\infty = +\infty \cdot 0 = 0 \quad (84)$$

$$0 \cdot -\infty = -\infty \cdot 0 = 0 \quad (85)$$

$$+\infty \cdot +\infty = -\infty \cdot -\infty = +\infty \quad (86)$$

$$+\infty \cdot -\infty = -\infty \cdot +\infty = +\infty \quad (87)$$

$$(88)$$

^aThe fact that $0 \cdot \infty = 0$ might sound odd. Look at the extension into hyperreals later.

It turns out that this is still Dedekind-complete, which is nice. Unfortunately this is not even a field since the multiplicative inverse of $\pm\infty$ is not defined. Furthermore, we lose the Archimedean property.

The general rule of thumb is that if one wishes to use cancellation, this is only safe if one can guarantee that the numbers we work with are all finite. If we must work with infinity, another way is to work with the nonnegative reals.

⁷as far as I know

Definition 1.16 (Extended Real Number Line)

The **extended nonnegative reals** is the set $\mathbb{R}_{\geq 0} \cup \{+\infty\}$ with the following operations.

1. *Order.* $x \leq +\infty$ for all $x \in \mathbb{R}$.
2. *Addition.*

$$\forall x \in [0, +\infty], +\infty + x = x + \infty = +\infty \quad (89)$$

3. *Multiplication.*

$$\forall x \in (0, +\infty], +\infty \cdot x = x \cdot +\infty = +\infty \quad (90)$$

$$0 \cdot +\infty = +\infty \cdot 0 = 0 \quad (91)$$

There is a tradeoff here: we can work with infinity, or negative numbers, but not both. Also, note that if we define $\infty \cdot 0 = 0$, the multiplication becomes *upward continuous*, but not *downwards continuous*. This leads to an asymmetry when defining integrals, but in univariate analysis we will only work with bounded functions, and this will not hinder us until measure theory.

1.8 Hyperreals

The loss of the field property of the extended reals is quite bad, and we might want to recover this. Therefore, we can add more elements that serve to be the multiplicative inverse of infinity. We call these inverses *infinitesimals* and the new set the *hyperreal numbers*.

Theorem 1.18 (Hyperreals)

The **hyperreals**

In fact, when Newton first invented calculus, the hyperreals were what he worked with, and you can surprisingly build a good chunk of calculus with this. Even though this is a dead topic at this point, a lot of modern notation is based off of this number system, so it's good to see how it works. For example, when we write the integral

$$\int f(x) dx \quad (92)$$

we are saying that we are taking the uncountable sum of the terms $f(x) dx$, the multiplication of the real number $f(x)$ and the infinitesimal number dx living in the hyperreals. Unfortunately, we cannot fully construct a rigorous theory of calculus with only infinitesimals. However, in practice (especially physics) people tend to manipulate and do algebra with infinitesimals, so having a good foundation on what you can and cannot do with them is practical. While the focus won't be on *smooth infinitesimal analysis (SIA)*, I will include some alternate constructions later purely with infinitesimals.

1.9 Some Algebraic Inequalities

We also introduce various inequalities that may be useful for producing future results. The following lemmas can be proved with elementary algebra on the field of reals.

Lemma 1.7 (Bernoulli's Inequality)

For any real $x \geq -1$ and $n \in \mathbb{N}$, we have

$$(1 + x)^n \geq 1 + nx \quad (93)$$

Proof. We prove by induction. For $n = 1$, it is trivial. Now given that the inequality is satisfied for some $n \in \mathbb{N}$, we have

$$(1+x)^{n+1} = (1+x)^n(1+x) \quad (94)$$

$$\geq (1+nx)(1+x) \quad (95)$$

$$= 1 + nx + x + nx^2 \quad (96)$$

$$= 1 + (n+1)x \quad (97)$$

Lemma 1.8 (Young's Inequalities)

If $a > 0$ and $b > 0$, and the numbers p and q are such that $p \neq 0, 1$ and $q \neq 0, 1$, and $\frac{1}{p} + \frac{1}{q} = 1$, then

$$a^{\frac{1}{p}} b^{\frac{1}{q}} \leq \frac{1}{p}a + \frac{1}{q}b \text{ if } p > 1 \quad (98)$$

$$a^{\frac{1}{p}} b^{\frac{1}{q}} \geq \frac{1}{p}a + \frac{1}{q}b \text{ if } p < 1 \quad (99)$$

and equality holds in both cases if and only if $a = b$.

Proof.

Lemma 1.9 (Holder's Inequalities)

Let $x_i \geq 0, y_i \geq 0$ for $i = 1, 2, \dots, n$, and let $\frac{1}{p} + \frac{1}{q} = 1$. Then,

$$\sum_{i=1}^n x_i y_i \leq \left(\sum_{i=1}^n x_i^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n y_i^q \right)^{\frac{1}{q}} \text{ for } p > 1 \quad (100)$$

$$\sum_{i=1}^n x_i y_i \geq \left(\sum_{i=1}^n x_i^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n y_i^q \right)^{\frac{1}{q}} \text{ for } p < 1, p \neq 0 \quad (101)$$

Proof.

Lemma 1.10 (Minkowski's Inequalities)

Let $x_i \geq 0, y_i \geq 0$ for $i = 1, 2, \dots, n$. Then,

$$\left(\sum_{i=1}^n (x_i + y_i)^p \right)^{\frac{1}{p}} \leq \left(\sum_{i=1}^n x_i^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n y_i^p \right)^{\frac{1}{p}} \text{ when } p > 1 \quad (102)$$

$$\left(\sum_{i=1}^n (x_i + y_i)^p \right)^{\frac{1}{p}} \geq \left(\sum_{i=1}^n x_i^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n y_i^p \right)^{\frac{1}{p}} \text{ when } p < 1, p \neq 0 \quad (103)$$

Proof.

1.10 Exercises

Exercise 1.1 (Math 531 Spring 2025, PS2.1)

Prove that the set of all matrices of the form:

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix}, \quad (104)$$

with $a, b \in \mathbb{R}$ forms a field with the usual sum and product operations of matrices. What does this field resemble? Give extensions to 3×3 and 4×4 matrices.

Solution.

Exercise 1.2 (Math 531 Spring 2025, PS2.2)

Why can't the field of complex numbers (with its usual operations) be made into an ordered field?

Solution. Solution is shown as theorem.

Exercise 1.3 (Math 531 Spring 2025, PS2.3)

Prove there are no finite ordered fields.

Solution. Solution is shown as theorem.

Exercise 1.4 (Math 531 Spring 2025, PS2.4)

Prove that if x and n are natural numbers, then

$$x^n - 1 = (x - 1)(1 + x + x^2 + \dots + x^{n-1}). \quad (105)$$

Solution. We use the commutative addition and multiplication, plus distributive property in \mathbb{Z} .

$$(x - 1) \left(\sum_{i=0}^{n-1} x^i \right) = x \sum_{i=0}^{n-1} x^i - \sum_{i=0}^{n-1} x_i \quad (106)$$

$$= \sum_{i=1}^n x^i - \sum_{i=0}^{n-1} x^i \quad (107)$$

$$= x^n + \sum_{i=1}^{n-1} x^i - \sum_{i=1}^{n-1} x^i - 1 \quad (108)$$

$$= x^n - 1 \quad (109)$$

Exercise 1.5 (Math 531 Spring 2025, PS2.7)

Prove that there is no $q \in \mathbb{Q}$ for which

$$q^2 + q = 4. \quad (110)$$

Solution. Assume that such a $q \in \mathbb{Q}$ in canonical form exists. Then by the field properties, since $\frac{1}{4} \in \mathbb{Q}$,

$$q^2 + q + \frac{1}{4} = 4 + \frac{1}{4} = \frac{17}{4} \in \mathbb{Q} \quad (111)$$

But by distributive properties, $(q + \frac{1}{2})^2 \in \mathbb{Q}$. We claim that there exists no rational $x = a/b$ (a, b coprime) s.t. $x^2 = 17/4$. If there were, then clearly $a \neq 0$ and

$$\frac{a^2}{b^2} = \frac{17}{4} \implies 4a^2 = 17b^2 \quad (112)$$

$$\implies 2|b, \text{ and so } b = 2b' \text{ for some } b' \in \mathbb{N} \quad (113)$$

$$\implies a^2 = 17(b')^2 \quad (114)$$

$$\implies 17|a \text{ and so } a = 17a' \text{ for some } a' \in \mathbb{Z} \quad (115)$$

$$\implies 17(a')^2 = (b')^2 \quad (116)$$

which implies that $17|b'$, but this contradicts the assumption that a, b are coprime. Therefore $q + \frac{1}{2} \notin \mathbb{Q} \implies q \notin \mathbb{Q}$.

Exercise 1.6 (Math 531 Spring 2025, PS2.8)

Let X be an ordered set with the least upper bound property. Prove that X has the greatest lower bound property.

Solution. Shown in theorem above.

Exercise 1.7 (Math 531 Spring 2025, PS2.9)

Prove that if $x, y \in \mathbb{Q}$ we have that

$$||x| - |y|| \leq |x - y|. \quad (117)$$

Solution. By subadditivity of the norm we have

$$|x| \leq |x - y| + |y| \implies |x| - |y| \leq |x - y| \quad (118)$$

$$|y| \leq |y - x| + |x| \implies |y| - |x| \leq |y - x| \quad (119)$$

But $|y - x| = |-1(x - y)| = |-1| \cdot |x - y| = |x - y|$, and so

$$\max\{|x| - |y|, |y| - |x|\} \leq |x - y| \quad (120)$$

and the LHS is the definition of the norm $||x| - |y||$ in \mathbb{Q} .

Exercise 1.8 (Rudin 1.1)

If r is rational ($r \neq 0$) and x is irrational, prove that $r + x$ and rx are irrational.

Solution. If we assume that $rx = t$ and $r + x = s$ are rational, then this violates the field axioms of \mathbb{Q} since then $x = tr^{-1}$ and $x = s + (-r)$ are rational.

Exercise 1.9 (Rudin 1.2)

Prove that there is no rational number whose square is 12.

Solution. Assume that there exists a number p/q such that p and q are both not even. Then,

$$\left(\frac{p}{q}\right)^2 = 12 \implies p^2 = 12q^2 = 3(2q)^2 \quad (121)$$

So p must be even $p = 2p'$. Therefore, $p'^2 = 3q^2$, and q must be odd. This means that p' must be odd. We can rewrite the equation

$$p'^2 - q^2 = 2q^2 \implies (p' + q)(p' - q) = 2q^2 \quad (122)$$

where the left hand side is divisible by 4 but the right hand side is divisible by at most 2, leading to a contradiction.

Exercise 1.10 (Rudin 1.3)

Prove that the axioms of multiplication imply the following.

1. If $x \neq 0$ and $xy = xz$, then $y = z$.
2. If $x \neq 0$ and $xy = x$, then $y = 1$.
3. If $x \neq 0$ and $xy = 1$, then $y = x^{-1}$.
4. If $x \neq 0$, then $(x^{-1})^{-1} = x$.

Solution. Listed.

1. $xy = xz \implies \frac{1}{x} \cdot xy = \frac{1}{x}xz \implies y = z$
2. $xy = x \implies \frac{1}{x}xy = \frac{1}{x}x \implies y = 1$
3. $xy = 1 \implies \frac{1}{x}xy = \frac{1}{x}1 \implies y = \frac{1}{x}$
4. $(x^{-1})^{-1} \cdot x^{-1} = 1 \implies (x^{-1})^{-1} \cdot x^{-1} \cdot x = 1 \cdot x \implies (x^{-1})^{-1} = x$

Exercise 1.11 (Rudin 1.4)

Let E be a nonempty subset of an ordered set; suppose α is a lower bound of E and β is an upper bound of E . Prove that $\alpha \leq \beta$.

Solution. Since E is nonempty, we choose any $x \in E$. By definition, $\alpha \leq x$ and $x \leq \beta$, and by transitive property of orderings, we have $\alpha \leq \beta$.

Exercise 1.12 (Rudin 1.5)

Let A be a nonempty set of real numbers which is bounded below. Let $-A$ be the set of all numbers $-x$, where $x \in A$. Prove that

$$\inf A = -\sup(-A) \quad (123)$$

Solution. We would like to prove that $\inf A \leq -\sup(-A)$ and $\inf A \geq -\sup(-A)$. For the first part, we start off with the definition of the infimum.

$$\begin{aligned}\inf A \leq x \quad \forall x \in A &\implies -\inf A \geq -x \quad \forall x \in A \\ &\implies -\inf A \geq x \quad \forall x \in -A \\ &\implies -\inf A \geq \sup(-A) \\ &\implies \inf A \leq -\sup(-A)\end{aligned}$$

For the second part, we start with the definition of the supremum.

$$\begin{aligned}\sup(-A) \geq x \quad \forall x \in -A &\implies \sup(-A) \geq -x \quad \forall x \in A \\ &\implies -\sup(-A) \leq x \quad \forall x \in A \\ &\implies -\sup(-A) \leq \inf A\end{aligned}$$

Exercise 1.13 (Rudin 1.6)

Fix $b > 1$.

1. If m, n, p, q are integers, $n > 0, q > 0$, and $r = m/n = p/q$, prove that

$$(b^m)^{1/n} = (b^p)^{1/q}. \quad (124)$$

Hence it makes sense to define $b^r = (b^m)^{1/n}$.

2. Prove that $b^{r+s} = b^r b^s$ if r and s are rational.
3. If x is real, define $B(x)$ to be the set of all numbers b^t , where t is rational and $t \leq x$. Prove that

$$b^r = \sup B(r) \quad (125)$$

when r is rational. Hence it makes sense to define

$$b^x = \sup B(x) \quad (126)$$

for every real x .

4. Prove that $b^{x+y} = b^x b^y$ for all real x and y .

Solution. Proved in theorem above.

Exercise 1.14 (Rudin 1.7)

Fix $b > 1, y > 0$, and prove that there is a unique real x such that $b^x = y$, by completing the following outline. (This x is called the logarithm of y to the base b .)

1. For any positive integer n , $b^n - 1 \geq n(b - 1)$.
2. Hence $b - 1 \geq n(b^{1/n} - 1)$.
3. If $t > 1$ and $n > (b - 1)/(t - 1)$, then $b^{1/n} < t$.
4. If w is such that $b^w > y$, then $b^{-(1/n)} < y$ for sufficiently large n ; to see this, apply part (c) with $t = y \cdot b^{-w}$.
5. If $b^w > y$, then $b^{w-(1/n)} > y$ for sufficiently large n .
6. Let A be the set of all w such that $b^w < y$, and show that $x = \sup A$ satisfies $b^x = y$.
7. Prove that this x is unique.

Solution. Proved in theorem above.

Exercise 1.15 (Rudin 1.8)

Prove that no order can be defined in the complex field that turns it into an ordered field.

Solution. Note that if $x \geq 0$, then $-x \leq 0$ for all x of any ordered field. Since if $x \geq 0$ and $-x > 0$, then $x - x > 0$, which is absurd. Therefore, one of either i or $-i$ should be greater than 0. But $i^2 = (-i)^2 = -1$, so this means that $-1 > 0$, which implies that $0 < 1$. But either 1 or -1 must ≥ 0 .

Exercise 1.16 (Rudin 1.9)

Equip \mathbb{C} with the dictionary order. That is, given $z = a + bi$ and $w = c + di$, $z < w$ if $a < c$, or if $a = c$ and $b < d$. Does this ordered set have a least upper bound property?

Solution. No it does not. Consider the set $S = \{a + bi \in \mathbb{C} \mid a \leq 3\}$. S is bounded by 4, but it doesn't have a least upper bound. Given any $3 + bi$, this is not an upper bound since we can construct $3 + (b + \epsilon)i \in S$. Given any $a + bi$ where $a > 3$, we can always find a lower bound of form $a + (b - \epsilon)i$ that also bounds S .

Exercise 1.17 (Rudin 1.10)

Suppose $z = a + bi$, $w = u + iv$, and

$$a = \left(\frac{|w| + u}{2} \right)^{1/2} \quad \text{and} \quad b = \left(\frac{|w| - u}{2} \right)^{1/2} \quad (127)$$

Prove that $z^2 = w$ if $v \geq 0$ and that $(\bar{z})^2 = w$ if $v \leq 0$. Conclude that every complex number (with one exception!) has two complex square roots.

Solution. We can calculate

$$z^2 = (a^2 - b^2) + 2abi = u + \sqrt{v^2}i = \begin{cases} u + vi & \text{if } v \geq 0 \\ u - vi & \text{if } v \leq 0 \end{cases} \quad (128)$$

Since if we assume $v \geq 0$, then we have $z^2 = w$. We also get

$$\bar{z}^2 = (a^2 - b^2) - 2abi = u - \sqrt{v^2}i = \begin{cases} u - vi & \text{if } v \geq 0 \\ u + vi & \text{if } v \leq 0 \end{cases} \quad (129)$$

and assuming $v \leq 0$, we have $\bar{z}^2 = w$. Therefore, every complex number w has both $\pm z$ as its square root if $v \geq 0$, $\pm \bar{z}$ if $v \leq 0$, and just one root if $z = 0$.

Exercise 1.18 (Rudin 1.11)

If z is a complex number, prove that there exists an $r \geq 0$ and a complex number w with $|w| = 1$ s.t. $z = rw$. Are w and r always uniquely determined by z ?

Solution. If $z = 0$, then $r = 0$ and there is no unique w . If $z = a + bi \neq 0$, then define

$$r = |z| = (a^2 + b^2)^{1/2}, \quad w = \frac{1}{r}z \quad (130)$$

which proves existence. As for uniqueness, assume that there are two forms

$$z = rw = r'w' \quad (131)$$

Then, $w = \frac{r'}{r}w' \implies |w| = \left|\frac{r'}{r}\right||w'| = 1$, which implies that $r'/r = 1$ and so $r = r'$. This means that $w = w'$.

Exercise 1.19 (Rudin 1.12)

If z_1, \dots, z_n are complex, prove that

$$|z_1 + z_2 + \dots + z_n| \leq |z_1| + \dots + |z_n| \quad (132)$$

Solution. By induction, it suffices to prove $|z_1 + z_2| \leq |z_1| + |z_2|$. We have

$$\begin{aligned} |z_1 + z_2|^2 &= (z_1 + z_2)(\overline{z_1 + z_2}) \\ &= (z_1 + z_2)(\bar{z}_1 + \bar{z}_2) \\ &= z_1\bar{z}_1 + z_1\bar{z}_2 + z_2\bar{z}_1 + z_2\bar{z}_2 \\ &= |z_1|^2 + |z_2|^2 + z_1\bar{z}_2 + z_2\bar{z}_1 \\ &= |z_1|^2 + |z_2|^2 + 2(ac + bd) \\ &\leq |z_1|^2 + |z_2|^2 + 2\sqrt{a^2 + b^2}\sqrt{c^2 + d^2} \quad (\text{Schwartz}) \\ &= |z_1|^2 + |z_2|^2 + 2|z_1||z_2| \\ &= (|z_1| + |z_2|)^2 \end{aligned}$$

since both sides are positive, we can take their square root to get the desired result.

Exercise 1.20 (Rudin 1.13)

If x, y are complex, prove that

$$||x| - |y|| \leq |x - y| \quad (133)$$

Solution. Since both sides are nonnegative, we can square both sides. Note that due to Cauchy Schwartz inequality, $2|x||y| \geq x\bar{y} + y\bar{x}$ since expanding them gives

$$2\sqrt{a^2 + b^2}\sqrt{c^2 + d^2} \geq 2(ac + bd) \quad (134)$$

Therefore, the following inequality is true:

$$|x|^2 + |y|^2 - 2|x||y| \leq x\bar{x} + y\bar{y} - x\bar{y} - y\bar{x} \quad (135)$$

which reduces to form $(|x| - |y|)^2 \leq |x - y|^2$.

Exercise 1.21 (Rudin 1.14)

If z is a complex number s.t. $|z| = 1$, that is such that $z\bar{z} = 1$, compute

$$|1 + z|^2 + |1 - z|^2 \quad (136)$$

Solution. Compute.

$$(1+z)(1+\bar{z}) + (1-z)(1-\bar{z}) = 1+z+\bar{z}+z\bar{z} + 1-z-\bar{z}+z\bar{z} = 4 \quad (137)$$

Exercise 1.22 (Rudin 1.15)

Under what conditions does equality hold in the Schwarz inequality?

Solution. If they are antiparallel, since

$$\langle x, y \rangle = \|x\| \|y\| \cos \theta \quad (138)$$

Exercise 1.23 (Rudin 1.16)

Suppose $k \geq 3$, $\mathbf{x}, \mathbf{y} \in \mathbb{R}^k$, $|\mathbf{x} - \mathbf{y}| = d > 0$, and $r > 0$. Prove:

a) If $2r > d$, there are infinitely many $\mathbf{z} \in \mathbb{R}^k$ s.t.

$$|\mathbf{z} - \mathbf{x}| = |\mathbf{z} - \mathbf{y}| = r \quad (139)$$

b) If $2r = d$, there is exactly one such \mathbf{z} .

c) If $2r < d$, there is no such \mathbf{z} .

Solution.

Exercise 1.24 (Rudin 1.17)

Prove that

$$|\mathbf{x} + \mathbf{y}|^2 + |\mathbf{x} - \mathbf{y}|^2 = 2|\mathbf{x}|^2 + 2|\mathbf{y}|^2 \quad (140)$$

Solution. This is trivial if we simply expand

$$|\mathbf{x} + \mathbf{y}|^2 + |\mathbf{x} - \mathbf{y}|^2 = \langle \mathbf{x} + \mathbf{y}, \mathbf{x} + \mathbf{y} \rangle + \langle \mathbf{x} - \mathbf{y}, \mathbf{x} - \mathbf{y} \rangle \quad (141)$$

$$= \langle \mathbf{x}, \mathbf{x} \rangle + 2\langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{y}, \mathbf{y} \rangle + \langle \mathbf{x}, \mathbf{x} \rangle - 2\langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{y}, \mathbf{y} \rangle \quad (142)$$

$$= 2\langle \mathbf{x}, \mathbf{x} \rangle + 2\langle \mathbf{y}, \mathbf{y} \rangle \quad (143)$$

$$= 2|\mathbf{x}|^2 + 2|\mathbf{y}|^2 \quad (144)$$

Exercise 1.25 (Rudin 1.18)

If $k \geq 2$ and $\mathbf{x} \in \mathbb{R}^k$, prove that there exists $\mathbf{y} \in \mathbb{R}^k$ s.t. $\mathbf{y} \neq \mathbf{0}$, but $\mathbf{x} \cdot \mathbf{y} = 0$. Is this also true if $k = 1$?

Solution. Let $x \in \mathbb{R}^k$ and $\ell \in \mathbb{R}^{k*}$, the dual space. By Riesz representation theorem, we can define the canonical isomorphism $\ell \mapsto y$ between these two spaces as

$$\ell(x) = (x, y) \quad (145)$$

Since $y \neq 0$ by assumption, $\ell \neq 0$, and so its rank is at least 1. Since ℓ maps to \mathbb{R} , the rank has to be 1. By rank nullity theorem, we have

$$\dim N(\ell) = k - \text{rank}(\ell) = k - 1 \quad (146)$$

and so there exists nontrivial annihilators ℓ of x , which can be mapped to a nontrivial $y \in \mathbb{R}^k$.

Exercise 1.26 (Rudin 1.19)

Suppose $\mathbf{a}, \mathbf{b} \in \mathbb{R}^k$. Find $\mathbf{c} \in \mathbb{R}^k$ and $r > 0$ s.t.

$$|\mathbf{x} - \mathbf{a}| = 2|\mathbf{x} - \mathbf{b}| \quad (147)$$

if and only if $|\mathbf{x} - \mathbf{c}| = r$.

Solution. If we draw out the circle, it must contain two points on the line drawn by connecting A and B . Since it must be symmetric, its center and radius can then be easily calculated to be

$$r = \frac{2}{3}|b - a|, \quad c = \frac{1}{3}(4b - a) \quad (148)$$

Exercise 1.27 (Zorich 2.2.1)

Using the principle of induction, show that

1. the sum $x_1 + \dots + x_n$ of real numbers is defined independently of the insertion of parentheses to specify the order of addition.
2. the same is true of the product $x_1 \dots x_n$
3. $|x_1 + \dots + x_n| \leq |x_1| + \dots + |x_n|$
4. $|x_1 \dots x_n| \leq |x_1| \dots |x_n|$
5. For any $n, m \in \mathbb{N}$ such that $m < n$, $(n - m) \in \mathbb{N}$.
6. $(1 + x)^n \geq 1 + nx$ for $x > -1$ and $n \in \mathbb{N}$, equality holding for when $n = 1$ or $x = 0$.
7. $(a + b)^n = a^n + {}_nC_1 a^{n-1} b^1 + \dots + b^n$ (aka binomial theorem).

Solution. Listed.

1. Let n denote the number of elements in the sum. We prove by strong law of induction. The base case for when $n = 1, 2, 3$ is trivially true.

$$x_1 = x_1 \quad (\text{identity})$$

$$x_1 + x_2 = x_1 + x_2 \quad (\text{identity})$$

$$(x_1 + x_2) + x_3 = x_1 + (x_2 + x_3) \quad (\text{associativity})$$

Then, the sum of $n = k$ parameters is defined by $k - 2$ pairs of parentheses defining the order of the sum. These parentheses define a sequence of $k - 1$ 2-fold additions. Now, assume that the claim is true for

$$S_n \equiv x_1 + \dots x_n \text{ for } n = 1, 2, \dots, k \quad (149)$$

Then, for a specific sum S_{k+1} of $k + 1$ elements with $k - 1$ parentheses, we can reduce the sum to its final 2-fold addition

$$S_{k+1} \equiv \underbrace{(x_1 + \dots + x_i)}_{\varphi_1} + \underbrace{(x_{i+1} + \dots + x_{k+1})}_{\varphi_2} \quad (150)$$

Since $i, k - i + 1 < k$, by the strong law φ_1 and φ_2 are independent of the order of sum.

2. Exactly identical to (a).
3. By the triangle inequality $|x_1 + x + 2| \leq |x_1| + |x_2|$. Now, assume for $n = k$ is true. Then, let

$S_k = x_1 + \dots + x_k$, so

$$|x_1 + \dots + x_k + x_{k+1}| = |S_k + x_{k+1}| \leq |S_k| + |x_{k+1}| \leq \sum_{i=1}^{k+1} |x_i| \quad (151)$$

4. Same as (c).

5. Let us fix m to be any element of \mathbb{N} . Then, the base case is for $n = m + 1$ (which is in \mathbb{N} since it is inductive), so

$$n - m = (m + 1) - m = 1 \in \mathbb{N} \quad (152)$$

Now, given that for some integer $n \geq m + 1$, $n - m \in \mathbb{N}$ is true, we have

$$\begin{aligned} (n + 1) - m &= n + (1 - m) && \text{(associativity)} \\ &= n + (-m + 1) && \text{(commutativity)} \\ &= (n - m) + 1 && \text{(associativity)} \end{aligned}$$

where $(n - m) + 1 \in \mathbb{N}$ by inductive property of \mathbb{N} .

6. We prove by induction. For $n = 1$, it is trivial that $(1 + x)^1 \geq 1 + 1 \cdot x$. Now assume that the claim is true for some $k \in \mathbb{N}$. Then,

$$\begin{aligned} (1 + x)^{k+1} &= (1 + x)^k(1 + x) \geq (1 + kx)(1 + x) \\ &= 1 + (k + 1)x + kx^2 \\ &\geq 1 + (k + 1)x \end{aligned}$$

where equality holds if $x = 0 \implies 1^{k+1} = 1^k \cdot 1 = 1$ or $n = 1 \implies$ trivial case.

7. The base case for $n = 1$ is trivial since $(a + b)^1 = \binom{1}{0}a + \binom{1}{1}b$. We introduce Newton's identity.

$$\begin{aligned} \binom{k}{j-1} + \binom{k}{j} &= \frac{k!}{(j-1)!(k-j+1)!} + \frac{k!}{j!(k-j)!} \\ &= k! \left(\frac{j}{j!(k-j+1)!} + \frac{k-j+1}{j!(k-j+1)!} \right) \\ &= k! \cdot \frac{k+1}{j!(k-j+1)!} \\ &= \frac{(k+1)!}{j!(k-j+1)!} = \binom{k+1}{j} \end{aligned}$$

Now assuming that the binomial formula holds for some $n = k$, we have

$$(a + b)^{k+1} = (a + b)^k(a + b) \quad (153)$$

$$= \left(\sum_{j=0}^k \binom{k}{j} a^j b^{k-j} \right) (a + b) \quad (154)$$

$$= \sum_{j=0}^k \binom{k}{j} a^{j+1} b^{k-j} + \sum_{j=0}^k \binom{k}{j} a^j b^{k-j+1} \quad (155)$$

$$= \binom{k}{0} a^0 b^{k+1} + \binom{k}{k} a^{k+1} b^0 + \sum_{j=1}^{k-1} \binom{k}{j} a^{j+1} b^{k-j} + \sum_{j=1}^k \binom{k}{j} a^j b^{k-j+1} \quad (156)$$

$$= \binom{k+1}{0} a^0 b^{k+1} + \binom{k+1}{k+1} a^{k+1} b^0 + \sum_{j=1}^k \left[\binom{k}{j-1} + \binom{k}{j} \right] a^j b^{k-j+1} \quad (157)$$

$$= \sum_{j=0}^{k+1} \binom{k+1}{j} a^j b^{k-j+1} \quad (158)$$

Exercise 1.28 (Zorich 2.2.3)

Show that an inductive set is not bounded above.

Solution. Assume that a X is a nonempty inductive set that is bounded above. By definition, there exists a number $B \in \mathbb{R}$ such that $\max X < B$. Then, this means that there exists no numbers in $[B, B + 1)$. Since X is inductive, this means that there cannot exist any elements of X in the interval $[B - 1, B)$, and similarly for the interval $[B - 2, B)$, and so on, meaning that if $x \in X$, then $x \notin [B - k, B - k + 1)$ for all $k \in \mathbb{Z}$. By the Archimidean principle, this implies that $X = \emptyset$, contradicting our assumption.

Exercise 1.29 (Zorich 2.2.4)

Prove the following.

1. An inductive set is infinite (that is, equipollent with one of its subsets different from itself).
2. The set $E_n = \{x \in \mathbb{N} \mid x \leq n\}$ is finite.

Solution. Listed.

1. Assume that an inductive set X is finite $\implies X$ is bounded above (we can choose upper bound $B = \max X + 1$). But from 2.2.3, an inductive set cannot be bounded above, contradicting our assumption.
2. It is trivial that $E_1 = \{1\}$ is finite since $\text{card} E_1 = 1$. Now, if for some k , E_k is finite with cardinality e_k , then $\text{card} E_{k+1} = e_k + 1$, which implies finiteness.

Exercise 1.30 (Zorich 2.2.5)

Listed.

1. Let $m, n \in \mathbb{N}$ and $m > n$. Their greatest common divisor $\gcd(m, n) = d \in \mathbb{N}$ can be found in a finite number of steps using the following algorithm of Euclid involving successive divisions with remainder.

$$m = q_1 n + r_1$$

$$n = q_2 r_1 + r_2$$

$$r_1 = q_3 r_2 + r_3$$

$$\dots = \dots$$

$$r_{k-2} = q_k r_{k-1} + r_k$$

$$r_{k-1} = q_{k+1} r_k + 0$$

Then $d = r_k$

2. If $d = \gcd(m, n)$, one can choose numbers $p, q \in \mathbb{Z}$ such that $pm + qn = d$.

Solution. Listed.

- 1.
2. Letting $n = r_0$, notice that the equations above satisfy for $i = 0, 1, \dots$

$$r_i = q_{i+2} r_{i+1} + r_{i+2} \implies r_i - q_{i+2} r_{i+1} = r_{i+2} \quad (1)$$

Note that the second-to-last equation allows us to write r_k as a linear combination of r_{k-2} and r_{k-1} : $r_k = r_{k-2} - q_k r_{k-1}$. Now by applying (1), we can reduce the above to a linear combination

of r_{k-3} and r_{k-2} .

$$\begin{aligned} r_k &= r_{k-2} - q_k r_{k-1} \\ &= r_{k-2} - q_k(r_{k-3} - q_{k-1} r_{k-3}) \\ &= (1 + q_{k-1} q_k) r_{k-2} - q_k r_{k-3} \end{aligned}$$

and repeatedly doing this allows us to reduce r_k to a linear combination $q_0 r_0 + q_1 r_1$. By the ring properties of \mathbb{Z} , the new linear coefficients are also in \mathbb{Z} . Reducing one last time using the first equation in the Euclidean algorithm gives

$$\begin{aligned} r_k &= q_0 r_0 + q_1 r_1 \\ &= q_0 n + q_1(m - q_1 n) \\ &= q_1 m + (q_0 - q_1) n \\ &= pm + qn \end{aligned}$$

Exercise 1.31 (Zorich 2.2.9)

Show that if the natural number n is not of the form k^m , where $k, m \in \mathbb{N}$, then the equation $x^m = n$ has no rational roots.

Solution. Assume that there is a rational solution $x = p/q$, with $p, q \in \mathbb{N}$ of the equation. Then,

$$\left(\frac{p}{q}\right)^m = \frac{p^m}{q^m} = n \implies p^m = q^m n \quad (159)$$

By the fundamental theorem of arithmetic, the exponents of the prime factors of p^m must all be multiples of m , and so it must be so for the right hand side $\implies x$ must be of form $x = k^m$ for some k . This is a contradiction.

Exercise 1.32 (Zorich 2.2.12)

Knowing that $\frac{m}{n} \equiv m \cdot n^{-1}$ by definition, where $m \in \mathbb{Z}$ and $n \in \mathbb{N}$, derive the “rules” for addition, multiplication, and division of fractions, and also the condition for two fractions to be equal.

Solution. We can construct a \mathbb{Q} as a quotient space $\mathbb{Z} \times \mathbb{N} / \sim$, where \sim is an equivalent relation where

$$(q_1, p_1) \sim (q_2, p_2) \text{ iff } q_1 p_2 = p_1 q_2 \quad (160)$$

which is the familiar equivalence relation from “simplifying” a fraction. We define addition and multiplication as the following

$$\begin{aligned} (a, b) + (c, d) &= (ad + bc, bd) \\ (a, b) \cdot (c, d) &= (ac, bd) \end{aligned}$$

which turns out to be algebraically closed in \mathbb{Q} . The additive identity is the equivalence class $0 = \{(0, c) \mid c \in \mathbb{N}\}$, and the multiplicative identity is the equivalence class $1 = \{(c, c) \mid c \in \mathbb{N}\}$. It is easy to check that $+$ is commutative, the additive inverse is $-(a, b) = (-a, b)$, and the multiplicative inverse is $(a, b)^{-1} = (b, a)$. We can subtract and divide these elements of \mathbb{Q} , called “fractions,” as such:

$$\begin{aligned} (a, b) - (c, d) &= (a, b) + (-(c, d)) = (a, b) + (-c, d) = (ad - bc, bd) \\ (a, b) \div (c, d) &= (a, b) \cdot (c, d)^{-1} = (a, b) \cdot (d, c) = (ad, bc) \end{aligned}$$

Exercise 1.33 (Zorich 2.2.13)

Verify that the rational numbers \mathbb{Q} satisfy all the axioms for real numbers except for the axiom of completeness.

Solution. From continuing the steps of 2.2.14, we can prove \mathbb{Q} is an algebraic field (associativity, commutativity of addition and multiplication, along with distributive property). We can actually define the order relation $\leq_{\mathbb{Q}}$ in two ways:

1. $(a, b) \leq (c, d)$ iff $ad \leq_{\mathbb{Z}} bc$, where $\leq_{\mathbb{Z}}$ is the order relation on \mathbb{Z} (which can be defined much more simply).
2. Recognizing that $\mathbb{Q} \subset \mathbb{R}$, we define the canonical injection map $i : \mathbb{Q} \rightarrow \mathbb{R}$ and by abuse of language, endow the relation $\leq_{\mathbb{Q}}$ as the restriction of $\leq_{\mathbb{R}}$ onto \mathbb{Q} . That is, for $(a, b), (c, d) \in \mathbb{Q}$,

$$(a, b) \leq_{\mathbb{Q}} (c, d) \text{ iff } i(a, b) \leq_{\mathbb{R}} i(c, d) \quad (161)$$

The ordering for the 1st step can be checked for consistency.

1. $(a, b) \leq (a, b)$ since $ab \leq ab$ (true in \mathbb{Z})
2. $(a, b) \leq (c, d), (c, d) \leq (a, b)$ means that $ad \leq bc$ and $bc \leq ad \implies ad = bc$ (true in \mathbb{Z})
3. $(a, b) \leq (c, d) \leq (e, f)$ implies $ad \leq bc, cf \leq de$. Multiplying positive (important that $f > 0$!) to the first inequality gives $adf \leq bcf$, and multiplying positive b to the second gives $bcf \leq bde$, and by interpreting \leq as the ordering defined on \mathbb{Z} , we use transitive property of $\leq_{\mathbb{Z}}$ to get $adf \leq bde \implies af \leq be \iff (a, b) \leq (e, f)$.
4. For any $(a, b), (c, d) \in \mathbb{Q}$, $(a, b) \leq (c, d)$ or $(a, b) \geq (c, d)$, which is equivalent to $ad \leq bc$ or $ad \geq bc$, which is true in \mathbb{Z} .

It is easy to prove $(a, b) \leq (c, d) \implies (a, b) + (p, q) \leq (c, d) + (p, q)$, and $0_{\mathbb{Q}} \leq (a, b), (c, d) \implies 0_{\mathbb{Q}} \leq (a, b) \cdot (c, d)$. However, \mathbb{Q} is **not** complete. We prove this by showing that the subset $X = \{x \in \mathbb{Q} \mid x^2 \leq 2\} \subset \mathbb{Q}$ does not satisfy the least upper bound property. Assume that there is a least upper bound $c \in \mathbb{Q}$. $c \neq \sqrt{2}$ (you should know how to prove irrationality of $\sqrt{2}$!), we have either $c > \sqrt{2}$ or $c < \sqrt{2}$.

1. Let $c < \sqrt{2} \iff c - \sqrt{2} > 0$. By the Archimidean principle, there exists a $k \in \mathbb{N}$ such that $0 < \frac{1}{k} < c - \sqrt{2}$. Then, $\frac{1}{k} \in \mathbb{Q}$ and \mathbb{Q} is a field, so $c - \frac{1}{k} \in \mathbb{Q}$.

$$c - \frac{1}{k} < c - c + \sqrt{2} = \sqrt{2} \quad (162)$$

So c is not least and so it must be the case that $c < \sqrt{2}$.

2. Let $c < \sqrt{2} \iff \sqrt{2} - c > 0$. By the Archimidean principle, there exists a $k \in \mathbb{N}$ such that $0 < \frac{1}{k} < \sqrt{2} - c$. Then, $c + \frac{1}{k} \in \mathbb{Q}$ and

$$c + \frac{1}{k} < c + (\sqrt{2} - c) = \sqrt{2} \quad (163)$$

So c is not an upper bound.

Note that given a well-defined $c = \sup X$ and in the case where $c < \sqrt{2}$, we have $2 - c^2 > 0$, so we can choose a well-defined δ satisfying (by Archimidean principle)

$$0 < \delta < \min \left\{ 1, \frac{2 - c^2}{2c + 1} \right\} \quad (164)$$

which gives us

$$\begin{aligned} (c + \delta)^2 &= c^2 + \delta(2c + \delta) \\ &< c^2 + \delta(2c + 1) & (\delta < 1) \\ &< c^2 + (2 - c^2) = 2 \end{aligned}$$

meaning that c is not an upper bound. Similarly for when $c > \sqrt{2}$.

Exercise 1.34 (Zorich 2.2.15)

Prove the equivalence of these two statements.

1. If X and Y are nonempty sets of \mathbb{R} having the property that $x \leq y$ for every $x \in X, y \in Y$, then there exists $c \in \mathbb{R}$ such that $x \leq c \leq y$ for all $x \in X$ and $y \in Y$.
2. Every set $X \subset \mathbb{R}$ that is bounded above has a least upper bound.

Solution. Let S_1 be the first statement and S_2 the second.

1. ($S_2 \implies S_1$). Let $X \subset \mathbb{R}$ be a set that is bounded above, and Y is a set such that $x \leq y$ for all $x \in X, y \in Y$. Then, by LUB principle, there exists $c = \sup X \in \mathbb{R}$. Now, we claim that $c \leq y$ for all $y \in Y$. Assume it doesn't: then there exists $y' \in Y$ such that $y' < c$. But since we assumed $x \leq y$ for all $x \in X, y \in Y$, we have $x \leq y'$ for all $x \in X$, which means that y' is an upper bound of X . But $y' < c$, contradicting the given fact that c was the least upper bound.
2. ($S_1 \implies S_2$). Given a nonempty set $X \subset \mathbb{R}$, we wish to show the existence of $\sup X$. We are guaranteed the existence of nonempty set $Y \subset \mathbb{R}$ such that $x \leq y$ for all $x \in X, y \in Y$, which implies that X must be bounded above. Then, by S_1 , there must exist a $c \in \mathbb{R}$ such that

$$x \leq c \leq y \text{ for all } x \in X, y \in Y \quad (165)$$

We claim that $c = \sup X$. It is an upper bound of X since $x \leq c$ for all $x \in X$. It is least since the set of all upper bounds of X is Y , and $c \leq y$ for all $y \in Y$.

Exercise 1.35 (Olmsted 1.15)

Prove **Dedekind's Theorem**: Let the real numbers be divided into two nonempty sets A and B such that (i) if $x \in A$ and if $y \in B$, then $x < y$ and (ii) if $x \in \mathbb{R}$ then either $x \in A$ or $x \in B$, then there exists a number c (which may belong to either A or B) such that any number less than c belongs to A and any number greater than c belongs to B .

Solution. This is really the same statement as Zorich 2.2.15.a, the original statement of completeness, but with the extra condition that the sets $A = X, B = Y$ must be disjoint.

Exercise 1.36 (Olmsted 1.7)

If x is an irrational number, under what conditions on the rational numbers a, b, c, d is $(ax + b)/(cx + d)$ rational?

Solution. Note that a trivial solution is $a = b = c = d = 1$ which gives 1. Since

$$\frac{ax + b}{cx + d} = \frac{acx + ad - ad + bc}{cx + d} = a + \frac{bc - ad}{cx + d} \quad (166)$$

for the above to be rational it is necessary that $1/(cx + d)$ is rational. But this cannot be the case, which leaves us with the condition that $bc = ad$.

Exercise 1.37 (Olmsted 1.8)

Prove that the system of integers satisfies the axiom of completeness.

Solution. Let $S \subset \mathbb{Z}$ be bounded from above. It must have a maximum element (justify?), call it c . Then we claim that $c \in \mathbb{Z}$ is the least upper bound. Being the maximum, it is an upper bound, and c is least since the next smallest element is $c - 1$, which is less than $c \in S$, and therefore cannot be an upper bound.

Exercise 1.38 (Zorich 2.2.16/Olmsted 1.16)

Prove the following.

1. If $A \subset B \subset \mathbb{R}$, then $\sup A \leq \sup B$ and $\inf A \geq \inf B$.
2. Let $\mathbb{R} \supset X \neq \emptyset$ and $\mathbb{R} \supset Y \neq \emptyset$. If $x \leq y$ for all $x \in X, y \in Y$, then X is bounded above, Y is bounded below, and $\sup X \leq \inf Y$.
3. If the sets X, Y in (b), are such that $X \cup Y = \mathbb{R}$, then $\sup X = \inf Y$.
4. If X and Y are the sets defined in (c), then either X has a maximal element or Y as a minimal element.
5. Show that Dedekind's theorem is equivalent to the axiom of completeness.

Solution. Listed.

1. Let

$$A' = \{x \in \mathbb{R} \mid x \geq a \forall a \in A\}$$

$$B' = \{x \in \mathbb{R} \mid x \geq b \forall b \in B\}$$

where we can easily verify that $B' \subset A'$. By definition, we get $\sup B = \min B'$ and $\sup A = \min A'$. But since $B' \subset A'$, for any $b' \in B'$, there exists an $a' \in A'$ such that $a' \leq b'$, which implies that $\sup B = \min B' \leq \min A' = \sup A$.

2. X is bounded above by any element of Y . Y is bounded below by any element of X . By the completion axiom, there exists a $c \in \mathbb{R}$ such that

$$x \leq c \leq y \text{ for all } x \in X, y \in Y \quad (167)$$

Since c is an upper bound of X , $\sup X \leq c$ by definition, and since c is a lower bound of Y , $\inf Y \geq c$ by definition. Therefore, $\sup X \leq c \leq \inf Y$.

3. From completeness there exists a $c \in \mathbb{R}$ such that $x \leq c \leq y$ for all $x \in X, y \in Y$. Y is, by definition, the set of *all* upper bounds of X (i.e. *every* upper bound of X is in Y , unlike Y defined in 2.2.16.b). Since $c \leq y$ for all $y \in Y$, c is minimal and so $c = \sup X$. X is the set of all lower bounds of Y by definition, so $c \geq x$ for all $x \in X \implies c = \inf Y$. So, $\inf Y = c = \sup X$.
4. We know that there exists $c = \inf Y = \sup X$. Since $X \cup Y = \mathbb{R}$, c must be in at least X or Y . If $c \in X$, then $c = \sup X = \max X$, and if $c \in Y$, then $c = \inf Y = \min Y$.
5. This is the same statement as Zorich 2.2.15.a (an iff equivalence, not just one way implying).

Exercise 1.39 (Olmsted 1.13)

Let S be a nonempty set of numbers bounded above, and let x be the least upper bound of S . Prove that x has the two properties corresponding to an arbitrary positive number ϵ :

1. every element $s \in S$ satisfies the inequality $s < x + \epsilon$
2. at least one element $s \in S$ satisfies the inequality $s > x - \epsilon$

Solution. Listed.

1. x is an upper bound $\implies s \leq x$ for all $s \in S$, which implies that $s \leq x < x + \epsilon$.
2. By definition, $x - \epsilon$ cannot be an upper bound, so $x - \epsilon \geq s$ for all $s \in S$ is not true. Therefore, there must exist one $s \in S$ such that $s > x - \epsilon$.

Exercise 1.40 (Zorich 2.2.18)

Let $-A$ be the set of numbers of the form $-a$, where $a \in A \subset \mathbb{R}$. Show that $\sup(-A) = -\inf(A)$.

Solution. If A is unbounded below, then $-\inf A = \infty$ and $-A$ is unbounded above, implying that $\sup A = \infty$. Now assume that A is bounded below, then by completeness, it must have a greatest lower bound. Let us define the set $B = \{b \in \mathbb{R} \mid b \leq a \forall a \in A\}$. From 2.2.16.b, we have $b \leq \inf A \leq a$ for all $a \in A, b \in B$. Multiplying by -1 gives $-b \geq -\inf A \geq -a$ for all $a \in A, b \in B$, which is equivalent to saying

$$a \leq -\inf A \leq b \text{ for all } a \in -A, b \in -B \quad (168)$$

by definition of $-A, -B$. $-\inf A$ is clearly an upper bound of $-A$, and since

$$\begin{aligned} B &= \{b \in \mathbb{R} \mid b \leq a \forall a \in A\} \\ &= \{b \in \mathbb{R} \mid -b \geq -a \forall a \in A\} \\ &= \{b \in \mathbb{R} \mid -b \geq a \forall a \in -A\} \end{aligned}$$

implies that $-B = \{b \in \mathbb{R} \mid b \geq a \forall a \in -A\}$ is the set of all upper bounds of A . So, $-\inf A$ is the least upper bound of $-A$, i.e. $-\inf A = \sup(-A)$.

Exercise 1.41 (Zorich 2.2.21)

Show that the set $\mathbb{Q}(\sqrt{n})$ of numbers of the form $a + b\sqrt{n}$ where $a, b \in \mathbb{Q}$, n is a fixed natural number that is not the square of any integer, is an ordered set satisfying the principle of Archimedes but not the axiom of completeness.

Solution. The order on $\mathbb{Q}(\sqrt{n})$ can be embedded from the ordering on the reals by defining the canonical injection map $i : \mathbb{Q}(\sqrt{n}) \rightarrow \mathbb{R}$ and defining for any $x, y \in \mathbb{Q}(\sqrt{n})$,

$$x \leq_{\mathbb{Q}(\sqrt{n})} y \iff i(x) \leq_{\mathbb{R}} i(y) \quad (169)$$

Now, let $h > 0$ be any fixed real number, and $x = (a, b) = a + b\sqrt{n}$. By the Archimidean principle, we can find a $k \in \mathbb{Z}$ such that

$$(k-1)h \leq x \leq kh \text{ for some } x \in \mathbb{Q}(\sqrt{n}) \subset \mathbb{R} \quad (170)$$

We now show that $\mathbb{Q}(\sqrt{n})$ is not complete since it doesn't satisfy the LUB property. Since there are infinite prime numbers in \mathbb{N} , choose a prime number p that is not a factor of n . Then, we are guaranteed that pn is not a perfect square, and can define the set

$$X = \{x \in \mathbb{Q}(\sqrt{n}) \mid x < \sqrt{pn}\} \subset \mathbb{Q}(\sqrt{n}) \quad (171)$$

and assume that $c = c_1 + c_2\sqrt{n} = \sup X$ exists ($c_1, c_2 \in \mathbb{Q}$). Clearly, $c \neq \sqrt{pn} \notin \mathbb{Q}(\sqrt{n})$.

1. Assume $c < \sqrt{pn} \iff 0 < \sqrt{pn} - c \in \mathbb{R}$. By the Archimidean principle, there exists a $k \in \mathbb{N}$ such that $0 < \frac{1}{k} < \sqrt{pn} - c$. Then, we can verify that $c + \frac{1}{k} = (c_1 + \frac{1}{k}) + c_2\sqrt{n} \in \mathbb{Q}(\sqrt{n})$ and

$$c + \frac{1}{k} < c + \sqrt{pn} - c = \sqrt{pn} \implies c + \frac{1}{k} \in X \quad (172)$$

implies that c is not an upper bound. So we must turn to case 2.

2. Assume $c > \sqrt{pn} \iff c - \sqrt{pn} > 0$. By AP, there exists a $k \in \mathbb{N}$ such that $0 < \frac{1}{k} < c - \sqrt{pn}$. Then, we can verify that $c - \frac{1}{k} \in \mathbb{Q}(\sqrt{n})$ and

$$c - \frac{1}{k} > c - c + \sqrt{pn} = \sqrt{pn} \quad (173)$$

implies that $c - \frac{1}{k}$ is an upper bound of X , so c is not least. Therefore, by contradiction, c does not exist.

Exercise 1.42 (Zorich 2.2.22)

Let $n \in \mathbb{N}$ and $n > 1$. In the set $E_n = \{0, 1, \dots, n-1\}$, we define the sum and product of two elements as the remainders when the usual sum and product in \mathbb{R} are divided by n . With these operations on it, the set E_n is denoted \mathbb{Z}_n .

1. Show that if n is not a prime number, then there are nonzero numbers $m, k \in \mathbb{Z}_n$ such that $m \cdot k = 0$, i.e. there exist nonzero zero divisors.
2. Show that if p is prime, then there are no zero divisors in \mathbb{Z}_p and \mathbb{Z}_p is a field.
3. Show that, no matter what the prime p , \mathbb{Z}_p cannot be ordered in a way consistent with the arithmetic operations on it.

Solution. Listed.

1. n is composite implies that there exist $1 < m, k < n$ such that $n = mk$. These factors m, k are precisely the zero divisors of \mathbb{Z}_n since $mk = n \equiv 0 \pmod{n}$.
2. With p prime, assume that there are nontrivial zero divisors $1 < m, k < p$ in \mathbb{Z}_p . Then, $mk \equiv 0 \pmod{p} \implies mk = lp$ for some $l \in \mathbb{N}$. But this implies that m or k must divide p , which is impossible since $1 < m, k < p$. Then prove field axioms.
3. For any field, we must have $0 \leq 1$, because if not, then

$$0 > 1 \implies 0 < 1^{-1} \cdot 1 = 1^{-1} \implies 0 \cdot 0 < 1^{-1} \cdot 1^{-1} = 1 \quad (174)$$

So, $0 \leq 1$ implies that $0 \leq 1 \leq 2 \leq \dots \leq p-1$. But

$$0 + 1 \leq (p-1) + 1 = 0 \quad (175)$$

is false, so any ordering is impossible.

Exercise 1.43 (Zorich 2.2.23)

Show that if \mathbb{R} and \mathbb{R}' are two models of the set of real numbers and $f : \mathbb{R} \rightarrow \mathbb{R}'$ (with $f \neq 0'$) is a mapping such that $f(x+y) = f(x) + f(y)$ and $f(x \cdot y) = f(x) \cdot f(y)$ for any $x, y \in \mathbb{R}$. Prove that f is an order-preserving isomorphism.

Solution. Let $0, 0'$ be the additive identity of \mathbb{R}, \mathbb{R}' , respectively, and $1, 1'$ the multiplicative identity. We claim that $f(0) = 0'$ since

$$\begin{aligned} f(0) &= f(0+0) && \text{(definition of additive identity)} \\ &= f(0) + f(0) && \text{(homomorphism over +)} \end{aligned}$$

which implies that $f(0) + f(0) = f(0) = 0' + f(0)$. Since $f(0)$ lives in field \mathbb{R}' , its additive identity $-f(0)$ is well defined, and we get $f(0) = f(0) + f(0) + (-f(0)) = 0' + f(0) + (-f(0)) = 0'$. We also claim that $f(1) = 1'$ since

$$\begin{aligned} f(1) &= f(1 \cdot 1) && \text{(definition of multiplicative identity)} \\ &= f(1) \cdot f(1) && \text{(homomorphism over \cdot)} \end{aligned}$$

which implies that $f(1) \cdot f(1) = 1' \cdot f(1)$. Since $f(1)$ lives in field \mathbb{R}' , its multiplicative identity $f(1)^{-1}$ is well defined, and we get $f(1) = f(1) \cdot f(1) \cdot f(1)^{-1} = 1' \cdot f(1) \cdot f(1)^{-1} = 1'$. Now that we have proved mapping of identities, this implies the mapping of inverses.

$$\begin{aligned} 0' &= f(0) = f(x-x) = f(x) + f(-x) \implies f(-x) = -f(x) \\ 1' &= f(1) = f(x \cdot x^{-1}) = f(x) \cdot f(x^{-1}) \implies f(x^{-1}) = f(x)^{-1} \end{aligned}$$

With these conditions, we have proved that f is a homomorphism of fields. Now we prove that f is a bijection, but first, we claim that $f(x) = 0' \implies x = 0$. Assume that there exists a nonzero $x \in \mathbb{R}$ such that $f(x) = 0'$. Then, x^{-1} is well defined, and

$$\begin{aligned} f(x) \cdot f(x^{-1}) &= f(x) \cdot f(x)^{-1} = 0' \\ f(x) \cdot f(x^{-1}) &= f(x \cdot x^{-1}) = f(1) = 1' \end{aligned}$$

which implies that $0' = 1'$. So, $f(1) = 1' = 0'$, and so for all $k \in \mathbb{R}$, $f(k) = f(k \cdot 1) = f(k) \cdot f(1) = f(k) \cdot 0' = 0' \implies f \equiv 0'$, leading to a contradiction of the assumption that $f' \not\equiv 0'$.

1. (f injective). Assume f is not injective, i.e. there exists distinct $x_1, x_2 \in \mathbb{R}$ s.t. $f(x_1) = f(x_2)$. Then, using that fact $f(x) = 0 \implies x = 0$,

$$0 = f(x_1) - f(x_2) = f(x_1 - x_2) \implies x_1 - x_2 = 0 \implies x_1 = x_2 \quad (176)$$

2. (f surjective). Let y be any nonzero element in \mathbb{R}' (clearly if $y = 0'$ then its preimage is 0) and y^{-1} its multiplicative inverse. Assume there exists no $x \in \mathbb{R}$ satisfying $f(x) = y$, meaning that there exist no x satisfying

$$f(x) \cdot y^{-1} = 1' \quad (177)$$

But since f maps inverses to inverses, we can choose $x = (y^{-1})^{-1}$, which leads to

$$f(x) \cdot y^{-1} = (\quad) \quad (178)$$

Finally, we prove that f is order preserving. Assume that $x \leq y \iff 0 \leq y - x$, we wish to prove that

$$f(x) \leq f(y) \iff 0 \leq f(y) - f(x) = f(y - x) \quad (179)$$

Therefore, since this preservation of ordering is really the statement $0 \leq y - x \implies 0 \leq f(y - x)$, it suffices to prove that $0 \leq x \implies 0 \leq f(x)$. Now, assume that we have a x such that $f(x) < 0'$. Adding it with the equation $f(1) = 1'$ gives us

$$f(x + 1) < 1' \quad (180)$$

It is easy to prove that $0 \leq x \iff 0 \leq x^{-1}$. Now assume that $0 > f(x)$. **INCOMPLETE**

Exercise 1.44 (Density of Rationals in \mathbb{R})

Prove that for any two distinct $a < b \in \mathbb{R}$, there exists an infinite number of rational numbers between a and b .

Solution. Since $a < b$, then $b - a > 0$ and by the Archimidean principle, there exists a $k \in \mathbb{N}$ such that

$$0 < \frac{1}{k} < b - a \implies 1 < kb - ka \quad (181)$$

which implies that the length of $[ka, kb)$ greater than 1. By the inductive property of \mathbb{Z} , there must be an integer $p \in [ka, kb)$. If there were not, then this would imply that $[ka + 1, kb + 1)$ and $[ka - 1, kb - 1)$ had no integers and repeating would mean that there were no integers in \mathbb{R} . Therefore,

$$ka \leq p < kb \implies a \leq \frac{p}{k} < b \quad (182)$$

for all $a, b \in \mathbb{R}$, with $p/k \in \mathbb{Q}$. If a is irrational we can replace the \leq to $<$, leaving $a < \frac{p}{k} < b$, and if a is rational, we can construct another rational $a + \frac{1}{k} \in (a, b)$.

Exercise 1.45 (Nested Interval Lemma)

With the fact that \mathbb{R} is complete, prove the following.

1. For a sequence of closed nested intervals $I_1 \supset I_2 \supset \dots$ of \mathbb{R} , there exists a point $c \in \mathbb{R}$ belonging to all these intervals.
2. Furthermore, if the hypothesis also satisfies the fact that for any $\epsilon > 0$, there exists a $k \in \mathbb{N}$ such that $|I_k| < \epsilon$ (i.e. the length of the intervals decreases to 0), then the point c common to all sets is unique.

Solution. Listed.

1. Let $I_n = [a_n, b_n]$, with $a_n < b_n$ finite for all $n \in \mathbb{N}$. For all $n \in \mathbb{N}$, we have $I_n = [a_n, b_n]$ and can take the two subsets $X_n = (-\infty, a_n)$ and $Y_n = (b_n, \infty)$, where $x \leq y$ for every $x \in X_n, y \in Y_n$. We also have the fact that $\mathbb{R} = X_n \cup I_n \cup Y_n$. Since \mathbb{R} is complete, there exists a c such that $x \leq c \leq y$ for all $x \in X, y \in Y$. But $x \leq c \iff c \notin X_n$ and $c \leq y \iff c \notin Y_n$, so for all $n \in \mathbb{N}$, c must be in I_n .
2. Since we have proved (a), it now suffices to prove uniqueness of c . Let there be two distinct points $c_1, c_2 \in \mathbb{R}$ belonging to these intervals. Without loss of generality, assume $c_1 - c_2 > 0$, and choose

$$\epsilon = \frac{c_1 - c_2}{3} \quad (183)$$

Then, there should exist a $k \in \mathbb{N}$ such that $|I_k| < \epsilon$. Since I_k must contain c_1 , it must be a subset of $[c_1 - \epsilon, c_1 + \epsilon]$ (should be able to see why) and similarly for c_2 .

$$\begin{aligned} I_k &\subset \left[c_1 - \frac{c_1 - c_2}{3}, c_1 + \frac{c_1 - c_2}{3} \right] = \left[\frac{2c_1 + c_2}{3}, \frac{4c_1 - c_2}{3} \right] = L \\ I_k &\subset \left[c_2 - \frac{c_1 - c_2}{3}, c_2 + \frac{c_1 - c_2}{3} \right] = \left[\frac{-c_1 + 4c_2}{3}, \frac{c_1 + 2c_2}{3} \right] = M \end{aligned}$$

But since $c_1 > c_2 \implies \frac{c_1 + 2c_2}{3} < \frac{2c_1 + c_2}{3}$, L and M are disjoint $\implies I_k$, as a subset of both, leaves us with $I_k = \emptyset$, contradicting that it is a closed interval.

Exercise 1.46 ()

Compactness of Closed Interval in \mathbb{R} Prove that any system of open intervals covering (i.e. an open cover of) a closed interval contains a finite subsystem that covers the closed interval. Another way to state this is by saying that every closed interval of \mathbb{R} is compact.

Solution. A closed interval with a finite open covering is trivially compact since any subcovering is also finite. We only need to deal with when a closed interval $I = [a, b]$ has an infinite open covering $\{U_\alpha\}_{\alpha \in A}$, which means that the set of indices A is infinite. Assume that there exists no finite covering of I . Then, we divide I into two halves

$$I_1 = \left[a, \frac{a+b}{2} \right], \quad I_2 = \left[\frac{a+b}{2}, b \right] \quad (184)$$

and define a subcovering for each of them. That is, we can define $A_1 \subset A$ and $A_2 \subset A$ such that $\{U_\alpha\}_{\alpha \in A_1} \subset \{U_\alpha\}_{\alpha \in A}$ is a covering of I_1 and $\{U_\alpha\}_{\alpha \in A_2} \subset \{U_\alpha\}_{\alpha \in A}$ is a covering of I_2 . At least one of A_1 or A_2 must be infinite, since if they were both finite, then we can define a finite covering $\{U_\alpha\}_{\alpha \in A_1 \cup A_2}$ of I . Choose the interval with the infinite covering and repeat this procedure, which will result in a nested interval that decreases in length by a half.

$$I \supset I_1 \supset I_2 \supset \dots \quad (185)$$

By the nested interval lemma, there exists a unique point c common to all these intervals. But since $c \in [a, b]$, the open cover $\{U\}$ should contain an open interval $(c - \delta_1, c + \delta_2)$ containing c . We wish to prove that this interval is a superset of some I_k in the sequence above, contradicting the fact that I_k has an infinite cover. Since the length of each I_i decreases arbitrarily (i.e. we can choose any $\epsilon > 0$ and find a I_k with length less than ϵ), we choose $\epsilon = \frac{1}{2} \min\{\delta_1, \delta_2\}$, and we should be able to find some I_k that is a subinterval of $[c - \epsilon, c + \epsilon]$, which itself is a subinterval of $(c - \delta_1, c + \delta_2)$.

$$I_k \subset \left[c - \frac{1}{2} \min\{\delta_1, \delta_2\}, c + \frac{1}{2} \min\{\delta_1, \delta_2\} \right] \subset (c - \delta_1, c + \delta_2) \quad (186)$$

Therefore, $(c - \delta_1, c + \delta_2)$ is a finite cover of I_k , contradicting the fact that all I_k 's have infinite covers.

Exercise 1.47 (Bolzano-Weierstrass Theorem)

Prove that every bounded infinite set of real numbers has at least one limit point. (A limit point p of set X is a point such that every open neighborhood of p contains an infinite number of elements of X).

Solution. Let the set of points be denoted X , and let a be the lower bound and b be the upper bound. Then, $X \subset [a, b] = I$. Now divide $[a, b]$ into halves $[a, \frac{a+b}{2}] \cup [\frac{a+b}{2}, b]$. At least one of the halves must have an infinite number of points; choose the interval with infinite points as I_1 and doing this repeatedly gives the nested sequence

$$I \supset I_1 \supset I_2 \supset \dots \quad (187)$$

By the nested interval lemma, there exists at least one point $c \in \mathbb{R}$ that is in all these intervals. Furthermore, since $|I_i| = \frac{1}{2^i}(b - a)$ decreases to 0, we can choose a $\epsilon > 0$ and find an interval I_k with $|I_k| < \epsilon$. We claim that c is a limit point of X . Given an ϵ , we wish to prove that there are an infinite number of points within the ϵ -neighborhood $(c - \epsilon, c + \epsilon)$ of c . Since we can find some I_k with $|I_k| < \epsilon$, we can see that

$$I_k \subset (c - \epsilon, c + \epsilon) \quad (188)$$

and therefore the ϵ -neighborhood of c contains I_k , which contains an infinite number of points in X . We can construct another proof that is dependent on the compactness lemma. This construction will be useful for problem 2.3.4. Let X be a given subset of \mathbb{R} , and it follows from the definition of boundedness that X is contained in some closed interval $I \subset \mathbb{R}$. We show that at least one point of I is a limit point of X . Assume that it is not. Then each point $x \in I$ would have a neighborhood $U(x)$ containing at most a finite number of points from X . The totality of such neighborhoods $\{U(x)\}$ constructed for the points $x \in I$ forms an open covering of X . Since I is closed, it is compact and therefore we can find a finite subcovering $\{U_i(x)\}_i$ of open intervals that cover I and therefore cover X . This open cover $\{U_i(x)\}_i$ of X is a finite union of sets that each contain at most a finite number of points from X , so the covering of X contains a finite number of points from X , a contradiction that X contains infinite points.

Exercise 1.48 (Zorich 2.3.1)

Show that

1. if I is any system of nested closed intervals, then

$$\sup\{a \in \mathbb{R} \mid [a, b] \in I\} = \alpha \leq \beta = \inf\{b \in \mathbb{R} \mid [a, b] \in I\}$$

and

$$[\alpha, \beta] = \bigcap_{[a, b] \in I} [a, b]$$

2. if I is a system of nested open intervals (a, b) , the intersection

$$\bigcap_{(a,b) \in I} (a, b)$$

may happen to be empty.

Solution. Listed.

1. (May be tempted to say that $a_1 \leq a_2 \leq \dots$, but this assumes that the indexing set I is countable). We claim that for any two intervals $[a_n, b_n]$ and $[a_m, b_m]$ in I ,

$$a_n \leq b_m$$

Assume that $a_n > b_m$. Then $b_n \geq a_n > b_m \geq a_m$ implies that $[a_n, b_n]$ and $[a_m, b_m]$ are disjoint, contradicting the fact that they are nested. Now given that X is the set of a_n 's and Y is the set of b_n 's, we have $x \leq y$ for all $x \in X, y \in Y$. So by 2.2.16.b, we have $\sup X \geq \inf Y$.

To prove the second statement, we show that trying to “expand” the interval $[\alpha, \beta]$ will lead to a contradiction. Since α is the LUB, given any $\epsilon > 0$, there exists a $(a_l, b_l) \in X$ such that $\alpha - \epsilon < a_l < \alpha$, which implies that $[\alpha, \beta] \subset [a_l, \beta] \subset [\alpha - \epsilon, \beta]$. Assuming that this extended interval is the intersection, we should be able to choose any point in $[\alpha - \epsilon, \beta]$ and find that it is in every element of I . We choose a point in $[\alpha - \epsilon, a_l)$, which is not in the interval (a_l, b_l) . We do the same for $\beta \mapsto \beta + \epsilon$. We also check that “shrinking” the interval $[\alpha, \beta] \mapsto [\alpha + \epsilon, \beta]$ is no good, since we can find an element in $[\alpha, \alpha + \epsilon)$ that is in every interval in I .

2. Take the system of nested open intervals

$$(0, 1) \supset (0, \frac{1}{2}) \supset (0, \frac{1}{3}) \dots (0, \frac{1}{n}) \supset \dots$$

Take their infinite intersection, denote it S , and assume that some $\epsilon \in (0, 1)$ is in S . Since ϵ is a real number, by the Archimidean principle there exists a $k \in \mathbb{N}$ such that $\frac{1}{k} < \epsilon$. Therefore, $\epsilon \notin (0, \frac{1}{k}) \implies \epsilon \notin S$.

Exercise 1.49 (Zorich 2.3.2)

Show that

1. from a system of closed intervals covering a closed interval it is not always possible to choose a finite subsystem covering the interval.
2. from a system of open intervals covering a open interval it is not always possible to choose a finite subsystem covering the interval.
3. from a system of closed intervals covering a open interval it is not always possible to choose a finite subsystem covering the interval.

Solution. We show with the interval $(0, 1)$ or $[0, 1]$. Using linear transformations it is easy to generalize this to any other interval (a, b) or $[a, b]$.

1. Consider the infinite covering

$$[0, 1] = [0, \frac{1}{2}] \cup [\frac{1}{2}, \frac{3}{4}] \cup [\frac{3}{4}, \frac{7}{8}] \cup \dots$$

2. Consider the infinite covering

$$(0, 1) = (0, \frac{1}{2}) \cup (\frac{1}{2}, \frac{3}{4}) \cup (\frac{3}{4}, \frac{7}{8}) \cup \dots$$

3. Consider the infinite covering

$$(0, 1) = [0, \frac{1}{2}] \cup [\frac{1}{2}, \frac{3}{4}] \cup [\frac{3}{4}, \frac{7}{8}] \cup \dots$$

Exercise 1.50 (Zorich 2.3.3)

Show that if we only take the set \mathbb{Q} of rational numbers instead of the complete set \mathbb{R} of real numbers, with the definitions of closed, open, and neighborhood of a point $r \in \mathbb{Q}$ to mean respectively the corresponding subsets of \mathbb{Q} , then none of the three lemmas is true.

Solution. We prove only for the nested interval lemma. We choose the series of nested intervals

$$\left(\sqrt{2} - \frac{1}{n}, \sqrt{2} + \frac{1}{n} \right)$$

with $n \in \mathbb{N}$. Assume that there is a $r \in \mathbb{Q}$ such that

$$r \in \left(\sqrt{2} - \frac{1}{n}, \sqrt{2} + \frac{1}{n} \right) \text{ for all } n \in \mathbb{N}$$

which is equivalent to saying that $|r - \sqrt{2}| < \frac{1}{n}$ for all $n \in \mathbb{N}$. Clearly, $r \neq \sqrt{2}$, and by the Archimidean principle, there exists a $k \in \mathbb{N}$ such that

$$0 < \frac{1}{k} < |r - \sqrt{2}|$$

which contradicts the above.

Exercise 1.51 (Zorich 2.3.4)

Show that the three lemmas above are equivalent to the axiom of completeness.

Solution. Note that from the proofs, completeness implies nested interval lemma, which implies compactness of closed intervals, which implies the Bolzano-Weierstrass theorem. So, it is sufficient to prove that Bolzano-Weierstrass theorem implies completeness to determine equivalence. There are not a lot of direct proofs, so we prove that Weierstrass implies nested interval, which implies completeness.

1. (Weierstrass \implies Nested) Assume that we have \mathbb{R} with the Bolzano-Weierstrass theorem. Take the series of nested closed intervals

$$I = [a, b] \supset I_1 = [a_1, b_1] \supset I_2 = [a_2, b_2] \supset \dots$$

We see that $a \leq a_i \leq b$, so the infinite sequence of monotonically nondecreasing values a_i is bounded. Therefore, it must have a limit point, which we will denote as c . We claim that $a_i \leq c$ for all a_i . Since if it were not, then $c < a_i$ for some i , and choosing $\epsilon = 0.5(a_i - c)$, the ϵ -neighborhood of c will not contain a_j for $j \geq i$ since

$$c < a_i \implies 0.5c < 0.5a_i \implies c + \epsilon = 0.5c + 0.5a_i < a_i < a_{i+1} < \dots$$

. With similar reasoning, we can conclude that $b_i \geq c$ for all b_i . This implies that $a_i \leq c \leq b_i$ for all i which is equivalent to saying that $c \in [a_i, b_i] = I_i$ for all $i \in \mathbb{N}$.

2. (Nested \implies LUB Principle) Let $X \subset \mathbb{R}$ be a set that is bounded above, with b_1 any upper bound. Since X is nonempty, there exists $a_1 \in X$ that is not an upper bound (otherwise, X would be a singleton set and it trivially has a least upper bound). Consider the well-defined

interval $[a_0, b_0]$. Take the mean $m_0 = 0.5(a_0 + b_0)$, and if m_0 is an upper bound, set it to b_1 (with $a_1 = a_0$) and a_1 if else (with $b_1 = b_0$). Then, we have a sequence of nested intervals

$$[a_0, b_0] \supset [a_1, b_1] \supset [a_2, b_2] \supset \dots$$

of decreasing lengths $|I_k| = \frac{1}{2^k}(b - a)$. All of them must contain a unique common point $c \in \mathbb{R}$ by the nested intervals lemma, which implies that

$$a_0 \leq a_1 \leq a_2 \leq \dots \leq c \leq \dots \leq b_2 \leq b_1 \leq b_0$$

I claim two things:

- (a) c is an upper bound for X . Suppose it were not, then there exists some $x \in X$ such that $c < x$, and let the distance between them be $\epsilon = x - c > 0$. By AP, we can choose $k \in \mathbb{N}$ such that $\frac{1}{k} < \epsilon$. All the b_n are upper bounds of X , so we have $x \leq b_n$. Subtracting c on both sides gives

$$0 < x - c = \epsilon \leq b_n - c \leq |I_n| = \frac{1}{2^n}(b_0 - a_0)$$

where the last inequality follows from $c \in I_n = [a_n, b_n]$, so the maximum distance it can be from the endpoint b_n is $|I_n|$. The inequality above holds for all $n \in \mathbb{N}$, so increasing n arbitrarily should decrease $\frac{1}{2^n}(b_0 - a_0)$ past ϵ . To formalize this, we use the inequality

$$\frac{1}{2^n} < \frac{1}{n} \text{ for all } n \in \mathbb{N}$$

and so we have

$$\epsilon \leq b_n - c < \frac{1}{n}(b_0 - a_0)$$

We choose the natural number $n = \lceil \frac{2(b_0 - a_0)}{\epsilon} \rceil$, which does not satisfy the inequality above since

$$\epsilon < \frac{1}{n}(b_0 - a_0) = \frac{1}{\lceil 2(b_0 - a_0)/\epsilon \rceil}(b_0 - a_0) \leq \frac{\epsilon}{2(b_0 - a_0)}(b_0 - a_0) = \frac{\epsilon}{2}$$

This leads to a contradiction.

- (b) We now prove that c is least. Assume that c is not least \implies there exists an upper bound B such that $B < c$ and $x \leq B$ for all $x \in X$. **INCOMPLETE**

Exercise 1.52 (Zorich 2.4.1)

Show that the set of real numbers has the same cardinality as the points of the interval $(-1, 1)$.

Solution. We define the bijective map $\rho : (-1, 1) \longrightarrow \mathbb{R}$

$$\rho(x) = \begin{cases} 0 & \text{if } x = 0 \\ \frac{1}{x} & \text{if } x \neq 0 \end{cases}$$

Exercise 1.53 (Zorich 2.4.2)

Give an explicit one-to-one correspondence between

1. the points of two open intervals
2. the points of two closed intervals
3. the point of a closed interval and an open interval
4. the points of the closed interval $[0, 1]$ and \mathbb{R}

Solution. Listed.

1. $\rho : (a, b) \longrightarrow (c, d)$ defined

$$\rho(x) = \frac{d-c}{b-a}(x-a) + c$$

2. the extension of ρ defined on (a) to $[a, b]$
3. From (a) and (b), it suffices to prove a bijection from $(0, 1)$ to $[0, 1]$. We extract a countably infinite sequence from $(0, 1)$, say

$$x_1 = \frac{1}{3}, x_2 = \frac{1}{4}, \dots, x_i = \frac{1}{i+2}$$

Then, we define bijection $\rho : (0, 1) \longrightarrow [0, 1]$ as

$$\rho(x) = \begin{cases} x & \text{if } x \notin \{x_i\} \\ 0 & \text{if } x = x_1 = \frac{1}{2} \\ 1 & \text{if } x = x_2 = \frac{1}{3} \\ x_{i-2} & \text{if } x = x_i \text{ for } i > 2 \end{cases}$$

Colloquially, we extract a copy of \mathbb{N} from $(0, 1)$ and use the bijection $\mathbb{N} \simeq \mathbb{N} \cup \{0, -1\}$ to “shift” the terms.

4. We compose the bijections $\rho_1 : [0, 1] \longrightarrow (0, 1)$ and $\rho_2 : (0, 1) \longrightarrow \mathbb{R}$.

Exercise 1.54 (Zorich 2.4.3)

Show that

1. every infinite set contains a countable subset
2. the set of even integers has the same cardinality as the set of all natural numbers
3. the union of an infinite set and an at most countable set has the same cardinality as the original infinite set.
4. the set of irrational numbers has the cardinality of the continuum
5. the set of transcendental numbers has the cardinality of the continuum

Solution. Listed.

1. Let A be an infinite set. By axiom of choice, choose $a_0 \in A$. Then, $A \setminus \{a_0\} \neq \emptyset$ since A is infinite. By induction, assume you have chosen $a_0, a_1, \dots, a_k \in A$. Then, since A is infinite, $A \setminus \{a_0, a_1, \dots, a_k\} \neq \emptyset$, so we can choose $a_{k+1} \in A \setminus \{a_0, \dots, a_k\}$. Thus, we have constructed a countable subset $\{a_k\}_{k \in \mathbb{N}}$ of A .
2. Given the quotient ring $2\mathbb{Z}$, define the bijection $\rho : 2\mathbb{Z} \longrightarrow \mathbb{N}$ as

$$p(x) = \begin{cases} x+2 & \text{if } x \geq 0 \\ -x-1 & \text{if } x < 0 \end{cases}$$

3. From (a), we can extract a countable set from original set A , call it X . Since the product of countable sets is countable ($\mathbb{N} \cup \mathbb{N}$ is countable), we can define a bijection $\tilde{\rho} : X \longrightarrow X \cup B$. Therefore, we can define a bijection $\rho : A \longrightarrow A \cup B$ as

$$\rho(x) = \begin{cases} x & \text{if } x \in A \setminus X \\ \tilde{\rho}(x) & \text{if } x \in X \end{cases}$$

4. \mathbb{Q} is countable and \mathbb{R} is uncountable. So, $\mathbb{R} \setminus \mathbb{Q}$ must be uncountable since if it were countable, then the union of the rationals and irrationals, which is \mathbb{R} , would be countable.

5. It suffices to prove that the set of algebraic numbers (numbers that are possible roots of a polynomial with integer coefficients with leading coefficient nonzero) is countable, since we can apply (d) right after. The set of all k th degree polynomials with integer coefficients is isomorphic to \mathbb{Z}^k through the map

$$a_k x^k + a_{k-1} x^{k-1} + \dots + a_2 x^2 + a_1 x^1 + a_0 \mapsto (a^{k-1}, a^{k-2}, \dots, a_1, a_0)$$

and the union of these countable sets (minus the 0 map)

$$P = \left(\bigcup_{k=1}^{\infty} \mathbb{Z}^k \right) \setminus \{0\} = (\mathbb{Z} \setminus \{0\}) \cup \mathbb{Z}^2 \cup \dots$$

is countable. For any element in \mathbb{Z}^k , there are at most k real roots, and so we can define the set of roots of an element $z \in \mathbb{Z}^k \subset P$ as a j -tuple of algebraic numbers, which can have at most $j = k$ roots.

$$r(z) = \underbrace{(r_{1z}, r_{2z}, \dots, r_{jz})}_{j \leq k}$$

Therefore, the union of all these j -tuples for all $z \in P$

$$\bigcup_{z \in P} r(z) = \bigcup_{k=1}^{\infty} \bigcup_{z \in \mathbb{Z}^k} r(z)$$

is a countable union of a countable union of finite sets, making it countable.

Exercise 1.55 (Zorich 2.4.4)

Show that

1. the set of increasing sequences of natural numbers has the same cardinality as the set of fractions of the form $0.\alpha_1\alpha_2\dots$
2. the set of all subsets of countable set has cardinality of the continuum

Solution. Listed.

1. Given a sequence of increasing naturals $S = (n_1, n_2, \dots)$, we can define a binary expansion $0.\alpha_1\alpha_2\dots$ where $\alpha_i = 1$ if and only if $i \in \mathbb{N}$ is in S and $\alpha_i = 0$ if not. This is clearly a bijection.
2. The set of all segments of increasing natural is equipotent with $2^{\mathbb{N}}$, since the elements of each sequence define a subset of \mathbb{N} . Cantor's diagonalization argument proves that the set of infinite binary expansions is uncountable, and by (a), this proves that $2^{\mathbb{N}}$ is uncountable.

This is very interesting since $\mathbb{N} \simeq \mathbb{R}$, but $2^{\mathbb{N}} \simeq \mathbb{R}$, and the set of all infinite q -ary expansions is equipotent to \mathbb{R} too.

Exercise 1.56 (Zorich 2.4.5)

Show that

1. the set $\mathcal{P}(X)$ of subsets of a set X has the same cardinality as the set of all functions $f : X \rightarrow \{0, 1\}$.
2. for a finite set X of n elements, $\text{card } \mathcal{P}(X) = 2^n$
3. one can write $\text{card } \mathcal{P}(X) = 2^{\text{card } X}$, which implies $\text{card } \mathcal{P}(\mathbb{N}) = 2^{\text{card } \mathbb{N}} = \text{card } \mathbb{R}$
4. for any set X , $\text{card } X < 2^{\text{card } X}$

Solution. Listed.

1. An element $Y \in \mathcal{P}(X)$ is a subset of X by definition. Letting

$$f_Y(x) = \begin{cases} 0 & \text{if } x \notin Y \\ 1 & \text{if } x \in Y \end{cases}$$

we can construct the bijective map $Y \mapsto f_Y$.

2. We can prove this using the identity (which can be proved using induction)

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

3. Let $F(X; \{0, 1\})$ be the set of all binary valued functions from X to $\{0, 1\}$. From (a), $\text{card } \mathcal{P}(X) \simeq F(X; \{0, 1\})$. Each binary-valued function f is determined by the assignment $f(x)$ for each $x \in X$. Since $f(x)$ has two possible values, the assignment of $f(x)$ for all $x \in X$ has $\{0, 1\}^{\text{card } X}$ possible choices. This gives another bijection $F(X; \{0, 1\}) \simeq \{0, 1\}^{\text{card } X}$, so

$$\mathcal{P}(X) \simeq \{0, 1\}^{\text{card } X} \implies \text{card } \mathcal{P}(X) = \text{card}(\{0, 1\}^{\text{card } X}) = 2^{\text{card } X}$$

4. If X is finite, then letting $n = \text{card } X$, we can simply prove $n < 2^n$ by induction (which we will not do here). If X is countable, then $\mathcal{P}(X)$ is uncountable (from 2.4.4) and so using (c),

$$\text{card } X = \text{card } \mathbb{N} < \text{card } \mathbb{R} = \text{card } \mathcal{P}(X) = 2^{\text{card } X}$$

For uncountable sets (and for the two cases mentioned above), we can use Cantor's theorem, which states that $\text{card } X < \text{card } \mathcal{P}(X)$, and so using (c), we have $\text{card } X < \text{card } \mathcal{P}(X) = 2^{\text{card } X}$.

Exercise 1.57 (Zorich 2.4.6)

Let X_1, \dots, X_m be a finite system of finite sets. Show that

$$\begin{aligned} \text{card} \left(\bigcup_{i=1}^m X_i \right) &= \sum_{i_1} \text{card } X_{i_1} - \sum_{i_1 < i_2} \text{card}(X_{i_1} \cap X_{i_2}) + \dots \\ &\quad \sum_{i_1 < i_2 < i_3} \text{card}(X_{i_1} \cap X_{i_2} \cap X_{i_3}) - \dots + (-1)^{m-1} \text{card}(X_1 \cap \dots \cap X_m) \\ &= \sum_{k=1}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k X_{i_j} \right) \end{aligned}$$

Solution. Ignoring Russell's paradox (defining the universe set of all sets), we can use the commutative, associative, and distributive properties of \cup, \cap on the algebra of sets. We prove using induction on m . For $m = 1$, we trivially have $\text{card } X_1 = \text{card } X_1$, and for $m = 2$, we claim

$$\text{card}(X_1 \cup X_2) = \text{card}(X_1) + \text{card}(X_2) - \text{card}(X_1 \cap X_2)$$

X_1 and $X_2 \setminus X_1$ are clearly exclusive sets by definition, with $X_1 \cup X_2 = X_1 \cup (X_2 \setminus X_1)$, so

$$\text{card}(X_1 \cup X_2) = \text{card}(X_1 \cup (X_2 \setminus X_1)) = \text{card}(X_1) + \text{card}(X_2 \setminus X_1) \quad (2)$$

By definition, the set $X_2 \setminus X_1$ and $X_1 \cap X_2$ are disjoint and satisfies $X_2 = (X_2 \setminus X_1) \cup (X_1 \cap X_2)$ (also by definition), so

$$\text{card}(X_2) = \text{card}(X_2 \setminus X_1) + \text{card}(X_1 \cap X_2) \quad (3)$$

and substituting (3) into (2) gives the claim for $m = 2$. Assuming that the claim is satisfied for some m , we have

$$\begin{aligned}
 \text{card} \left(\bigcup_{i=1}^{m+1} X_i \right) &= \text{card} \left(\left[\bigcup_{i=1}^m X_i \right] \cup X_{m+1} \right) \\
 &= \text{card} \left(\bigcup_{i=1}^k X_i \right) + \text{card}(X_{m+1}) - \text{card} \left(\left[\bigcup_{i=1}^m X_i \right] \cap X_{m+1} \right) \quad (\text{claim for } m = 2) \\
 &= \text{card} \left(\bigcup_{i=1}^k X_i \right) + \text{card}(X_{m+1}) - \text{card} \left(\bigcup_{i=1}^m (X_i \cap X_{m+1}) \right) \quad (\text{distributive prop.}) \\
 &= \sum_{k=1}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k X_{i_j} \right) + \text{card}(X_{m+1}) \\
 &\quad - \sum_{k=1}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k (X_{i_j} \cap X_{m+1}) \right)
 \end{aligned}$$

With a bit of thought, we can see that the k th term of the second summation contributes to adding another term to the $k+1$ th summation term of the first. Therefore, we must try to shift the summation over by 1 index. Let us simplify this by taking the summations and extracting the first and last term, respectively. We have

$$\begin{aligned}
 \sum_{k=1}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k X_{i_j} \right) &= \sum_{1 \leq i_1 \dots i_k \leq m} \text{card}(X_{i_1}) \\
 &\quad + \sum_{k=2}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k X_{i_j} \right)
 \end{aligned}$$

and

$$\begin{aligned}
 \sum_{k=1}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k (X_{i_j} \cap X_{m+1}) \right) \\
 &= \sum_{k=1}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\left[\bigcap_{j=1}^k X_{i_j} \right] \cap X_{m+1} \right) \\
 &= \sum_{k=1}^{m-1} \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k (X_{i_j} \cap X_{m+1}) \right) + (-1)^{m-1} \text{card} \left(\bigcap_{j=1}^{m+1} X_j \right) \\
 &= \sum_{k=2}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-2} \text{card} \left(\bigcap_{j=1}^{k-1} (X_{i_j} \cap X_{m+1}) \right) + (-1)^{m-1} \text{card} \left(\bigcap_{j=1}^{m+1} X_j \right)
 \end{aligned}$$

So subtracting the summations gives

$$\begin{aligned}
& \sum_{k=1}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k X_{i_j} \right) - \sum_{k=1}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k (X_{i_j} \cap X_{m+1}) \right) + |X_{m+1}| \\
&= \sum_{1 \leq i_1 \dots i_k \leq m} \text{card}(x_i) + \sum_{k=2}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k X_{i_j} \right) + \text{card}(X_{m+1}) \\
&\quad + \sum_{k=2}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^{k-1} (X_{i_j} \cap X_{m+1}) \right) + (-1)^m \text{card} \left(\bigcap_{j=1}^{m+1} X_j \right) \\
&= \sum_{1 \leq i_1 \dots i_k \leq m+1} \text{card}(X_i) + \sum_{k=2}^m \sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \left[\text{card} \left(\bigcap_{j=1}^k X_{i_j} \right) \right. \\
&\quad \left. + \text{card} \left(\left[\bigcap_{j=1}^{k-1} X_{i_j} \right] \cap X_{m+1} \right) \right] + (-1)^m \text{card} \left(\bigcap_{j=1}^{m+1} X_j \right)
\end{aligned}$$

and since the set of sequences of k terms bounded by $m+1$ (of form $1 \leq i_1 \dots i_k \leq m+1$) is the set of sequences of k terms bounded by m (of form $1 \leq i_1 \dots i_k \leq m$) unioned with the set of sequences of k terms with max element $m+1$ (of form $1 \leq i_1 \dots i_k = m+1$), we have

$$\sum_{1 \leq i_1 \dots i_k \leq m} (-1)^{k-1} \left[\text{card} \left(\bigcap_{j=1}^k X_{i_j} \right) + \text{card} \left(\left[\bigcap_{j=1}^{k-1} X_{i_j} \right] \cap X_{m+1} \right) \right] = \sum_{1 \leq i_1 \dots i_k \leq m+1} \text{card} \left(\bigcap_{j=1}^k X_{i_j} \right)$$

and therefore, substituting the above and observing that the independent terms are the first and last terms of the summation gives

$$\begin{aligned}
\text{card} \left(\bigcup_{i=1}^{m+1} X_i \right) &= \sum_{1 \leq i_1 \dots i_k \leq m+1} \text{card}(X_i) + \sum_{k=2}^m \sum_{1 \leq i_1 \dots i_k \leq m+1} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k X_{i_j} \right) \\
&\quad + \dots + (-1)^m \text{card} \left(\bigcap_{j=1}^{m+1} X_j \right) \\
&= \sum_{k=1}^{m+1} \sum_{1 \leq i_1 \dots i_k \leq m+1} (-1)^{k-1} \text{card} \left(\bigcap_{j=1}^k X_{i_j} \right)
\end{aligned}$$

Exercise 1.58 (Zorich 2.4.7)

On the closed interval $[0, 1] \subset \mathbb{R}$, describe the sets of numbers $x \in [0, 1]$ whose ternary representation $x = 0.\alpha_1\alpha_2\dots$, $\alpha_i \in \{0, 1, 2\}$ has the property.

1. $\alpha_1 \neq 1$
2. $\alpha_1 \neq 1$ and $\alpha_2 \neq 1$
3. For all $i \in \mathbb{N}$, $\alpha_i \neq 1$ (the Cantor set)

Solution. Listed.

1. $[0, \frac{1}{3}) \cup [\frac{2}{3}, 1)$
2. $[0, \frac{1}{9}) \cup [\frac{2}{9}, \frac{3}{9}) \cup [\frac{6}{9}, \frac{7}{9}) \cup [\frac{8}{9}, 1)$
3. Made by recursively removing the middle third of every partitioned intervals.

Exercise 1.59 (Zorich 2.4.8)

Show that

1. the set of numbers $x \in [0, 1]$ whose ternary representation does not contain 1 has the same cardinality as the set of all numbers whose binary representation has the form $0.\beta_1\beta_2\dots$
2. the Cantor set has the same cardinality as the closed interval $[0, 1]$

Solution. Listed.

1. We can define a bijection $0.\alpha_1\alpha_2\dots \mapsto 0.\beta_1\beta_2\dots$ as $\alpha_i = 0 \iff \beta_i = 0$ and $\alpha_i = 1 \iff \beta_i = 2$.
2. The map above defines a bijection between the Cantor set and the set of all infinite binary expansions in $[0, 1]$, which is uncountable by Cantor's diagonalization theorem.

2 Euclidean Topology

With the construction of the real line and the real space, the extra properties of completeness, norm, and order (for the real line) allows us to restate these topological properties in terms of these “higher-order” properties. It also proves much more results than for general topological spaces. Therefore, the next few sections will focus on reiterating the topological properties of \mathbb{R} and \mathbb{R}^n (this can be done slightly more generally for metric spaces, but we talk about this in point-set topology).

Theorem 2.1 (Euclidean Topology)

The **Euclidean topology** \mathcal{T} on \mathbb{R}^n is defined in the equivalent ways:

1. If $n = 1$, it is the order topology generated by the basis of open intervals (a, b) for $a < b$.
2. It is the metric topology generated by the basis of open balls under the L^p metric for $p \geq 1$.
3. It is the n -fold product topology of the Euclidean topology on \mathbb{R} .

Proof. Let's call these sets $\mathcal{T}_1, \mathcal{T}_2, \mathcal{T}_3$. We first show that these are indeed topologies.

1. This is proven in the definition of the order topology.
2. This is proven for the L^2 metric in the definition of the metric topology, and then we prove equivalence to L^p metrics.
3. This is proven in the definition of the product topology.

Now it remains to show that these topologies are all the same. For $n = 1$, it suffices to show $\mathcal{T}_1 = \mathcal{T}_2$.

We do this by comparing their bases; let \mathcal{B}_k be the basis of \mathcal{T}_k .

1. $\mathcal{T}_1 \subset \mathcal{T}_2$. Given any open interval $(a, b) \in \mathcal{B}_1$, (a, b) is the open ball with center $\frac{a+b}{2}$ and radius $\frac{b-a}{2}$ and so it is in \mathcal{B}_2 .
2. $\mathcal{T}_2 \subset \mathcal{T}_1$. Given an open interval with center x and radius r , the corresponding interval $(x-r, x+r)$ is in \mathcal{B}_1 .

Now we show that $\mathcal{T}_2 = \mathcal{T}_3$ for \mathbb{R}^n .

1. We show that the product topology is the same as the metric topology under the L^∞ norm.
2. The equivalence of all L^p norms is already proven here.

Example 2.1 (Examples of Open and Not Open Sets)

Here are some examples of sets which are open and not open.

1. $U = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \neq 1\}$ is open since for every point $x \in U$, we just need to find a radius $\epsilon > 0$ that is smaller than its distance to the unit circle.
2. $(a, b) \times (c, d) \subset \mathbb{R}^2$ is open since given a point x , we can take the minimum of its distance between the two sides of the rectangle and construct an open ball.
3. $S = \{(x, y) \in \mathbb{R}^2 : xy \neq 0\}$ is open since given a point $x \in S$, we can take the minimum of the distance between it and the x and y axes.
4. The set of all complex z such that $|z| \leq 1$ is not open since we cannot construct open balls at the boundary points that are fully contained in the set.
5. The set $S = \{1/n\}_{n \in \mathbb{N}}$ is not open since given any point $x = 1/n$, we can construct an open ball with radius $\epsilon < 1/(n+1)$, which contains irrationals that are not in S .

Corollary 2.1 ()

Every closed bounded interval in \mathbb{R} is compact.

Proof. Let $I = [a, b]$. Then if $x, y \in I$, $|x - y| \leq b - a = \delta$. Now by contradiction, suppose that there exists an open cover $\{U_\alpha\}$ of I which contains no finite subcover of I . Then letting $c = (a + b)/2$, at least one of the two intervals $[a, c], [c, b]$ cannot have a finite subcovering (otherwise their finite union

can be covered). WLOG let it be $[a, c]$. We keep subdividing and get the sequence of nested intervals.

$$I \supset I_1 \supset I_2 \supset \dots \quad (189)$$

We know that I_n is not covered by any finite subcollection of $\{U_\alpha\}$ and if $x, y \in I_n$, then $|x - y| < 2^{-n}\delta$. From the nested intervals theorem, there exists a point z lying in every I_n . There must then be an open neighborhood U_z in the open cover, and by definition of openness there exists a $\epsilon > 0$ s.t. $z \in B_\epsilon(z) \subset U_z$. By the Archimidean property, we can set n so large that $2^{-n}\delta < \epsilon$ and this means that $B_\epsilon(z) \supset I_n$, which contradicts the fact that I_n is not covered by a finite subcollection. Therefore I is compact.

Now here are some useful properties.

Theorem 2.2 (Countability)

\mathbb{R}^n is 2nd countable.

Theorem 2.3 (Separability)

\mathbb{R}^n is Hausdorff.

Proof. It is a metric topology.

Theorem 2.4 (Connectedness)

\mathbb{R}^n is

1. connected,
2. path connected,
3. locally connected,
4. locally path connected

Theorem 2.5 (Compactness)

\mathbb{R}^n is

1. not compact,
2. locally compact.

Theorem 2.6 (Convexity)

An open ball is convex in a normed vector space. What happens if we weaken it to a metric?

Proof. The normed part is important here, as the properties of the metric is not sufficient. Given $B_r(p)$, $x, y \in B_r(p)$ implies that $\|x - p\| < r$ and $\|y - p\| < r$. Therefore,

$$\|tx + (1 - t)y - p\| = \|tx - tp + (1 - t)y - (1 - t)p\| \quad (190)$$

$$\leq t\|x - p\| + (1 - t)\|y - p\| \quad (191)$$

$$= tr + (1 - t)r = r \quad (192)$$

Theorem 2.7 ()

Let E be a nonempty set of real numbers which is bounded above. Let $y = \sup E$. Then $y \in \overline{E}$. Hence $y \in E$ if E is closed.

Proof. Assume that y is not a limit point of E . Then, there exists some $\epsilon > 0$ s.t. $(y - \epsilon, y + \epsilon)$ does not intersect with E . This means that $y - \epsilon$ is an upper bound of E , and so y is not the supremum.

The general notion of compactness for topological spaces is not needed for analysis. Rather, we make use of the following theorem which allows us to focus on the compactness of subsets in Euclidean spaces \mathbb{R}^n .

Theorem 2.8 (Heine-Borel)

Let $E \subset \mathbb{R}^n$. The following are equivalent.

1. E is closed and bounded.
2. E is compact.
3. E is sequentially compact.
4. E is limit point compact.

Proof. To prove the equivalence of the last three, note that X is a metric space, so it is Hausdorff. Therefore, all three forms of compactness is the same.

Example 2.2 (Open Sets in Real Plane are Not Compact)

An open set in \mathbb{R}^2 is not compact. Take the open rectangle $R = (0, 1)^2 \subset \mathbb{R}^2$. There exists an infinite cover of R

$$R = \bigcup_{n=0}^{\infty} (0, 1) \times \left(0, \frac{2^{n+1} - 1}{2^{n+1}}\right) \quad (193)$$

that does not have a finite subcover.

Corollary 2.2 (Bolzano-Weierstrass Theorem)

Every bounded sequence in \mathbb{R}^n has a limit point, i.e. contains a convergent subsequence.

Proof. The fact that the infinite sequence is bounded means that there exists some closed subset $I \in \mathbb{R}^n$ that contains all point of the sequence. By definition I is compact, so by the Heine-Borel theorem, every cover of I has a finite subcover.

Now, assume that there exists an infinite sequence in I that is not convergent, i.e. has no limit point. Then, each point $x_i \in I$ would have a neighborhood $U(x_i)$ containing at most a finite number of points in the sequence. We can define I such that the union of the neighborhoods is a cover of I . That is,

$$I \subset \bigcup_{i=1}^{\infty} U(x_i) \quad (194)$$

However, since every $U(x_i)$ contains at most a finite number of points, we must have an infinite open neighborhoods to cover $I \implies$ we cannot have a finite subcover. This contradicts the fact that I is compact.

Proof. It suffices to prove that there exists a monotonic sequence within a bounded sequence (x_n) .

Theorem 2.9 (Connectedness)

A subset E of the real line \mathbb{R} is connected if and only if it has the following property: if $x \in E, y \in E$ and $x < z < y$, then $z \in E$.

Proof.

Definition 2.1 (Perfect Sets)

A set P is perfect if it is closed and all of its points are limit points of P . In other words, the limit points of P and P itself coincide.

$$P' = P \quad (195)$$

Theorem 2.10 ()

Let P be a nonempty perfect set in \mathbb{R}^k . Then P is uncountable.

2.1 Exercises**Exercise 2.1 (Math 531 Spring 2025, PS5.1)**

We know what it means for a metric space (X, d) to be compact. We say that it is sequentially compact if every sequence in (X, d) has a convergent subsequence. Prove that a metric space is compact if and only if it is sequentially compact.

Solution. We prove bidirectionally.

1. (\rightarrow). Assume that X is compact and let (x_n) be a sequence in X . For any $\epsilon > 0$, let

$$\mathcal{C}_\epsilon := \{B_\epsilon(x) \mid x \in X\} \quad (196)$$

be an open cover of open balls. Then there exists a finite subcover $\mathcal{F}_\epsilon \subset \mathcal{C}_\epsilon$. There is a countable sequence (x_n) , with each point in at least one open set in \mathcal{C} . By the pigeonhole principle, at least one open set must be hit infinitely many times, call this $B(x^*, \epsilon)$. Now consider for $\epsilon = \frac{1}{n}$ for $n \in \mathbb{N}$.

2. (\leftarrow).

Exercise 2.2 (Math 531 Spring 2025, PS5.2)

Give an example of a sequence of real numbers x_n for which

$$|x_{n+1} - x_n| \rightarrow 0 \quad (197)$$

as $n \rightarrow \infty$, but x_n is not convergent.

Solution. Consider the sequence

$$x_n = \sum_{i=1}^n \frac{1}{i} \quad (198)$$

It is the case that $x_{n+1} - x_n = 1/(n+1)$ which tends to 0, but this is a harmonic series which is not convergent.

Exercise 2.3 (Math 531 Spring 2025, PS5.3)

Let $\{x_n\}_{n=0}^{\infty}$ be a sequence of real numbers. Assume that

$$|x_{n+1} - x_n| \leq c|x_n - x_{n-1}| \quad (199)$$

for all $n \geq 1$, for some fixed $c < 1$. Prove that x_n is convergent. Hint: you may want to use the formula that you proved in a previous homework for $1 + c + c^2 + \cdots + c^N$.

Solution.

Exercise 2.4 (Math 531 Spring 2025, PS5.4)

Let x_n be a sequence of rational numbers defined recursively by:

$$x_0 = 1 \quad (200)$$

$$x_{n+1} = \frac{1}{x_n + 2} \quad (201)$$

when $n \geq 0$. The first few terms of this sequence are $1, \frac{1}{3}, \frac{3}{7}, \dots$. Prove that the sequence is convergent and find its limit.

Solution.

Exercise 2.5 (Math 531 Spring 2025, PS5.5)

As we know, every bounded sequence of real numbers has a convergence subsequence.

1. Let's say we have two sequences a_n and b_n that are bounded. Find a single sequence of indices $\{n_k\}$ so that *both* a_{n_k} and b_{n_k} are convergent. This is called a common convergent subsequence for a_n and b_n .
2. Show that for any finite number of bounded sequences of real numbers, we can find a common convergent subsequence.
3. Now suppose have a sequence of bounded sequences. Find a common convergent subsequence. What does this remind you of?

Solution.

Exercise 2.6 (Math 531 Spring 2025, PS5.6)

Given a sequence of real numbers x_n , we can define the sequence of its means by:

$$x_1, \frac{x_1 + x_2}{2}, \frac{x_1 + x_2 + x_3}{3}, \dots \quad (202)$$

Call the sequence of means y_n . Prove that if $x_n \rightarrow x$, then $y_n \rightarrow x$. Discuss the examples $x_n = (-1)^n$, $x_n = n$, $x_n = (-1)^n n$, and $x_n = (-1)^n \sqrt{n}$. The fourth example shows that it is possible to average out chaotic behavior (so long as it isn't focused in one direction). The third example shows that this is impossible if the system becomes too chaotic.

Solution.

Exercise 2.7 (Math 531 Spring 2025, PS4.1)

Let (X, d) be a metric space. Assume that K is compact and F is closed in (X, d) . Assume $K \cap F = \emptyset$. Prove that

$$\inf_{x \in F, y \in K} d(x, y) > 0. \quad (203)$$

Show by an example that this number could be zero if K is only assumed to be closed (rather than compact).

Solution. $K \cap F = \emptyset \implies K \subset F^c$ with F^c open. This means that for every $x \in K \subset F^c$, there exists a $r_x > 0$ s.t. $B(x, r_x) \subset F^c \iff B(x, r_x) \cap F = \emptyset$.

Now we take the covering $\{B(x, \frac{r_x}{2}) \mid x \in K\}$, and since K is compact there must be a finite subcovering, which we denote $C = \{B(x_i, \frac{r_i}{2}) \mid i = 1, \dots, n\}$. Denote $r^* = \min\{r_i\}$, which is positive since we take the minimum of a finite number of positive elements.

Now for any $x \in K$ and $y \in F$, x must be in some $B(x_i, \frac{r_i}{2}) \iff d(x, x_i) < \frac{r_i}{2} \iff -d(x, x_i) > -\frac{r_i}{2}$. With the same i , since $B(x_i, r_i)$ is disjoint from F , we have $d(x_i, y) \geq r_i$. Therefore, by the triangle inequality,

$$d(x, y) \geq d(x_i, y) - d(x_i, x) = r_i - \frac{r_i}{2} = \frac{r_i}{2} \geq \frac{r^*}{2} \quad (204)$$

and thus we have found a nontrivial lower bound.

Exercise 2.8 (Math 531 Spring 2025, PS4.2)

Consider \mathbb{R} with the usual metric. Find an open cover of \mathbb{Q} that does not cover \mathbb{R} .

Solution. $\mathbb{Q} = (-\infty, \sqrt{2}) \cup (\sqrt{2}, +\infty)$.

Exercise 2.9 (Math 531 Spring 2025, PS4.3)

\mathbb{R} is not compact with the usual metric since it is not bounded. Let us, however, define the following metric on \mathbb{R} :

$$d_*(x, y) = \frac{|x - y|}{(1 + |x|)(1 + |y|)}. \quad (205)$$

Verify that (\mathbb{R}, d_*) is a metric space. Prove that all subsets of (\mathbb{R}, d_*) are bounded. Show that \mathbb{R} still isn't compact with this metric. What is the problem?

Solution. We first verify metric.

1. Since $|x - y| \geq 0, 1 + |x| \geq 1, 1 + |y| \geq 1, d_*(x, y) \geq 0$. We also see that

$$d_*(x, y) = 0 \iff |x - y| = 0 \iff x = y \quad (206)$$

2. It is symmetric since

$$d_*(x, y) = \frac{|x - y|}{(1 + |x|)(1 + |y|)} = \frac{|y - x|}{(1 + |y|)(1 + |x|)} = d_*(y, x) \quad (207)$$

3. It satisfies triangle inequality since

$$d_*(x, y) + d_*(y, z) = \frac{|x - y|}{(1 + |x|)(1 + |y|)} + \frac{|y - z|}{(1 + |y|)(1 + |z|)} \quad (208)$$

$$= \frac{|x - y|(1 + |z|) + |y - z|(1 + |x|)}{(1 + |x|)(1 + |y|)(1 + |z|)} \quad (209)$$

$$= \frac{|x - y| + |x - y| \cdot |z| + |y - z| + |y - z| \cdot |x|}{(1 + |x|)(1 + |y|)(1 + |z|)} \quad (210)$$

$$\geq \frac{|x - z| + (|x| - |y|) \cdot |z| + (|y| - |z|) \cdot |x|}{(1 + |x|)(1 + |y|)(1 + |z|)} \quad (211)$$

$$= \frac{|x - z| + |x||y| - |y||z|}{(1 + |x|)(1 + |y|)(1 + |z|)} \quad (212)$$

$$= \frac{(1 + |y|)|x - z|}{(1 + |x|)(1 + |y|)(1 + |z|)} \quad (213)$$

$$= \frac{|x - z|}{(1 + |x|)(1 + |z|)} \quad (214)$$

$$= d_*(x, z) \quad (215)$$

where the inequality comes from $|x - y| + |y - z| \geq |x - z|$, $|x - y| \geq ||x| - |y|| \geq |x| - |y|$, and $|y - z| \geq ||y| - |z|| \geq |y| - |z|$.

This is bounded since for any $x, y \in \mathbb{R}$, we can show that the numerator is bounded by the denominator. Since both are positive from (1), it suffices to prove $|x - y|^2 \leq (1 + |x|)^2(1 + |y|)^2$.

$$(1 + |x|)^2(1 + |y|)^2 = (1 + 2|x| + |x|^2)(1 + 2|y| + |y|^2) \quad (216)$$

$$= 1 + 2|x| + 2|y| + 4|x||y| + |x|^2 + |y|^2 + 2|x||y|^2 + 2|y||x|^2 + |x|^2|y|^2 \quad (217)$$

$$\geq |x|^2 + |y|^2 + 2|x||y| \quad (218)$$

$$\geq |x|^2 + |y|^2 - 2xy \quad (219)$$

$$= |x - y|^2 \quad (220)$$

where the first inequality holds since all the terms in the expansion are nonnegative and the second holds since $|x||y| \geq xy$. Therefore, $d_*(x, y) \leq 1$. \mathbb{R} is still not compact since we can construct the set of open balls $B_r(0)$ around 0 w.r.t. d_* . Consider the cover

$$\mathcal{C} = \{B_{1-\frac{1}{n}}(0)\}_{n \in \mathbb{N}, n \geq 2} \quad (221)$$

Now assume that there is a finite subcover. Then there must be a maximum index $N \in \mathbb{N}$ in this cover. I claim that this does not cover \mathbb{R} . Consider the element $y = N - 1 \in \mathbb{R}$. The distance is

$$d_*(x, y) = \frac{|0 - (N - 1)|}{(1 + |0|)(1 + |N - 1|)} = \frac{N - 1}{N} = 1 - \frac{1}{N} \quad (222)$$

and so $y \notin \mathcal{C}$. Hence \mathcal{C} is not a cover of \mathbb{R} .

Exercise 2.10 (Math 531 Spring 2025, PS4.4)

Consider the set $X = \mathbb{R} \cup \{Gandalf\}$. Define a metric d_* on X by:

$$d_*(x, y) = \frac{|x - y|}{(1 + |x|)(1 + |y|)} \quad (223)$$

for $x, y \in \mathbb{R}$, while

$$d_*(Gandalf, x) = \frac{1}{1 + |x|}, \quad (224)$$

for all $x \in \mathbb{R}$. Verify that d_* is a metric on X (you don't have to do much for this, since you already did part of it in the previous problem). Prove that (X, d_*) is a compact metric space.

Solution.

Exercise 2.11 (Math 531 Spring 2025, PS4.5)

Let X be any set and endow it with the metric $d(x, y) = 1$ if $x \neq y$ and $d(x, x) = 0$. Check that this is a metric. Find all compact sets in (X, d) .

Solution. It trivially satisfies nonnegativity since it's either 0 or 1, and $d(x, x) = 0$. It is symmetric as well. As for triangle inequality this is trivial. All compact sets are finite sets.

Exercise 2.12 (Math 531 Spring 2025, PS3.1)

Determine for each of the following sets, whether or not it is countable. Justify your answers

1. The set of all functions $f : \{0, 1\} \rightarrow \mathbb{N}$.
2. The set B_n of all functions $f : \{1, \dots, n\} \rightarrow \mathbb{N}$
3. The set $C = \cup_{n \in \mathbb{N}} B_n$
4. The set of all functions $f : \mathbb{N} \rightarrow \{0, 1\}$.
5. The set of all functions $f : \mathbb{N} \rightarrow \{0, 1\}$ that are "eventually zero" (We say that f is eventually zero if there exists some $N \geq 1$ so that $f(n) = 0$ for all $n \geq N$.)
6. G the set of all functions $f : \mathbb{N} \rightarrow \mathbb{N}$ that are eventually constant.

Solution. Listed.

1. Countable since bijective to $\mathbb{N} \times \mathbb{N}$. We define the bijection as

$$(a_0, a_1) \in \mathbb{N} \times \mathbb{N} \mapsto f(i) = a_i \quad (225)$$

2. Countable since bijective to \mathbb{N}^n . We define the bijection as

$$(a_1, \dots, a_n) \in \mathbb{N}^n \mapsto f(i) = a_i \quad (226)$$

3. Countable since we proved that B_n is countable, and a countable union of countable sets are countable from Rudin Theorem 2.12.
4. Uncountable since we can create a bijection from the set of all sequences (a_i) of 0 or 1, which from Rudin Theorem 2.14 is uncountable.

$$(a_i)_{i \in \mathbb{N}} \mapsto f(i) = a_i \quad (227)$$

5. Countable. Call this set B , and call the set of functions f that have their final 1 at index k to be A_k . Then,

$$B = \cup_{k=1}^{\infty} A_k \quad (228)$$

where $A_0 = A_1 = 1$, and $|A_k| = 2^{k-1}$ for $k \geq 2$. Since B is the countable union of at most countable sets, B must be countable.

6. Countable. Call this set B . Let A_k be the set of functions that are eventually constant to value k . Let A_{ki} be the set of functions that are always k starting from index i (where i is the smallest element). Since everything is determined to be k at i and beyond, A_{ki} can be divided up into the

first $i - 1$ elements of any natural number, followed by a sequence of k 's. Therefore $|A_{ki}| \approx \mathbb{N}^{i-1}$, where \approx means equipotent, and so A_{ki} is countable. Therefore since countable unions of countable sets are countable,

$$A_k = \bigcup_{i=1}^{\infty} A_{ki} \text{ is countable} \implies B = \sum_{k=1}^{\infty} A_k \text{ is countable} \quad (229)$$

Exercise 2.13 (Math 531 Spring 2025, PS3.2)

Tell if the following subsets $A \subset \mathbb{R}$ (with the usual metric $d(x, y) = |x - y|$) are open or closed. Also, find (i) the limit points of A , (ii) the interior of A , (iii) \bar{A} .

1. $A = \mathbb{Q}$
2. $A = (0, 1]$
3. $A = \{1, \frac{1}{2}, \frac{1}{4}, \dots\}$
4. $A = \{0, 1, \frac{1}{2}, \frac{1}{4}, \dots\}$
5. $A = \mathbb{Z}$

Solution. Listed. We denote A' as the limit points of A and the interior as A° .

1. Not open nor closed. $A' = \mathbb{R}$, $A^\circ = \emptyset$. $\bar{A} = \mathbb{R}$.
2. Not open nor closed. $A' = [0, 1]$, $A^\circ = (0, 1)$. $\bar{A} = [0, 1]$.
3. Not open nor closed. $A' = \{0\}$, $A^\circ = \emptyset$. $\bar{A} = \{0\} \cup A$.
4. Closed. $A' = \{0\}$, $A^\circ = \emptyset$. $\bar{A} = A$.
5. Closed. $A' = \emptyset$. $A^\circ = \emptyset$. $\bar{A} = A$.

Exercise 2.14 (Math 531 Spring 2025, PS3.3)

Prove the following statements subsets A, B of a general metric space (X, d) .

- $\overline{A \cup B} = \bar{A} \cup \bar{B}$.
- Show by example that $\overline{A \cap B} \neq \bar{A} \cap \bar{B}$.

Solution. For the first part, we show bidirectionally.

1. $\overline{A \cup B} \subset \bar{A} \cup \bar{B}$. Let $x \in \overline{A \cup B}$. If $x \in A \cup B$, then it must be the case that either $x \in A \subset (A \cup A') = \bar{A}$ or $x \in B \subset (B \cup B') = \bar{B}$, which means $x \in \bar{A} \cup \bar{B}$. Now assume not. Then $x \in (A \cup B)'$. Therefore, for any $r > 0$, we know that $B(x, r) \cap (A \cup B) \neq \emptyset$. Now let us take a sequence $(r_n = \frac{1}{n})_{n \in \mathbb{N}}$, and for each r_n we have some element $x_n \in (A \cup B)$. Given that we have a countably infinite sequence of x_n , each which may be in A or B , by the pigeonhole principle either A or B must be hit infinitely many times. If $x_n \in A$ infinitely many times, then $x \in \bar{A}$, and analogous for B .
2. $\bar{A} \cup \bar{B} \subset \overline{A \cup B}$. WLOG let $x \in \bar{A}$. If $x \in A$, then $x \in (A \cup B) \subset \overline{A \cup B}$. If $x \notin A$, then $x \in A'$. Therefore for every $r > 0$, $B(x, r) \cap A \neq \emptyset$. But this means

$$\emptyset \neq (B(x, r) \cap A) \cup (B(x, r) \cap B) = B(x, r) \cap (A \cup B) \implies x \in (A \cup B)' \subset \overline{A \cup B} \quad (230)$$

For a counterexample, consider the sequences

$$A = (x_n) = \frac{1}{n} \quad B = (y_n) = -\frac{1}{n} \quad (231)$$

for $n \in \mathbb{N}$. $\bar{A} = A \cup \{0\}$, $\bar{B} = B \cup \{0\}$, and so $\bar{A} \cap \bar{B} = \{0\}$. However, $A \cap B = \emptyset \implies \overline{A \cap B} = \emptyset$.

Exercise 2.15 (Math 531 Spring 2025, PS3.4)

Consider the set of rationals in canonical form (such that numerator and denominator are relatively prime) with potential distance:

$$d_1\left(\frac{p_1}{q_1}, \frac{p_2}{q_2}\right) = |q_1 - q_2|. \quad (232)$$

Is this a metric? Prove that the following defines a metric

$$d_2\left(\frac{p_1}{q_1}, \frac{p_2}{q_2}\right) = |p_1 - p_2| + |q_1 - q_2|. \quad (233)$$

Solution. This is not a metric since

$$d_1\left(\frac{2}{1}, \frac{3}{1}\right) = 1 - 1 = 0 \quad (234)$$

when $2/1 \neq 3/1$. For d_2 , we show that it satisfies the three properties.

1. *Nonnegativity.* Since it is the sum of 2 absolute values which are norms and therefore nonnegative, it must be nonnegative by ordered field properties. We see that

$$\frac{p_1}{q_1} = \frac{p_2}{q_2} \iff p_1 = p_2 \text{ and } q_1 = q_2 \quad (235)$$

$$\iff |p_1 - p_2| = |q_1 - q_2| = 0 \quad (236)$$

$$\iff |p_1 - p_2| + |q_1 - q_2| = 0 \quad (237)$$

2. For symmetricity, note that

$$d_2\left(\frac{p_1}{q_1}, \frac{p_2}{q_2}\right) = |p_1 - p_2| + |q_1 - q_2| = |p_2 - p_1| + |q_2 - q_1| = d_2\left(\frac{p_2}{q_2}, \frac{p_1}{q_1}\right) \quad (238)$$

3. For triangle inequality, we see that for any $p_1/q_1, p_2/q_2, p_3/q_3$,

$$d_2\left(\frac{p_1}{q_1}, \frac{p_3}{q_3}\right) = |p_1 - p_3| + |q_1 - q_3| \quad (239)$$

$$= |(p_1 - p_2) + (p_2 - p_3)| + |(q_1 - q_2) + (q_2 - q_3)| \quad (240)$$

$$\leq |p_1 - p_2| + |p_2 - p_3| + |q_1 - q_2| + |q_2 - q_3| \quad (\text{subadditivity of norm})$$

$$= d_2\left(\frac{p_1}{q_1}, \frac{p_2}{q_2}\right) + d_2\left(\frac{p_2}{q_2}, \frac{p_3}{q_3}\right) \quad (241)$$

Exercise 2.16 (Math 531 Spring 2025, PS3.5)

Let $M = \{x_1, \dots, x_3\}$ be a set with three points. Describe the set of all metrics on M . What if M has four points?

Solution. If M has 3 points call them x_1, x_2, x_3 , then the metric is completely defined by the three values

$$d(x_1, x_2) = d(x_2, x_1) \quad (242)$$

$$d(x_2, x_3) = d(x_3, x_2) \quad (243)$$

$$d(x_3, x_1) = d(x_1, x_3) \quad (244)$$

where $d(x, x) = 0$. We must make sure that the triangle inequality satisfies for these 3 numbers. Therefore we can think of this as the set of all triangles in \mathbb{R}^2 (that are equivalent under translation

and rotation, but not permutation of points).

Similarly for 4 points, we can visualize the metrics as the set of all tetrahedra in \mathbb{R}^3 (since each face is a triangle, and therefore for any three points the triangle inequality is guaranteed to be satisfied), equivalent under translation and rotation, but not permutation of the 4 points.

Exercise 2.17 (Math 531 Spring 2025, PS3.6)

Let P be a polynomial of degree $n \geq 1$. Prove that if $P(0) = 0$, then $P(x) = xQ(x)$, for some polynomial Q of degree $n - 1$. Deduce that if $P(a) = 0$, then we can write $P(x) = (x - a)Q(x)$ for some Q of degree $n - 1$.

Solution. A n th degree polynomial will have the form

$$p(x) = \sum_{i=0}^n c_i x^i \quad (245)$$

Since $p(0) = c_0 = 0 \implies c_0 = 0$. This means that

$$p(x) = \sum_{i=1}^n c_i x^i = x \sum_{i=0}^{n-1} c_{i+1} x^i \text{ where } Q(x) = \sum_{i=0}^{n-1} c_{i+1} x^i \quad (246)$$

If $p(a) = 0$, we can construct $f(x) = p(x + a)$, where f is a polynomial since the expansion does not increase its degree. Since $f(x) = p(a) = 0$, by above f can be factorized $f(x) = xg(x)$ for some $(n-1)$ th degree polynomial g , and by substitution this means that $p(x) = f(x - a) = (x - a)g(x - a)$.

Exercise 2.18 (Math 531 Spring 2025, PS3.7)

Consider all polynomials $P : \mathbb{R} \rightarrow \mathbb{R}$ of degree less than or equal to n . Call this set \mathcal{P}_n . Let's define potential distances on \mathcal{P}_n .

$$d_1(p, q) = |p(0) - q(0)|. \quad (247)$$

Show this defines a distance on \mathcal{P}_0 but not on \mathcal{P}_n for $n \geq 1$. Now consider

$$d_N(p, q) = \sum_{j=0}^N |p(j) - q(j)| \quad (248)$$

Show that this defines a distance on \mathcal{P}_n , for every $n \leq N$. What does the solution say about polynomials of degree N ?

Solution. If $n = 0$, \mathcal{P}_n is a set of constant functions P , where each constant function P is determined completely by its value at any point, e.g. 0. We check the properties.

1. $d_1(p, q) \geq 0$ since we take the norm at the end. We can see that

$$d_1(p, q) = 0 \iff |p(0) - q(0)| \quad (249)$$

$$\iff p(0) = q(0) \quad (250)$$

$$\iff p = q \quad (251)$$

2. It is clearly symmetric.

$$d_1(p, q) = |p(0) - q(0)| = |q(0) - p(0)| = d_1(q, p) \quad (252)$$

3. It satisfies the triangle inequality by subadditivity of the norm.

$$d_1(p, r) = |p(0) - r(0)| \quad (253)$$

$$= |(p(0) - q(0)) + (q(0) - r(0))| \quad (254)$$

$$\leq |p(0) - q(0)| + |q(0) - r(0)| \quad (255)$$

$$= d_1(p, q) + d_1(q, r) \quad (256)$$

It doesn't satisfy for P_n because consider $p(x) = x$ and $q(x) = x^2$. They are not the same function but $d_1(p, q) = |p(0) - q(0)| = 0$. For d_N defined on \mathcal{P}_n for $n \leq N$, we verify the properties.

1. This is the sum of norms, so it must be nonnegative. Now we see that if $p = q$, then $p(x) = q(x) \implies |p(x) - q(x)| = 0 \implies d_N(p, q) = 0$. For the other way around, suppose $d_N(p, q) = 0$. Then from problem 3.8, we are solving the linear equation $0 = Vb - Vc$, where b, c are the vectors representing the coefficients of p, q , and V is the Vandermonde matrix with $a_i = i$. By linearity, this is equivalent to solving $0 = V(b - c)$, and since we showed that V is invertible (since a_i 's are distinct), V has a trivial kernel and therefore $b - c = 0 \iff b = c \implies p = q$.
2. Symmetricity is trivial.

$$d_N(p, q) = \sum_{j=0}^N |p(j) - q(j)| = \sum_{j=0}^N |q(j) - p(j)| = d_N(q, p) \quad (257)$$

3. For triangle inequality,

$$d_N(p, r) = \sum_{j=0}^N |p(j) - r(j)| \quad (258)$$

$$= \sum_{j=0}^N |(p(j) - q(j)) + (q(j) - r(j))| \quad (259)$$

$$\leq \sum_{j=0}^N |p(j) - q(j)| + |q(j) - r(j)| \quad (260)$$

$$= \sum_{j=0}^N |p(j) - q(j)| + \sum_{j=0}^N |q(j) - r(j)| \quad (261)$$

$$= d_N(p, q) + d_N(q, r) \quad (262)$$

This shows that we need to “sample” more points from higher-degree polynomials to get the metric as they are higher-dimensional.

Exercise 2.19 (Math 531 Spring 2025, PS3.8)

Given distinct numbers a_0, \dots, a_N and numbers b_0, \dots, b_N , prove that there exists a polynomial P of degree N with the property that

$$P(a_i) = b_i, \quad (263)$$

for $0 \leq i \leq N$. The most direct way to solve this problem, in my view, is to write the system equations you are trying to solve as a linear system for the coefficients of P . This will give you some matrix M that depends on the numbers a_0, \dots, a_N . The key is to show that the determinant of this matrix is non-zero. It turns out that the determinant of this matrix is equal to

$$\prod_{0 \leq i < j \leq N} (a_i - a_j), \quad (264)$$

up to a potential – sign depending on how you defined M . Prove this and deduce the result.

Solution. We can write the system of equations using the Vandermonde matrix $V \in \mathbb{R}^{(N+1) \times (N+1)}$ and c is the vector of coefficients of P .

$$b = Vc \iff \begin{bmatrix} b_0 \\ b_1 \\ \vdots \\ b_N \end{bmatrix} = \begin{bmatrix} 1 & a_0 & a_0^2 & \dots & a_0^N \\ 1 & a_1 & a_1^2 & \dots & a_1^N \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & a_N & a_N^2 & \dots & a_N^N \end{bmatrix} \begin{bmatrix} c_0 \\ c_1 \\ \vdots \\ c_N \end{bmatrix} \quad (265)$$

To calculate the determinant of V , we prove using induction. Clearly for $N = 1$ we have

$$\det \begin{pmatrix} 1 & a_0 \\ 1 & a_1 \end{pmatrix} = a_1 - a_0 \quad (266)$$

Now assume that this formula holds for some $N - 1 \in \mathbb{N}$. Then for N , we can take V and subtract a_0 times the i th column from the $(i + 1)$ st column. This gives us

$$V = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 1 & a_1 - a_0 & a_1^2 - a_0 a_1 & \dots & a_1^N - a_0 a_1^{N-1} \\ 1 & a_2 - a_0 & a_2^2 - a_0 a_2 & \dots & a_2^N - a_0 a_2^{N-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & a_N - a_0 & a_N^2 - a_0 a_N & \dots & a_N^N - a_0 a_N^{N-1} \end{bmatrix} \quad (267)$$

When calculating the determinant, we can perform the cofactor expansion by the first row, and then for each i th row factor out $(a_i - a_0)$ to get

$$\det V = \prod_{j=1}^N (a_j - a_0) \det \begin{bmatrix} 1 & a_1 & \dots & a_1^{N-1} \\ 1 & a_2 & \dots & a_2^{N-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & a_N & \dots & a_N^{N-1} \end{bmatrix} \quad (268)$$

which is the $(N - 1) \times (N - 1)$ Vandermonde matrix. Therefore, we can apply our inductive hypothesis to get

$$\det V = \prod_{j=1}^N (a_j - a_0) \prod_{1 \leq i < j \leq N} (a_j - a_i) = \prod_{0 \leq i < j \leq N} (a_j - a_i) \quad (269)$$

Note that this has a 0 determinant iff $a_i = a_j$ for some $i \neq j$. Therefore since a_i 's are distinct, it must be nonzero. Therefore, this matrix is nonsingular, i.e. invertible, and we can solve the matrix equation to get

$$c = V^{-1}b \quad (270)$$

which from linear algebra is guaranteed to exist and is unique.

Exercise 2.20 (Rudin 2.1)

Prove that the empty set is a subset of every set.

Solution. It must suffice that if $x \in \emptyset$, then $x \in A$ for any arbitrary set A . This is vacuously true, since the initial condition is never met.

Exercise 2.21 ()

Show that the empty function $f : \emptyset \rightarrow X$, where X is an arbitrary set, is always injective. If $X = \emptyset$, then f is bijective.

Solution. Given distinct $x, y \in \emptyset$, $f(x) \neq f(y)$ is vacuously true, but if $X \neq \emptyset$, then there exists a $w \in X$ with no preimage. If $X = \emptyset$, then the statement for all $w \in X$, there exists an $x \in \emptyset$ s.t. $f(x) = w$ is vacuously true.

Exercise 2.22 (Rudin 2.2)

A complex number z is said to be algebraic if there are integers a_0, a_1, \dots, a_n , not all zero, such that

$$a_0 z^n + a_1 z^{n-1} + \dots + a_{n-1} z + a_n = 0. \quad (271)$$

Prove that the set of all algebraic complex numbers is countable. Hint: For every positive integer N there are only finitely many equations with

$$n + |a_0| + |a_1| + \dots + |a_n| = N \quad (272)$$

Solution. Consider all polynomials s.t. $n + \sum_{i=0}^n |a_i| = N$. There is only a finite number of them, and each polynomial has at most n distinct complex roots. So this set is finite, an unioning over all $N \in \mathbb{N}$ gives an at most countable set of roots.

Exercise 2.23 (Rudin 2.3)

Prove there exists real numbers which are not algebraic.

Solution. From the previous exercise, if there were no real numbers which are not algebraic, then every real number is algebraic. This contradicts the fact that the set of all complex numbers is countable.

Exercise 2.24 (Rudin 2.4)

Is the set of all irrational real numbers countable?

Solution. No. Assume that it is countable. We have \mathbb{Q} countable. Then, by assumption, we must have $\mathbb{R} = \mathbb{Q} \cup \mathbb{Q}^c$ be the union of countable sets, which must be countable, contradicting the fact that it is uncountable.

Exercise 2.25 (Rudin 2.5)

Construct a bounded set of real numbers which exactly 3 limit points.

Solution. We can construct the union of 3 sequences that converge onto the limit points 0, 1, 2.

$$\left\{\frac{1}{n}\right\}_{n \in \mathbb{N}} \cup \left\{\frac{1}{n} + 1\right\}_{n \in \mathbb{N}} \cup \left\{\frac{1}{n} + 2\right\}_{n \in \mathbb{N}} \quad (273)$$

Exercise 2.26 ()

Prove that the union of the limit points of sets is equal to the limit points of the union of the sets.

$$\bigcup_{k=1}^m A'_k = \left(\bigcup_{k=1}^m A_k \right)' \quad (274)$$

Solution.

Exercise 2.27 (Rudin 2.6)

Let E' be the set of all limit points of a set E . Prove that E' is closed. Prove that E and \bar{E} have the same limit points. (Recall that $\bar{E} = E \cup E'$). Do E and E' always have the same limit points?

Solution. Listed.

1. Let x be a limit point of E' . Then, for every $\epsilon > 0$, $U = B_\epsilon(x) \cap E' \neq \emptyset$. Take a $y \in U$. Since $y \in B_\epsilon(x)$, which is open, we can construct an open ball $B_\delta(y) \subset B_\epsilon(x)$. Since $y \in E'$, $B_\delta(y)$ must contain elements of E , which means that $B_\epsilon(x)$ must also contain elements of E , and so x is a limit point of $E \implies x \in E'$ and E' is closed.
2. To prove that $E' \subset \bar{E}'$, we know that if $x \in E'$, then for every $\epsilon > 0$, there exists a $B_\epsilon(x)$ that has a nontrivial intersection with E which means that it has a nontrivial intersection with $E \cup E'$. To prove that $\bar{E}' \subset E'$, we know that if $y \in \bar{E}'$, then for every $\delta > 0$ there exists a $B_\delta(x)$ that has a nontrivial intersection with \bar{E} . If $B_\delta(x)$ intersects E then we are done. If $B_\delta(x)$ intersects E' , then we can find a $y \in E' \cap B_\delta(x)$. Since $B_\delta(x)$ is open, we can construct $B_\epsilon(y) \subset B_\delta(x)$ and since $y \in E'$, we know that $B_\epsilon(y)$ contains an element of E , which means that $B_\delta(x)$ contains an element of E . Therefore, $E' = \bar{E}'$.
3. No. Consider the set $E = \{1/n\}_{n \in \mathbb{N}}$. $E' = \{0\}$, but $E'' = \emptyset$.

Exercise 2.28 (Rudin 2.7)

Let A_1, A_2, \dots be subsets of a metric space.

1. If $B_n = \bigcup_{i=1}^n A_i$, prove that $\bar{B}_n = \bigcup_{i=1}^n \bar{A}_i$ for $n = 1, 2, 3, \dots$
2. If $B = \bigcup_{i=1}^\infty A_i$, prove that $\bar{B} \supset \bigcup_{i=1}^\infty \bar{A}_i$.

Solution. Listed.

1. We will prove that $\bar{B}_n \subseteq \bigcup_{i=1}^n \bar{A}_i$ and $\bigcup_{i=1}^n \bar{A}_i \subseteq \bar{B}_n$. If $x \in B_n$, then $x \in \bigcup_{i=1}^n A_i$. Therefore, assume that $x \in B'_n$. Then for every $\epsilon > 0$, there exists a $B_\epsilon(x)$ s.t.

$$B_\epsilon(x) \cap B_n \neq \emptyset \implies B_\epsilon(x) \cap \left(\bigcup_{i=1}^n A_i \right) \neq \emptyset \quad (275)$$

This means that there exists some $i = i(\epsilon)$, a function of ϵ , s.t. $B_\epsilon(x) \cap A_i \neq \emptyset$. However, this i may change if we unfix ϵ . We have so far proved that just for one $\epsilon > 0$ there exists an i . Now if we take a sequence of $\epsilon = 1, \frac{1}{2}, \frac{1}{3}, \dots$, we have a sequence of $i(\epsilon)$'s living in $\{1, \dots, n\}$. By the pigeonhole principle, there must be at least one i that is hit infinitely many times, and so we can choose this i , that works for all $\epsilon > 0 \implies x \in A'_i \subseteq \bigcup_{i=1}^n \bar{A}_i$. If $x \in \bigcup_{i=1}^n \bar{A}_i$, then there exists an \bar{A}_i s.t. $x \in \bar{A}_i$. If $x \in A_i$, then we are done. If $x \in A'_i$, then for every $\epsilon > 0$, there exists a $B_\epsilon(x)$ s.t.

$$B_\epsilon(x) \cap A_i \neq \emptyset \implies B_\epsilon(x) \cap \left(\bigcup_{i=1}^n A_i \right) \neq \emptyset \quad (276)$$

and so $x \in B'_n \subset \overline{B_n}$.

2. $x \in \bigcup_{i=1}^{\infty} \overline{A_i} \implies x \in \overline{A_i}$ for some i . If $x \in A_i$, then $x \in B$ and we are done. If $x \in A'_i$, then for every $\epsilon > 0$ there exists $B_\epsilon(x)$ s.t.

$$B_\epsilon^\circ(x) \cap A_i \neq \emptyset \implies B_\epsilon^\circ(x) \cap \left(\bigcup_{i=1}^{\infty} A_i \right) \neq \emptyset \quad (277)$$

and so $B_\epsilon^\circ(x) \cap B \neq \emptyset \implies x \in B' \subset \overline{B}$.

Exercise 2.29 (Rudin 2.8)

Is every point of every open set $E \subset \mathbb{R}^2$ a limit point of E ? Answer the same question for closed sets in \mathbb{R}^2 .

Solution. Yes for open. Given any $x \in U$ open, there always exists an $\epsilon > 0$ s.t.

$$B_\epsilon^\circ(x) \subset B_\epsilon(x) \subset U \quad (278)$$

and so $B_\epsilon^\circ(x)$ has a nontrivial intersection with U . If U is closed, then no. Note that for closed U , we have that every limit point is in U , but not every point in U is a limit point. Consider the isolated point $U = \{x\}$. x is not a limit point of U .

Exercise 2.30 (Rudin 2.9)

Let E° denote the set of all interior points of E in X . Prove the following:

- E° is always open.
- E is open if and only if $E^\circ = E$.
- If $G \subseteq E$ and G is open, then $G \subset E^\circ$.
- Prove that the complement of E° is the closure of the complement of E .
- Do E and \bar{E} always have the same interiors?
- Do E and E° always have the same closures?

Solution. Listed.

- We assume that E° is not open (this does not mean that E° is necessarily closed!). That is, there exists an $x \in E^\circ$ s.t. we can't construct an open ball $B_\epsilon(x) \subseteq E^\circ$. Since $x \in E^\circ \subset E$, by definition of an interior point we can construct a $B_\epsilon(x) \subset E$. But from our assumption $B_\epsilon(x) \not\subset E^\circ$. We choose a $y \in B_\epsilon(x) \setminus E^\circ$. Since $B_\epsilon(x)$ is open, there exists a $\delta > 0$ s.t.

$$B_\delta(y) \subset B_\epsilon(x) \subset E \quad (279)$$

But the fact that we can construct an open ball around y means that $y \in E^\circ$, leading to a contradiction.

- If E is open, then by definition $E \subset E^\circ$. Now $E^\circ \subset E$ holds for all sets since E° must be composed of points from E . If $E = E^\circ$, then for every $x \in E$, $x \in E^\circ$, so by definition there exists an $\epsilon > 0$ s.t. $B_\epsilon(x) \subset E$, which means that E is open.
- Let $x \in G$ open. Then there exists an $\epsilon > 0$ s.t. $B_\epsilon(x) \subset G$, and so $B_\epsilon(x) \subset E$. Since we can always construct an open ball around x contained within E , $x \in E^\circ$ and $G \subset E^\circ$.
- $((E^\circ)^c \subset \overline{E^c})$ If $x \in (E^\circ)^c$, then there exists no $\epsilon > 0$ s.t. $B_\epsilon(x) \subset E$. Then, for any $\epsilon > 0$, $B_\epsilon(x) \not\subset E \implies B_\epsilon(x) \cap E^c \neq \emptyset \implies x \in E^c \subset \overline{E^c}$. $(\overline{E^c} \subset (E^\circ)^c)$ If $x \in \overline{E^c}$, then $x \in E^c$ or $x \in E^{c'}$. If $x \in E^c$, note $E^\circ \subset E \implies (E^\circ)^c \supset E^c \implies x \in (E^\circ)^c$. If $x \in E^{c'}$, then for all $\epsilon > 0$ $B_\epsilon(x) \cap E^c \neq \emptyset \implies B_\epsilon(x) \not\subset E \implies x \in E^\circ$.

5. No. Consider the rationals $\mathbb{Q} \subset \mathbb{R}$. $\mathbb{Q}^\circ = \emptyset$ but $\overline{\mathbb{Q}}^\circ = \mathbb{R}^\circ = \mathbb{R}$. It is true and straightforward to prove that $E^\circ \subset \overline{E}^\circ$. Let $x \in E^\circ$. Then there exists an $\epsilon > 0$ s.t. $B_\epsilon(x) \subset E \implies B_\epsilon(x) \subset \overline{E} \implies x \in \overline{E}^\circ$.
6. No. Consider $\mathbb{Q} \subset \mathbb{R}$. Then $\overline{\mathbb{Q}} = \mathbb{R}$ and $\overline{\mathbb{Q}}^\circ = \overline{\emptyset} = \emptyset$.

Exercise 2.31 (Rudin 2.10)

Let X be an infinite set. For $p \in X$ and $q \in X$, define

$$d(p, q) = \begin{cases} 1 & \text{if } p \neq q \\ 0 & \text{if } p = q \end{cases} \quad (280)$$

Prove that this is a metric. Which subsets of the resulting metric space are open? Which are closed? Which are compact?

Solution. This is a metric since clearly it satisfies symmetry and the fact that $d(p, p) = 0$. The triangle inequality

$$d(p, r) \leq d(p, q) + d(q, r) \quad (281)$$

is trivially satisfied if $p = r$, and if $p \neq r$, then either $p \neq q$ or $q \neq r$, and so the RHS ≥ 1 . An open ϵ -ball around $x \in X$ is either X , when $\epsilon > 1$, or $\{x\}$ when $\epsilon \leq 1$. Therefore

Exercise 2.32 (Rudin 2.11)

For $x \in \mathbb{R}$ and $y \in \mathbb{R}$, define

$$\begin{aligned} d_1(x, y) &= (x - y)^2 \\ d_2(x, y) &= \sqrt{|x - y|} \\ d_3(x, y) &= |x^2 - y^2| \\ d_4(x, y) &= |x - 2y| \\ d_5(x, y) &= \frac{|x - y|}{1 + |x - y|} \end{aligned}$$

Determine, for each of these, whether it is a metric or not.

Solution. Listed. Positive semidefiniteness and symmetry are easy to check.

1. The triangle inequality gives

$$d_1(x, z) \leq d_1(x, y) + d_1(y, z) \iff (x - z)^2 \leq (x - y)^2 + (y - z)^2 \quad (282)$$

$$\iff 0 \leq (x - y)(y - z) \quad (283)$$

which is not satisfied if $x < y < z$, so this is not a valid metric.

2. The triangle inequality gives $\sqrt{|x - z|} \leq \sqrt{|x - y|} + \sqrt{|y - z|}$, and since both sides are positive this inequality is equivalent to squaring both sides to get

$$|x - z| \leq |x - y| + |y - z| + 2\sqrt{|x - y||y - z|} \quad (284)$$

which is true since $|x - z| \leq |x - y| + |y - z|$ of the Euclidean distance satisfies the triangle inequality and $0 \leq \sqrt{|x - y||y - z|}$.

3. This does not satisfy triangle inequality, as taking 0, 1, 2 gives

$$d_3(0, 2) = 4 > 1 + 1 = d_3(0, 1) + d_3(1, 2) \quad (285)$$

4. This does not satisfy symmetry.

5. For simplicity, let us set $A = |x - y|, B = |y - z|, C = |x - z|$. Then, we get

$$\frac{C}{1+C} \leq \frac{A}{1+A} + \frac{B}{1+B} \iff C \leq A + B + 2AB + ABC \quad (286)$$

where $C \leq A + B$ is true by triangle inequality of Euclidean distance, $0 \leq AB$, and $0 \leq ABC$. Intuitively, we want a metric that doesn't "blow up" the distance between x and y . More precisely, we want a valid metric $d(x, y)$ to be $O(|x - y|)$. Having something like a quadratic growth rate $(x - y)^2$ will blow the distance $d(x, z)$ up too much overpowering the individual $d(x, y) + d(y, z)$.

Exercise 2.33 (Rudin 2.12)

Let $K \subset \mathbb{R}$ consist of 0 and the numbers $1/n$ for $n = 1, 2, 3, \dots$. Prove that K is compact directly from the definition (without using the Heine-Borel theorem).

Solution. Every open cover of K must have an open set G s.t. $0 \in G$. Since G is open, there exists an open neighborhood $B_\epsilon(0) \subset G$ around 0. By the Archimidean principle, there exists an $N \in \mathbb{N}$ s.t.

$$\epsilon N > 1 \implies \epsilon > \frac{1}{N} \quad (287)$$

and so, $B_\epsilon(0)$ contains all points $\{1/n\}$ for $n > N$. For the rest of the points $1, 1/2, \dots, 1/N$, we can simply construct a finite cover over each of them, hence getting a finite cover.

Exercise 2.34 (Rudin 2.13)

Construct a compact set of real numbers whose limit points form a countable set.

Solution. Consider the set

$$E = \left\{ \left(\frac{1}{10} \right)^n + \left(\frac{1}{10} \right)^{n+k} : n \in \{0\} \cup \mathbb{N}, k \in \mathbb{N} \right\} \cup \{0\} \quad (288)$$

This is clearly bounded by 0 and 1.1. Let us represent the elements of this set by (n, k) . We can show that

$$(n_1, k_1) > (n_2, k_2) \quad (289)$$

if $n_1 < n_2$ or $n_1 = n_2$ and $k_1 < k_2$. Therefore, to prove closedness, we must prove that every limit point is a point in E . We can do this by proving that a point not in E cannot be a limit point. Choose any $x \notin E$. Then, due to the ordering, we can see that there exists a (n, k) s.t.

$$A = \left(\frac{1}{10} \right)^n + \left(\frac{1}{10} \right)^{n+k} < x < \left(\frac{1}{10} \right)^n + \left(\frac{1}{10} \right)^{n+k+1} = B \quad (290)$$

and so we can take $\epsilon = \min\{x - A, B - x\}$ and show that $B_\epsilon(x)$ does not contain A nor B , and so has an empty intersection with E . Therefore, it cannot be a limit point of E and is closed. Since E is bounded and closed in \mathbb{R} , it is compact. Its limit points contain $1, 0.1, 0.01, \dots, 0$ (simply by fixing n and letting $k \rightarrow \infty$, and so E' is infinite. We have just shown that since E is closed, $E' \subset E$. But E is countable, so E' is countable.

Exercise 2.35 (Rudin 2.14)

Given an example of an open cover of the segment $(0, 1)$ which has no finite subcover.

Solution. Consider

$$(0, 1/2) \cup \left(\bigcup_{i=1}^{\infty} \left[1 - \frac{1}{2^i}, 1 - \frac{1}{2^{i+1}} \right) \right) \quad (291)$$

Exercise 2.36 (Rudin 2.15)

Solution.

Exercise 2.37 (Rudin 2.16)

Regard \mathbb{Q} , the set of all rational numbers, as a metric space, with $d(p, q) = |p - q|$. Let E be the set of all $p \in \mathbb{Q}$ s.t. $2 < p^2 < 3$. Show that E is closed and bounded in \mathbb{Q} but is not compact. Is E open in \mathbb{Q} ?

Solution. E is clearly bounded by 0 and 2 since $0^2 < 2 < p^2 < 3 < 2^2$. It is closed and we can show this by showing that E^c is open. Let $x \in E^c$. Then, $x^2 < 2$ or $x^2 > 3$.

1. $x^2 < 2 \iff -\sqrt{2} < x < \sqrt{2}$. Now let $\epsilon = \min\{\sqrt{2} - x, x + \sqrt{2}\} > 0$. Then by the Archimidean property there exists a $n \in \mathbb{N}$ s.t. $0 < \frac{1}{n} < \epsilon$. Therefore, the image of $B_{1/n}(x) \subset \mathbb{Q}$ will map onto $(0, 2)$.
2. $x^2 > 3 \iff x > \sqrt{3}$ or $x < -\sqrt{3}$. If $x > \sqrt{3}$, then by AP there exists a $n \in \mathbb{N}$ s.t. $x - \frac{1}{n} > \sqrt{3} \implies (x - \frac{1}{n})^2 > 3$. If $x < -\sqrt{3}$, then by AP there exist $n \in \mathbb{N}$ s.t. $x + \frac{1}{n} < -\sqrt{3} \implies (x + \frac{1}{n})^2 > 3$. Either way, the image of $B_{1/n}(x)$ will map within E^c .

It is not compact because E is not closed in \mathbb{R} . The limit points of E in \mathbb{R} is $[\sqrt{2}, \sqrt{3}] \cup [-\sqrt{3}, -\sqrt{2}]$, which contains irrationals and is clearly not a subset of E . Since it is not closed in \mathbb{R} , it is not compact in \mathbb{R} , and it is not compact in $\mathbb{Q} \subset \mathbb{R}$. It is open because

$$E = ((\sqrt{2}, \sqrt{3}) \cup (-\sqrt{3}, \sqrt{2})) \cup \mathbb{Q} \subset \mathbb{R} \quad (292)$$

which is the union of open $(\sqrt{2}, \sqrt{3}) \cup (-\sqrt{3}, \sqrt{2})$ and subset $\mathbb{Q} \subset \mathbb{R}$, and so it is open.

Exercise 2.38 (Rudin 2.17)

Let E be the set of all $x \in [0, 1]$ whose decimal expansion consists of only the digits 4 and 7. Is E countable? Is E dense in $[0, 1]$? Is E compact? Is E perfect?

Solution.

Exercise 2.39 (Rudin 2.18)

Is there a nonempty perfect set in \mathbb{R} which contains no rational number?

Solution.

Exercise 2.40 (Rudin 2.19)

Listed.

1. If A and B are disjoint closed sets in some metric space X , prove that they are separated.
2. Prove the same for disjoint open sets.
3. Fix $p \in X$, $\delta > 0$, define A to be the set of all $q \in X$ for which $d(p, q) < \delta$. Define B similarly, with $>$ in place of $<$. Prove that A and B are separated.
4. Prove that every connected metric space with at least two points is uncountable.

Solution. Listed.

1. This is trivial with the fact that the closure of the closure of A is the closure of A .
2. Let A, B be open. We wish to show that if $x \in A'$, then $x \notin B$. Assume $x \in B$. Then there exists $\epsilon > 0$ s.t. $B_\epsilon(x) \subset B$. But $B \cap A = \emptyset \implies B_\epsilon(x) \cap A = \emptyset$ and so $x \notin A'$, which is a contradiction.
3. Clearly, $A \cap B = \emptyset$. Not let $x \in A \implies$ there exists $\epsilon > 0$ s.t. $B_\epsilon(x) \subset A \implies B_\epsilon(x) \cap B = \emptyset \implies x \in B'$. We can prove similarly to show that $x \in B \implies x \notin A'$.
4. Assume X is countable (solution is very similar for finite). Then, we can enumerate a $X = \{x_i\}_{i=1}^\infty$. We wish to show that X can be decomposed into the union of an open ball and the interior of its complement as shown in (3). We fix $p \in X$. Then, we take the set $D = \{d(p, x)\}_{x \neq p} \subset \mathbb{R}$. Since D is a countable subset of \mathbb{R} , there must exist some $\alpha > 0$ s.t. $\alpha \notin D$. This α partitions the distances into two sets, and we can define

$$X = \{q \in X \mid d(p, q) < \alpha\} \cup \{q \in X \mid d(p, q) > \alpha\} \quad (293)$$

and by (3), these two sets are separated, which means that X is not connected, leading to a contradiction.

Exercise 2.41 (Rudin 2.20)

Are closures and interiors of connected sets always connected? Look at subsets of \mathbb{R}^2 .

Solution. The interiors are not always connected. Consider the two closed balls $\overline{B_1((1,0))}$ and $\overline{B_1((-1,0))}$ as subsets of \mathbb{R}^2 . They are connected but their interiors, which are the two open balls, are not connected.

As for closures, they are always connected. Let W be connected. Then for any partition $A \cup B = W$, $\overline{A} \cap B \neq \emptyset$ WLOG. Consider $\overline{W} = W \cup W'$ and take any partition $\overline{W} = C \cup D$. Then, label $A = C \cap W, A^* = C \cap W', B = D \cap W, B^* = D \cap W'$. This implies that $C = A \cup A^*, D = B \cup B^*$, and $A \cup B = W$ (which is connected). Then, we can show that

$$\begin{aligned} \overline{C} \cap D &= (\overline{A \cup A^*}) \cap D = (\overline{A} \cup \overline{A^*}) \cap D = (\overline{A} \cap D) \cup (\overline{A^*} \cap D) \\ &= (\overline{A} \cap B) \cup (\overline{A} \cap B^*) \cup (\overline{A^*} \cap D) \end{aligned}$$

which cannot be empty since by connectedness of W , $\overline{A} \cap B \neq \emptyset$. Therefore, \overline{W} is connected.

Exercise 2.42 (Rudin 2.21)

Let A and B be separated subsets of some \mathbb{R}^k . Suppose $a \in A, b \in B$ and define

$$p(t) = (1-t)a + tb \quad (294)$$

for $t \in \mathbb{R}$. Put $A_0 = p^{-1}(A), B_0 = p^{-1}(B)$.

1. Prove that A_0 and B_0 are separated subsets of \mathbb{R} .
2. Prove that there exists a $t_0 \in (0, 1)$ s.t. $p(t_0) \notin A \cup B$.

3. Prove that every convex subset of \mathbb{R}^k is connected.

Solution.

Exercise 2.43 (Rudin 2.22)

A metric space is called *separable* if it contains a countable dense subset. Show that \mathbb{R}^k is separable.

Solution. Consider the set $\mathbb{Q}^k \subset \mathbb{R}^k$. It is a finite Cartesian product (and hence, a countable union) of countable \mathbb{Q} , and so it is countable. \mathbb{Q}^k is dense in \mathbb{R}^k since given any $x \in \mathbb{R}^k$, we claim x is a limit point of \mathbb{Q}^k . Given any $\epsilon > 0$, we can construct $B_\epsilon^\circ(x)$. For each coordinate x_i , by density of rationals in \mathbb{R} we can choose a $q_i \in \mathbb{Q}$ s.t. $0 < d(x_i, q_i) < \epsilon/k$. Then, using triangle inequality, we can take the distances between each coordinate changed from x_i to q_i . Let q^k be the vector x with the components x_1, \dots, x_k changed to q_1, \dots, q_k , respectively.

$$d(x, q) = d(x, q^1) + d(q^1, q^2) + \dots + d(q^{k-1}, q_k) < \frac{\epsilon}{k} + \dots + \frac{\epsilon}{k} = \epsilon \quad (295)$$

and so $q \in B_\epsilon^\circ(x)$. Hence the intersection of \mathbb{Q}^k and $B_\epsilon^\circ(x)$ for any $\epsilon > 0$ is nontrivial, so x is a limit point of \mathbb{Q}^k .

Exercise 2.44 (Rudin 2.23)

A collection $\{V_\alpha\}$ of open subsets of X is said to be a *base* for X if the following is true: For every $x \in X$ and every open set $G \subset X$ such that $x \in G$, we have $x \in V_\alpha \subset G$ for some α . In other words, every open set in X is the union of a subcollection of $\{V_\alpha\}$. Prove that every separable metric space has a countable base.

Solution. Since X is separable it contains a countable dense subset, call it S . Then for every $x \in S$, we can look at the set of all open balls with center x and rational radii, call it \mathcal{B} . Then \mathcal{B} is countable. Now consider an open set U . By definition, for every $x \in U$, there exists an $\epsilon > 0$ s.t. $B_\epsilon(x) \subset U$. By AP, we can find a $n \in \mathbb{N}$ s.t. $0 < \frac{1}{n} < \epsilon$, and therefore we can find an open ball $B \in \mathcal{B}$ s.t. $B(x) \subset U$. We claim that

$$W := \bigcup_{x \in U} B(x) = U \quad (296)$$

If $x \in U$, then by construction it is contained in $B(x) \subset \bigcup_{x \in U} B(x)$, and so $U \subset W$. If $x \in W$, then it is in $B(x)$, which is fully contained in U and so $W \subset U$. Therefore every open set can be constructed by a countable union of open balls in countable \mathcal{B} .

Exercise 2.45 (Rudin 2.24)

Let X be a metric space in which every infinite subset has a limit point. Prove that X is separable.

Solution. We fix $\delta > 0$. Choose $x_1 \in X$. Then choose $x_2 \in X$ s.t. $d(x_1, x_2) \geq \delta$, and keep doing this until we choose $x_{j+1} \in X$ s.t. $d(x_{j+1}, x_i) \geq \delta$ for all $i \in 1, \dots, j$.

1. We claim that this must stop after a finite number of steps. Assume it doesn't. Then by assumption $V = \{x_i\}_{i=1}^\infty$ should have a limit point in X , denote it x . Choose $\frac{\delta}{2} > 0$. Then, $B_{\delta/2}^\circ(x) \cap V \neq \emptyset$. This intersection can only have one point since if it had two x', x'' , then since

both are in $B_{\delta/2}(x)$, then

$$d(x', x'') \leq d(x', x) + d(x, x'') \leq \frac{\delta}{2} + \frac{\delta}{2} = \delta \quad (297)$$

and since they are both in V , then $d(x', x'') \geq \delta$, which is a contradiction. Since there is a finite number of points in $B_{\delta/2}(x)$ of V , x cannot be a limit point. So this must terminate at some finite $J < \infty$.

2. Denote $W = \{x_i\}_{i=1}^J$. Then, $\mathcal{B}_\delta = \{B_\delta(x) \mid x \in W\}$ must cover X , since if it didn't, there would exist a $y \in X$ s.t. $d(y, x) \geq \delta$ for all $x \in W$, and we can add another element in W .
3. Consider $\delta = 1, 1/2, 1/3, \dots$ and construct the same cover

$$\mathcal{B}_k = \{B_{1/k}(x_{ki}) \mid i = 1, \dots, J_k\} \quad (298)$$

which is finite. Therefore, $\mathcal{B} = \cup_{k=1}^{\infty} \mathcal{B}_k$ must be countable.

4. We claim that countable $\{x_{ki}\}_{k,i}$ is dense. Consider any $x \in X$. For every $\epsilon > 0$, we can find an arbitrarily large $n \in \mathbb{N}$ s.t. $0 < \frac{1}{n} < \epsilon$. Since \mathcal{B}_n is an open cover, there must exist some x_{ni} s.t. $x \in B_{1/n}(x_{ni})$, which by symmetry implies that $x_{ni} \in B_{1/n}(x) \subset B_\epsilon(x)$. Therefore, there always exists an x_{ni} in every $B_\epsilon(x)$, and so $B_\epsilon(x) \cap \{x_{ki}\} \neq \emptyset \implies x$ is a limit point of $\{x_{ki}\}$ and so it is dense.

Exercise 2.46 (Rudin 2.25)

Prove that every compact metric space K has a countable base, and that K is therefore separable.

Solution. For every $n \in \mathbb{N}$, let us consider an open covering $\mathcal{F}_n := \{B_{1/n}(x_n) \mid x_n \in K\}$. Since K is compact, it has a finite subcovering

$$\mathcal{G}_n := \{B_{1/n}(x_{ni}) \mid i = 1, \dots, k(n)\} \quad (299)$$

Now consider the union $\mathcal{G} = \cup_{n=1}^{\infty} \mathcal{G}_n$, which is countable. We claim that \mathcal{G} is a base. Consider any open set U . Then for every $x \in U$, we want to show that x is contained in a $B_{1/n}(x_{ni}) \subset U$. Since U is open, there exists a $\epsilon > 0$ s.t. $B_\epsilon(x) \subset U$. Now by AP, there exists a $n \in \mathbb{N}$ s.t. $0 < \frac{1}{n} < \frac{\epsilon}{2}$. Therefore $B_{1/n}(x) \subset B_\epsilon(x)$. Since \mathcal{G} is an open covering, there must exist some $B_{1/n}(x_{ni})$ that contains x . Now we wish to show that $B_{1/n}(x_{ni})$ is fully contained in U . Let $y \in B_{1/n}(x_{ni})$. Then, by triangle inequality,

$$d(y, x) = d(y, x_{ni}) + d(x_{ni}, x) < \frac{1}{n} + \frac{1}{n} < \epsilon \quad (300)$$

and therefore $x \in B_{1/n}(x_{ni}) \subset B_\epsilon(x)$. Therefore, for every $x \in U$, we can construct an open ball of \mathcal{G} containing x and contained in U , proving that this is a base.

We claim that the set of all $\mathcal{P} = \{x_{ni}\}_{n,i}$ forms a countable dense subset. This is clearly countable since \mathcal{G} is countable. We must prove that the closure of $\mathcal{P} = K$. Let $x \in K$. Given any $\epsilon > 0$, we wish to show that $B_\epsilon(x) \cap \mathcal{P} \neq \emptyset$. Since $B_\epsilon(x)$ is open, it can be covered by a subcollection of \mathcal{G} , and so their centers must be in $B_\epsilon(x)$, proving that $B_\epsilon(x) \cap \mathcal{P} \neq \emptyset$. Therefore, x is a limit point of \mathcal{P} .

Exercise 2.47 (Rudin 2.26)

Solution.

Exercise 2.48 (Rudin 2.27)*Solution.***Exercise 2.49 (Rudin 2.28)***Solution.***Exercise 2.50 (Rudin 2.29)**

Prove that every open set in \mathbb{R} is the union of an at most countable collection of disjoint segments.

Solution. Let $U \subset \mathbb{R}$ be open. Then for all $x \in U$ there exists $\epsilon > 0$ s.t. $(x - \epsilon, x + \epsilon) \subset U$. Now since \mathbb{R} is separable (by exercise Rudin 2.22), it has a countable dense subset \mathbb{Q} . Consider all segments of rational radius and rational centers

$$\mathcal{B} = \{(q - p, q + p) \subset \mathbb{R} \mid q, p \in \mathbb{Q}\} \quad (301)$$

This is clearly countable. We claim that every open U can be expressed as the union of a subset of \mathcal{B} . Now by AP, there exists $n \in \mathbb{N}$ s.t. $0 < \frac{1}{n} < \frac{\epsilon}{2}$, so for all $x \in U$, there exists $n \in \mathbb{N}$ s.t. $(x - \frac{1}{n}, x + \frac{1}{n}) \subset U$. Now since \mathbb{Q} is dense in \mathbb{R} , $x \in \mathbb{Q}' \implies (x - \frac{1}{n}, x + \frac{1}{n}) \cap \mathbb{Q} \neq \emptyset$. Say r is in this intersection. Then, by symmetry of metric, $x \in (r - \frac{1}{n}, r + \frac{1}{n})$. Therefore, for all $x \in U$, we have found an open ball in \mathcal{B} that contains x . Now, we must show that this actually is fully contained in U . This is easy, since if $y \in B_{1/n}(r)$, then

$$d(y, x) \leq d(y, r) + d(r, x) \leq \frac{1}{n} + \frac{1}{n} < \epsilon \quad (302)$$

and so $B_{1/n}(r)$ is completely contained in the ϵ -ball around x , which is a subset of U . So for all $x \in U$, we found an open set $U_x \in \mathcal{B}$ covering x and fully contained in U , which means that $\cup_{x \in U} U_x = U$. Now for some intervals $B_1, B_2 \in \mathcal{B}$, if $B_1 \cap B_2 \neq \emptyset$, take their union, which is another segment, and keep doing this until $B_i \cap B_j \neq \emptyset$ for all i, j . The cardinality of this new pruned set will be less than or equal to \mathcal{B} , which is countable, and so this must be at most countable.

3 Sequences in Euclidean Space

We assume that you know what sequences are, and what it means for them to converge in \mathbb{R}^n . Let's primarily focus on sequences in \mathbb{R} . Note that there are many way in which a sequence can be divergent.

1. Increasing/decreasing indefinitely
2. Oscillating between two constant values
3. Oscillating between a value tending to $+\infty$ and a value tending to $-\infty$
4. Many other classes of divergence

Definition 3.1 (Sequence Tending to Infinity)

The sequence $\{x_n\}$ **tends to positive infinity** if for each number c there exists $N \in \mathbb{N}$ such that $x_n > c$ for all $n > N$. It is denoted

$$x_n \rightarrow +\infty \text{ or } \lim_{n \rightarrow \infty} x_n = +\infty \quad (303)$$

We define sequences that **tend to negative infinity** similarly. And $\{x_n\}$ **tends to infinity** if for each c there exists $N \in \mathbb{N}$ such that $|x_n| > c$ for all $n > N$, which is written

$$x_n \rightarrow \infty \quad (304)$$

Note that

$$x_n \rightarrow +\infty \text{ or } x_n \rightarrow -\infty \implies x_n \rightarrow \infty \quad (305)$$

but the converse is not necessarily true. The simple example is the sequence $x_n = (-1)^n n$. Also, it is important to know that a sequence may be unbounded and yet not tend to $+\infty$, $-\infty$, or ∞ .

Example 3.1 (Unbounded Sequence that Doesn't tend to ∞)

The sequence $x_n = n^{(-1)^n}$ is divergent yet does not tend to positive infinity, negative infinity, nor infinity.

Definition 3.2 (Monotonic Sequences)

Let X be an ordered set. (x_n) is

1. **strictly increasing** if $x_{n+1} > x_n$ for all n .
2. **strictly decreasing** if $x_{n+1} < x_n$ for all n .
3. **increasing (nondecreasing)** if $x_{n+1} \geq x_n$ for all n .
4. **decreasing (nonincreasing)** if $x_{n+1} \leq x_n$ for all n .

Sequences of these types are called **monotonic**.

The properties of a general metric space give us some general conditions to determine whether a sequence converges in \mathbb{R}^n . To add to our toolbox in determining convergence, we will focus on sufficient conditions for convergence in \mathbb{R} , which has the additional properties of being an ordered field. These additional structures unlock a whole new suite of theorems in convergence. Why do we want to focus on just real-valued, i.e. numerical, sequences? First is that since \mathbb{R}^n is constructed as the product topology of \mathbb{R} , we can prove a lot about continuity of functions in \mathbb{R}^n by proving limits in \mathbb{R} , and letting the construction of the product topology do the rest. Second, the codomain of many natural structures such as inner products, norms, and measures lie in the reals, and we often need to prove convergence of these values.

3.1 Convergence Theorems from Completeness and Order

So far, in order to show that a sequence is convergent, we must identify a real number first and then show using the ϵ - δ definition that it converges. This might be overkill in a case where we just want to prove that a sequence converges, but we don't care what it converges to. The following theorems are direct consequences of the Cauchy-completeness and the least upper bound property of the reals.

Theorem 3.1 (Cauchy-Convergence Criterion)

A Cauchy sequence in \mathbb{R} converges.

Proof. This is by definition true, and to see how other forms of completeness implies Cauchy convergence, see above.

The second result is an immediate consequence of Dedekind completeness, which is equivalent to Cauchy completeness in the reals.

Theorem 3.2 (Convergence Criterion for Monotonic Sequences)

Let $(x_n)_n$ be a sequence.

1. If (x_n) is monotonically increasing/decreasing and bounded above/below in \mathbb{R} , then it converges.
2. If (x_n) is monotonically increasing/decreasing in $\mathbb{R} \cup \{\pm\infty\}$, then it converges.

Proof. It satisfies to prove the first case, as the second case can be done similarly without much difficulty. Let $x_n \leq x_{n+1}$. Then the set $\{x_n\}$ is bounded above in \mathbb{R} , which has the least upper bound property, and so there exists a least upper bound x . We claim that the sequence converges to x . For every $\epsilon > 0$, since it is least, there exists at least one $x_N \in (x - \epsilon, x)$. By monotonicity, this means that $x_n \in (x - \epsilon, x)$ for all $n \geq N$, and so the sequence converges to x .

We are able to see how both Cauchy and Dedekind completeness of the reals define convergence in \mathbb{R} . Now let's squeeze a bit more out of the total ordering to gain some properties of convergence and divergence.

Theorem 3.3 (Preservation of Ordering Between Sequences and Limits)

Given convergent sequences (x_n) and (y_n) , if

$$\lim_{n \rightarrow \infty} x_n < \lim_{n \rightarrow \infty} y_n \quad (306)$$

then there exists an index $N \in \mathbb{N}$ such that $x_n < y_n$ for all $n > N$.

Proof. Given $x_n \rightarrow x, y_n \rightarrow y$ and $x < y$ ($\in \mathbb{R}$), then for every $\epsilon > 0$, there exists $N_1, N_2 \in \mathbb{N}$ s.t. $d(x, x_n) < \epsilon$ for all $n > N_1$ and $d(y, y_n) < \epsilon$ for all $n > N_2$. Setting $N = \max\{N_1, N_2\}$, we can say the same for all $n > N$. We choose $\epsilon = \frac{y-x}{2} > 0$. Then, there exists $N \in \mathbb{N}$ s.t. $x_n \in (x - \epsilon, x + \epsilon)$ and $y_n \in (y - \epsilon, y + \epsilon)$ for all $n > N$. Therefore, if $a \in (x - \epsilon, x + \epsilon)$ and $b \in (y - \epsilon, y + \epsilon)$, then

$$a < \sup B_\epsilon(x) = x + \epsilon = y - \epsilon = \inf B_\epsilon(y) < b \quad (307)$$

which implies that $x_n < y_n$ for all $n > N$.

Theorem 3.4 (Squeeze Theorem for Sequences)

Given sequences $(x_n), (y_n), (z_n)$ such that

$$x_n \leq y_n \leq z_n \quad (308)$$

for all $n > N$, if $\{x_n\}$ and $\{z_n\}$ both converge to the same limit, then the sequence $\{y_n\}$ also converges to that limit. That is,

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} z_n = A \implies \lim_{n \rightarrow \infty} y_n = A \quad (309)$$

Proof. We first prove that if there exists a $N \in \mathbb{N}$ s.t. $a_n \leq b_n$ for all $n > N$, then $\lim_{n \rightarrow \infty} a_n = a \leq b = \lim_{n \rightarrow \infty} b_n$. Assume this weren't true, that $a > b$. Then for $\epsilon = \frac{a-b}{2} > 0$, there must exist $M \in \mathbb{N}$ s.t. $a_n \in (a - \epsilon, a + \epsilon)$ and $b_n \in (b - \epsilon, b + \epsilon)$ for all $n > M$. But

$$b_n < \sup(b - \epsilon, b + \epsilon) = b + \epsilon = a - \epsilon = \inf(a - \epsilon, a + \epsilon) < a_n \quad (310)$$

which contradicts $a_n \leq b_n$. Therefore, $a \leq b$. Therefore, we can use this to get

$$A = \lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n \leq \lim_{n \rightarrow \infty} z_n = A \implies \lim_{n \rightarrow \infty} y_n = A \quad (311)$$

Theorem 3.5 (Unbounded Sequence has Subsequence Tending to Infinity)

If (x_n) is not bounded above then it has a subsequence $x_{n_k} \rightarrow +\infty$.

Proof. We can construct such a subsequence.

Therefore, we can construct a subsequential limit to $\pm\infty$ if (x_n) is not bounded. If it is bounded, then by the Bolzano-Weierstrass theorem it contains a convergent subsequence. Therefore, we have the following.

Corollary 3.1 ()

From each sequence of real numbers there exists either a convergent subsequence or a subsequence tending to infinity.

Proof. If it's bounded, then Bolzano-Weierstrass. If it's not bounded, then use previous theorem.

Example 3.2 ()

We claim that

$$\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1 \quad (312)$$

3.2 Arithmetic**Theorem 3.6 (Arithmetic on Limits)**

Given that $(x_n), (y_n)$ are numerical sequences with $y_n \neq 0$ for all n , and let

$$\lim_{n \rightarrow \infty} x_n = x, \quad \lim_{n \rightarrow \infty} y_n = y \neq 0 \quad (313)$$

then,

$$\lim_{n \rightarrow \infty} (x_n + y_n) = x + y \quad (314)$$

$$\lim_{n \rightarrow \infty} (cx_n) = cx \quad (315)$$

$$\lim_{n \rightarrow \infty} (x_n \cdot y_n) = x \cdot y \quad (316)$$

$$\lim_{n \rightarrow \infty} \frac{x_n}{y_n} = \frac{x}{y} \quad (317)$$

It immediately follows that the set of all convergent sequences in $\mathbb{R}^{\mathbb{N}}$ is a subspace of $\mathbb{R}^{\mathbb{N}}$.

Proof. For every $\epsilon > 0$, there exists $N_1, N_2 \in \mathbb{N}$ such that

$$|x_n - x| < \epsilon \text{ for all } n > N_1 \quad (318)$$

$$|y_n - y| < \epsilon \text{ for all } n > N_2 \quad (319)$$

Therefore, for a given ϵ , we wish to prove that there exists a N such that for all $n > N$,

$$1. |(x_n + y_n) - (x + y)| < \epsilon \quad (320)$$

$$2. |cx_n - cx| < \epsilon \quad (321)$$

$$3. |(x_n y_n) - (xy)| < \epsilon \quad (322)$$

$$4. \left| \frac{x_n}{y_n} - \frac{x}{y} \right| < \epsilon \quad (323)$$

1. By the triangle inequality, we can see that

$$|(x_n + y_n) - (x + y)| = |x_n - x| + |y_n - y| \quad (324)$$

Since we can choose the error between x_n and x for $n > N_1$, and y_n and y for $n > N_2$ as small as we want, we set it to $\epsilon/2$. Then, we have

$$|(x_n + y_n) - (x + y)| = |x_n - x| + |y_n - y| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \quad (325)$$

for all $n > N = \max\{N_1, N_2\}$. Therefore, for a given ϵ , there exists an N such that

$$|(x_n + y_n) - (x + y)| < \epsilon \text{ for all } n > N \quad (326)$$

2. This proof is easy. For a given ϵ , we choose the error to be $\frac{\epsilon}{c}$.

$$|x_n - x| < \frac{\epsilon}{c} \text{ for all } n > N_1 \quad (327)$$

Then, there exists natural number N_1 such that

$$|cx_n - cx| < c|x_n - x| = c \frac{\epsilon}{c} = \epsilon \text{ for all } n > N_1 \quad (328)$$

3. We first observe that since the limit of $\{y_n\}$ exists, it must be bounded by a value, say y . That is,

$$|y_n| < Y \text{ for all } n \in \mathbb{N} \quad (329)$$

Then, we see that

$$|x_n y_n - xy| = |(x_n y_n - xy_n) + (xy_n - xy)| \quad (330)$$

$$< |x_n y_n - xy_n| + |xy_n - xy| \quad (331)$$

$$= |y_n||x_n - x| + |x||y_n - y| \quad (332)$$

Suppose $\epsilon > 0$ is given. Then, we can set the error bounds freely; there exists $N_1, N_2 \in \mathbb{N}$ such that

$$|x_n - x| < \frac{\epsilon}{2Y} \text{ for all } n > N_1 \quad (333)$$

$$|y_n - y| < \frac{\epsilon}{2|x|} \text{ for all } n > N_2 \quad (334)$$

Then, we can see that

$$|x_n y_n - xy| \leq |y_n||x_n - x| + |x||y_n - y| < Y \cdot \frac{\epsilon}{2Y} + |x| \frac{\epsilon}{2|x|} = \epsilon \quad (335)$$

for all $n > N = \max\{N_1, N_2\}$.

4. We use the estimate

$$\left| \frac{x}{y} - \frac{x_n}{y_n} \right| = \frac{|x_n||y_n - y| + |y_n||x_n - x|}{y_n^2} \cdot \frac{1}{1 - \delta(y_n)}, \quad \delta(y_n) = \frac{|y_n - y|}{|y_n|} \quad (336)$$

For a given $\epsilon > 0$, we find natural numbers N_1, N_2 such that

$$|x_n - x| < \min \left\{ 1, \frac{\epsilon|y|}{8} \right\} \text{ for all } n > N_1 \quad (337)$$

$$|y_n - y| < \min \left\{ \frac{|y|}{4}, \frac{\epsilon y^2}{16(|x| + 1)} \right\} \text{ for all } n > N_2 \quad (338)$$

From this we can deduce that

$$|x_n| = |x_n - x + x| < |x_n - x| + |x| < |x| + 1 \quad (339)$$

and

$$|y| = |y_n + y - y_n| < |y_n| + |y - y_n| \quad (340)$$

$$\implies |y_n| > |y| - |y_n - y| > |y| - \frac{|y|}{4} > \frac{|y|}{2} \quad (341)$$

$$\implies \frac{1}{|y_n|} < \frac{2}{|y|} \quad (342)$$

$$\implies 0 < \delta(y_n) = \frac{|y_n - y|}{|y_n|} < \frac{|y|/4}{|y|/2} = \frac{1}{2} \quad (343)$$

$$\implies 1 - \delta(y_n) > \frac{1}{2} \quad (344)$$

$$\implies 0 < \frac{1}{1 - \delta(y_n)} < 2 \quad (345)$$

So, we can substitute

$$|x_n| \cdot \frac{1}{y_n^2} \cdot |y_n - y| < (|x| + 1) \cdot \frac{4}{y^2} \cdot \frac{\epsilon \cdot y^2}{16(|x| + 1)} = \frac{\epsilon}{4} \quad (346)$$

$$\left| \frac{1}{y_n} \right| \cdot |x_n - x| < \frac{2}{|y|} \cdot \frac{\epsilon|y|}{8} = \frac{\epsilon}{4} \quad (347)$$

into the final equation to get

$$\left| \frac{x}{y} - \frac{x_n}{y_n} \right| < \epsilon \text{ for all } n > N = \max\{N_1, N_2\} \quad (348)$$

Example 3.3 ()

We claim that

$$\lim_{n \rightarrow \infty} \frac{n}{q^n} = 0 \text{ if } q > 1 \quad (349)$$

Since $x_n = \frac{n}{q^n} \implies x_{n+1} = \frac{n+1}{nq} x_n$ for $n \in \mathbb{N}$. Since

$$\lim_{n \rightarrow \infty} \frac{n+1}{nq} = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right) \frac{1}{q} = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right) \cdot \lim_{n \rightarrow \infty} \frac{1}{q} = 1 \cdot \frac{1}{q} = \frac{1}{q} < 1 \quad (350)$$

there exists an index N such that $\frac{n+1}{nq} < 1$ for $n > N$. Thus, we have

$$x_n > x_{n+1} = x_n \cdot \frac{n+1}{nq} \text{ for } n > N \quad (351)$$

which means that the sequence will be monotonically decreasing from index N on. The terms of the sequence

$$x_{N+1} > x_{N+2} > x_{N+3} > \dots \quad (352)$$

are positive (bounded below) and are monotonically decreasing, so it must have a limit.

Finding the actual limit is easy. Let $x = \lim_{n \rightarrow \infty} x_n$. It follows from the relation $x_{n+1} = \frac{n+1}{nq} x_n$ that

$$x = \lim_{n \rightarrow \infty} (x_{n+1}) = \lim_{n \rightarrow \infty} \left(\frac{n+1}{nq} x_n \right) = \lim_{n \rightarrow \infty} \frac{n+1}{nq} \cdot \lim_{n \rightarrow \infty} x_n = \frac{1}{q} x \quad (353)$$

which implies that $(1 - \frac{1}{q}) = 0 \implies x = 0$.

Definition 3.3 (Cauchy Product)

The Cauchy Product is the direct convolution of two sequences.

Definition 3.4 (Recursive Sequence)

Sometimes, a sequence may be defined **recursively**, where the n th term contains a combination of the $n - 1$ terms before it.

3.3 Limsup and Liminf

The superior and inferior limits represent some sort of "bound" on the sequence in the long run. That is, on the long run, the terms of the sequence (x_n) cannot be greater than its superior limit and cannot be less than its inferior limit. With this interpretation, the following definition should be clear.

Definition 3.5 (Inferior, Superior Limits)


The **superior/inferior limit** of a sequence (x_n) is defined in the equivalent ways.

1. Given that E is the set of all partial limits, the limsup/liminf is the supremum/infimum of E .

$$\limsup_{n \rightarrow \infty} x_n := \sup\{E\} \quad \liminf_{n \rightarrow \infty} x_n := \inf\{E\} \quad (354)$$

2. The limsup/liminf is the limit of the sequence of supremums/infimums of the elements up to k .

$$\limsup_{n \rightarrow \infty} x_n := \lim_{n \rightarrow \infty} \sup_{k \geq n} x_k \quad \liminf_{n \rightarrow \infty} x_n := \lim_{n \rightarrow \infty} \inf_{k \geq n} x_k \quad (355)$$



Example 3.4 (Computing Limsup and Liminf)

We give some basic examples.

1. Let $x_n = (-1)^n$. Then $E = \{-1, +1\}$ and

$$\limsup_{n \rightarrow \infty} x_n = 1, \quad \liminf_{n \rightarrow \infty} x_n = -1 \quad (356)$$

2. Let $x_n = (-1)^n / [1 + (1/n)]$. Then

$$\limsup_{n \rightarrow \infty} x_n = 1, \quad \liminf_{n \rightarrow \infty} x_n = -1 \quad (357)$$

Let's give two warnings. First, limsup and liminf do *not* behave like limits under addition and multiplication. That is,

$$\limsup x_n + \limsup y_n \neq \limsup x_n + y_n \quad (358)$$

Example 3.5 (Counterexamples of Arithmetic Consistency of Limit superior)

Consider $(x_n) = (-1)^n$ and $y_n = (-1)^{n+1}$. Then

$$\limsup x_n = \limsup y_n = 1, \quad \liminf x_n = \liminf y_n = -1 \quad (359)$$

But $(x_n + y_n) = 0$, so

$$\limsup x_n + y_n = \liminf x_n + y_n = 0 \quad (360)$$

Second, note that even though we are talking about subsequential limits, the limsup and liminf are *not* subsequential limits! It is the supremum of subsequential limits E , which may or may not be in E .

Example 3.6 (Limsup that is not attained by any subsequential limit)

This should be a sequence not in \mathbb{R} .

However, in \mathbb{R} , it turns out that the limsup and liminf are both contained in E , so we are fine.

Lemma 3.1 ()

If (x_n) is a sequence in \mathbb{R} , then

1. the limsup is indeed a subsequential limit, i.e. $\limsup x_n \in E$.
2. If $x > \limsup x_n$, then $\exists N \in \mathbb{N}$ s.t. $n \geq N \implies x_n < x$.

Proof. For the first claim, there are two cases to consider. If (x_n) is unbounded from above, then $\exists(x_{n_k})$ such that $x_{n_k} \rightarrow +\infty \implies \infty = \limsup x_n \in E$. If (x_n) is bounded from above, then the subsequential limits of (x_n) are either in (x_n) or they are limit points of x_n . This implies that the set E consists of points either in $\{x_n\}$ or are limit points of the set $\{x_n\} \implies \sup E$ is in E since it's a limit point.

For the second claim, if there are infinitely many terms of the sequence larger than x , then we could find a subsequence (x_{n_k}) with $x_{n_k} > x$ for all k . Therefore (x_n) has a subsequential limit which must be $\geq x$. Every subsequential limit of (x_{n_k}) is also a subsequential limit of (x_n) . This contradicts $\limsup x_n = \sup E$.

Theorem 3.7 (Requirements of Partial Limits for Limit to Exist)

Here are two results in which we can use partial limits to determine if a sequence has a limit or not.

1. A sequence has a limit or tends to $\pm\infty$ if and only if its inferior and superior limits are the same.

$$\limsup x_n = \liminf x_n = x \implies \lim_{n \rightarrow +\infty} x_n = x \quad (361)$$

2. A sequence converges if and only if every subsequence of it converges.

Proof. For (1), we pick $x + \epsilon > x$. Then every term past some N_1 must be less than $x + \epsilon$. By the same logic, we have N_2 for $x - \epsilon < x$. So take $N = \max\{N_1, N_2\}$, which is contained in the ϵ -ball around x .

Theorem 3.8 (Ordering on Subsequential Limits)

If $s_n \leq t_n$ for $n \geq N$, where N is fixed, then

$$\begin{aligned} \liminf_{n \rightarrow \infty} s_n &\leq \liminf_{n \rightarrow \infty} t_n \\ \limsup_{n \rightarrow \infty} s_n &\leq \limsup_{n \rightarrow \infty} t_n \end{aligned}$$

Example 3.7 ()

We claim

$$\lim_{n \rightarrow \infty} n^{1/n} = 1 \quad (362)$$

We can consider $x_n = n^{1/n} - 1$ and want to show that $x_n \rightarrow 0$. We have $x_n \geq 0$. If $n > 1$, then $n = (x_n + 1)^n \geq x_n^2 \cdot \frac{n(n-1)}{2}$ from the binomial theorem. This means that

$$x_n^2 \leq \frac{2}{n-1} \implies 0 \leq x_n \leq \sqrt{\frac{2}{n-1}} \rightarrow 0 \quad (363)$$

And so by the squeeze theorem, $x_n \rightarrow 0$.

Example 3.8 ()

If $x > 1, \alpha \in \mathbb{R}$, then

$$\lim_{n \rightarrow +\infty} \frac{n^\alpha}{x^n} = 0 \quad (364)$$

3.4 Convergence Tests for Real Series**Definition 3.6 (Series over \mathbb{R})**

Given a sequence of real numbers (x_n) , the **series (of partial sums)** is the sequence

$$(s_n) = \sum_{k=1}^n x_k \quad (365)$$

The **sum of the series** is the limit of (s_n) . Usually we define (s_n) implicitly and use the summation notation.

$$\sum_{n=1}^{\infty} x_n := \lim_{n \rightarrow \infty} s_n \quad (366)$$

1. If the sequence (s_n) converges to s , the series is **convergent**, written

$$\sum x_n < +\infty \quad (367)$$

2. If the sequence does not converge, it is **divergent**.
3. If the series of partial sums of $(|x_n|)$ converges, then it is said to be **absolutely convergent**.^a

$$\sum_{n=1}^{\infty} |x_n| \quad (368)$$

^aClearly, every absolutely convergent series is convergent because $|\sum_{n=1}^{\infty} a_n| \leq \sum_{n=1}^{\infty} |a_n|$. This is sort of the infinite analogue of the triangle inequality.

We must reiterate a few warnings here. Note that the series $\sum x_n$ is simply notation and should *not* be treated as an “infinite sum.” Such a thing does not exist for algebraic structures which have finary operations. More specifically, given a series, we cannot in general split nor combine series, and we cannot reindex nor rearrange (an infinite number of) terms. However, we can manipulate each term for a fixed index.

Example 3.9 (Disasters of Reindexing and Rearranging)

Let us take the series $\sum 0$. We clearly know that the corresponding sequence of partial sums $0, 0, \dots$ is

convergent to 0. But if we do this series of steps.

$$\begin{aligned}
 \sum_{n=1}^{\infty} 0 &= \sum_{n=1}^{\infty} n - n && \text{(Can manipulate terms)} \\
 &= \sum_{n=1}^{\infty} n - \sum_{n=1}^{\infty} n && \text{(Cannot split series)} \\
 &= 1 + \sum_{n=2}^{\infty} n - \sum_{n=1}^{\infty} n && \text{(Can take 1st term out)} \\
 &= 1 + \sum_{n=1}^{\infty} (n+1) - \sum_{n=1}^{\infty} n && \text{(Cannot reindexing)} \\
 &= 1 + \sum_{n=1}^{\infty} (n+1) - n && \text{(Cannot combine series)} \\
 &= 1 + \sum_{n=1}^{\infty} 1 && \text{(Can manipulate terms)} \\
 &= 1 + \infty = +\infty && (369)
 \end{aligned}$$

The wrong steps show that the series is divergent.

We have seen the consequences of these mistakes that beginners make and are often on popular media. However, note that we can always do splitting, combining, reindexing, and rearranging for *finite sums*, which are algebraically defined. Later on, we will show that some of these operations are allowed for series that we know are convergent.

Lemma 3.2 (All Series in Extended Positive Reals Converge)

All series in $\mathbb{R}_{\geq 0} \cup \{+\infty\}$ converge.

$$\sum_{n=1}^{\infty} x_n \in [0, +\infty] \quad (370)$$

Proof.

Since the convergence of a series is equivalent to convergence of its sequence of partial sums, applying the Cauchy convergence criterion to the sequence $\{s_n\}$ leads to the following theorem.

Theorem 3.9 (Cauchy Convergence Criterion for Series)

The series $a_1 + \dots + a_n + \dots$ converges if and only if for every $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that for all $m \geq n > N$,

$$|a_n + \dots + a_m| < \epsilon \quad (371)$$

Corollary 3.2 (nth Term Test)

A necessary (but not sufficient) condition for convergence of the series $a_1 + \dots + a_n + \dots$ is that the terms tend to 0 as $n \rightarrow \infty$. That is, it is necessary that

$$\lim_{n \rightarrow \infty} a_n = 0 \quad (372)$$

Proof. It suffices to set $m = n$ in the Cauchy convergence criterion. This would mean that for every $\epsilon > 0$ there exists a $N \in \mathbb{N}$ such that

$$|a_n| = |a_n - 0| < \epsilon \text{ for all } n > N \quad (373)$$

which, by definition, means that $\{a_n\}$ converges to 0.

Nothing so far is really suprising here. The Cauchy convergence criterion really just follows from the definition of Cauchy completeness, and the n th term test is pretty trivial. The way that we will build up convergence tests is by proving some special cases of convergence and then using the direct comparison test to then classify further series.

Example 3.10 (Telescoping Series)

A **telescoping series** is a series in which the partial sums can cancel out. An example is the series of partial sums of the sequence $(x_n) = \frac{1}{n(n+1)}$. In here, the series term is

$$s_n = \sum_{k=1}^n \frac{1}{k(k+1)} \quad (374)$$

$$= \sum_{k=1}^n \frac{1}{k} - \frac{1}{k+1} \quad (375)$$

$$= \sum_{k=1}^n \frac{1}{k} - \sum_{k=1}^n \frac{1}{k+1} \quad (376)$$

$$= \sum_{k=1}^n \frac{1}{k} - \sum_{k=2}^{n+1} \frac{1}{k} \quad (377)$$

$$= \frac{1}{1} + \left(\sum_{k=2}^n \frac{1}{k} \right) - \left(\sum_{k=2}^n \frac{1}{k} \right) - \frac{1}{n+1} \quad (378)$$

$$= 1 + \left(\sum_{k=2}^n \frac{1}{k} - \frac{1}{k} \right) - \frac{1}{n+1} \quad (379)$$

$$= 1 - \left(\sum_{k=2}^n 0 \right) - \frac{1}{n+1} \quad (380)$$

$$= 1 - \frac{1}{n+1} \quad (381)$$

Note that all of the examples that we have done here are for finite sums, so they are all legal.

Example 3.11 (Geometric Series)

The series $\sum_{n=0}^{\infty} q^n$ is called a **geometric series**.

$$1 + q + q^2 + \dots + q^n + \dots \quad (382)$$

is called the **geometric series**. We can see that

1. $|q| \geq 1 \iff \sum q^n$ is divergent. $|q| \geq 1 \implies |q|^n \geq 1$, and so the terms q^n does not converge to 0, and the n th term test is not met.

2. $|q| < 1 \iff \sum q^n$ is convergent. We can use the identity

$$s_n = 1 + q + \dots + q^{n-1} = \frac{1 - q^n}{1 - q} \implies \lim_{n \rightarrow \infty} \frac{1 - q^n}{1 - q} = \frac{1}{1 - q} \quad (383)$$

since $\lim_{n \rightarrow \infty} q^n = 0$ if $|q| < 1$.

The Cauchy convergence criterion can be used to prove the direct comparison test.

Theorem 3.10 (Direct Comparison Test)

For some fixed N , if

1. If $|x_n| \leq y_n$ for all $n \geq N$ and $\sum y_n$ converges, then $\sum x_n$ converges.
2. If $x_n \geq y_n \geq 0$ for all $n \geq N$ and $\sum y_n$ diverges, then $\sum x_n$ diverges.

Example 3.12 (Comparison with Telescoping Series)

We can prove the special case a geometric series with the direct comparison test. We claim that $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is finite. We can see that

$$\frac{1}{n^2} \leq \frac{2}{n(n+1)} \quad (384)$$

where the series of the terms in the RHS is telescoping and therefore converges. So by the direct comparison test, $\sum \frac{1}{n^2}$ converges.

Now we prove another corollary of the Cauchy convergence criterion.

Theorem 3.11 (Cauchy Condensation Test)

If $a_1 \geq a_2 \geq \dots \geq 0$, the series $\sum_{n=1}^{\infty} a_n$ converges if and only if the series

$$\sum_{k=0}^{\infty} 2^k a_{2^k} = a_1 + 2a_2 + 4a_4 + 8a_8 + \dots \quad (385)$$

converges.

Proof. Letting $A_k = a_1 + a_2 + \dots + a_k$ and $S_n = a_1 + 2a_2 + \dots + 2^n a_{2^n}$, it is clear that by adding up the inequalities

$$\begin{aligned} a_2 &\leq a_2 \leq a_1 \\ 2a_4 &\leq a_3 + a_4 \leq 2a_2 \\ 4a_8 &\leq a_5 + a_6 + a_7 + a_8 \leq 4a_4 \\ &\dots \\ 2^n a_{2^{n+1}} &\leq a_{2^n+1} + \dots + a_{2^{n+1}} \leq 2^n a_{2^n}, \end{aligned}$$

we get

$$\frac{1}{2}(S_{n+1} - a_1) \leq A_{2^{n+1}} - a_1 \leq S_n \quad (386)$$

Since the sequences $\{A_k\}$ and $\{S_k\}$ are nondecreasing, and hence from the inequalities we can conclude that they are either both bounded above (which means that they are both convergent since it is a bounded, nondecreasing series) or both unbounded above (which means that they are both divergent since they are nondecreasing and unbounded).

Corollary 3.3 (p-series Test)

The series

$$\sum_{n=1}^{\infty} \frac{1}{n^p} \quad (387)$$

converges for $p > 1$ and diverges for $p \leq 1$.^a

^aThis sort of reminds you of u -substitution. For example, look at $\int_1^{\infty} f(t) dt = \int_0^{\infty} e^u f(e^u) du$, where the convergence of LHS \iff convergence of RHS.

Proof. Suppose $p \geq 0$. By the previous theorem, the series converges or diverges simultaneously with the series

$$\sum_{k=0}^{\infty} 2^k \frac{1}{(2^k)^p} = \sum_{k=0}^{\infty} (2^{1-p})^k \quad (388)$$

which is really just a geometric series. A necessary and sufficient condition for the convergence of this series is that $2^{1-p} < 1$, that is, $p > 1$.

Now suppose $p \leq 0$. The series is then clearly divergent since all of the terms are larger than 1.

Example 3.13 (Harmonic Series)

The **harmonic series**

$$1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} + \dots \quad (389)$$

seems at first glance to be converging since the terms converge to 0. However, it does not pass the Cauchy condensation test since

$$\sum_{n=1}^{\infty} 2^n x_n = \sum_{n=1}^{\infty} 2^n \frac{1}{2^n} = \sum_{n=1}^{\infty} 1 = +\infty \quad (390)$$

As you can see, this increases logarithmically, so in early calculators it was hard to numerically detect divergence (you would have to double the number of series terms to get a linear increase).

3.5 Ratio and Root Tests

Now we introduce the root and ratio tests, which are derived by the comparison test with a geometric series. The ratio test is used more day-to-day, but not as decisive as the root test. Both tests have a similar flavor.

Theorem 3.12 (Ratio Test)

Suppose the limit $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \alpha$ exists for the series $\sum_{n=1}^{\infty} a_n$. Then,

1. $\alpha < 1 \implies \sum a_n$ converges absolutely.
2. $\alpha > 1 \implies \sum a_n$ diverges.
3. $\alpha = 1 \implies \sum a_n$ is inconclusive.

Alternatively, if

1. $\limsup |a_{n+1}/a_n| = \alpha < 1$, then $\sum a_n$ converges
2. If $\exists N$ s.t. $|a_{n+1}/a_n| \geq 1$ for all $n \geq N$, then $\sum a_n$ diverges.

Proof. Since $\limsup \left| \frac{a_{n+1}}{a_n} \right| = \alpha < 1$, fix any $\alpha < \beta < 1$. Then $\exists N$ s.t. if $n > N$, $|a_{n+1}/a_n| < \beta$. So $|a_{N+1}| < \beta|a_N| \implies |a_{N+2}| < \beta^2|a_N|$. So letting $C = |a_N|$, for all $m \geq N$,

$$|a_m| \leq \frac{C}{\beta^N} \beta^m \implies |a_m| \leq \tilde{C} \beta^m \text{ for all } m \geq N \quad (391)$$

So $\sum a_n$ converges by comparison test since $\sum \beta^m < \infty$ when $\beta < 1$.

Theorem 3.13 (Root Test)

Let $\sum_{n=1}^{\infty} a_n$ be a given series and

$$\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|} \quad (392)$$

Then,

1. $\alpha < 1 \implies \sum a_n$ converges.
2. $\alpha > 1 \implies \sum a_n$ diverges.
3. $\alpha = 1 \implies \sum a_n$ is inconclusive.

Proof. Listed.

1. If $\limsup \sqrt[n]{|a_n|} = \alpha < 1$, take any $\alpha < \beta < 1$. Then $\exists N \in \mathbb{N}$ s.t. if $n \geq N$, then $|a_n|^{1/n} < \beta \iff |a_n| < \beta^n$. Since $\beta < 1$, $\sum \beta^n < \infty$, and by comparison test, $\sum a_n$ converges.
2. Suppose $\alpha > 1$. Then $\limsup |a_n|^{1/n} = \alpha > 1$. So there exists a subsequence (a_{n_k}) s.t. $(|a_{n_k}|^{1/n_k}) \rightarrow \alpha > 1$. This means $\exists N$ s.t. for $n \geq N$, $|a_{n_k}|^{1/n_k} > 1 \implies |a_{n_k}| > 1$. But this fails the n th term test.
3. We do not claim anything and so there's nothing to prove.

Example 3.14 (Root Test Inconclusive Results)

Consider $\sum \frac{1}{n} = +\infty$, but from the root test

$$\sqrt[n]{\frac{1}{n}} \rightarrow 1, \text{ so } \alpha = 1 \quad (393)$$

Consider $\sum \frac{1}{n^2} < +\infty$, but from the root test

$$\sqrt[n]{\frac{1}{n^2}} = \left(\frac{1}{n^{1/n}} \right)^2 \rightarrow 1, \text{ so } \alpha = 1 \quad (394)$$

Example 3.15 ()

The sequence $\sum \frac{c^n}{n!}$ always converges for $c \in \mathbb{R}$.

Theorem 3.14 (Weierstrass M-test for Absolute Convergence)

Let $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ be series. Suppose there exists an index $N \in \mathbb{N}$ such that $|a_n| \leq b_n$ for all $n > N$. Then,

$$\sum_{n=1}^{\infty} b_n \text{ converges} \implies \sum_{n=1}^{\infty} a_n \text{ converges absolutely} \quad (395)$$

We finally conclude by giving a theorem about the convergence of some special sequences.

Theorem 3.15 (Special Sequences)

Some special sequences:

1. If $p > 0$, then $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 0$.
2. If $p > 0$, then $\lim_{n \rightarrow \infty} \sqrt[p]{p} = 1$.
3. $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$.
4. If $p > 0$ and α is real, then $\lim_{n \rightarrow \infty} \frac{n^\alpha}{(1+p)^n} = 0$.
5. If $|x| < 1$, then $\lim_{n \rightarrow \infty} x^n = 0$.

3.6 Euler's Number and Trigonometric Functions

Definition 3.7 (Euler's Number)

We define **Euler's number** as

$$e := \sum_{n=0}^{\infty} \frac{1}{n!} \quad (396)$$

The first thing we should do is show that it converges, this is a one-liner.

$$\sum_{n=0}^{\infty} \frac{1}{n!} = 1 + 1 + \sum_{n=2}^{\infty} \frac{1}{n!} \leq 2 + \sum_{n=2}^{\infty} \frac{1}{n(n-1)} \quad (397)$$

Theorem 3.16 (Euler's Number as a Limit)

We have

$$\lim_{n \rightarrow +\infty} \left(1 + \frac{1}{n}\right)^n = e \quad (398)$$

Proof. Let us define the sequence

$$t_n = \sum_{k=0}^n \frac{1}{k!}, \quad s_n = \left(1 + \frac{1}{n}\right)^n \quad (399)$$

We know that $t_n \rightarrow e$, and we want to show that $s_n \rightarrow e$. We do this with the squeeze theorem.

1. We can see that

$$s_n = \left(1 + \frac{1}{n}\right)^n \quad (400)$$

$$= \sum_{k=0}^n \binom{n}{k} 1^{n-k} \left(\frac{1}{n}\right)^k \quad (401)$$

$$= 1 + 1 + \frac{n(n-1)}{2!} \frac{1}{n^2} + \frac{n(n-1)(n-2)}{3!} \frac{1}{n^3} + \dots \quad (402)$$

$$= 1 + 1 + \frac{1}{2!}(1) \left(1 - \frac{1}{n}\right) + \frac{1}{3!}(1) \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) + \dots + \frac{1}{n!}(1) \prod_{k=1}^{n-1} \left(1 - \frac{k}{n}\right) \quad (403)$$

$$\leq \frac{1}{0!} + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{n!} = t_n \quad (404)$$

and so $s_n < t_n \implies \limsup s_n \leq \limsup t_n = e$.

2. Let $m \leq n$ be fixed. Then,

$$s_n \geq 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n}\right) + \dots + \frac{1}{m!} \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \dots \left(1 - \frac{m-1}{n}\right) \quad (405)$$

since we are just taking the first m positive terms of the element. Therefore, letting $n \rightarrow +\infty$ and keeping m fixed, we get

$$\liminf_{n \rightarrow +\infty} s_n \geq 1 + 1 + \frac{1}{2!} + \dots + \frac{1}{n!} \text{ for all } m \in \mathbb{N} \quad (406)$$

which implies $\liminf s_n \geq t_m$ for all $m \in \mathbb{N}$, and now letting $m \rightarrow +\infty$, we have $\liminf s_n \geq \liminf t_m = e$.

Now we prove the irrationality of e . It is usually extremely difficult to prove that an arbitrary number is irrational, e.g. π^e or π^{e^e} .

Theorem 3.17 (e is Irrational)

e is irrational.

Proof. Letting $t_n = \sum_{k=0}^n \frac{1}{k!}$, we have

$$e - t_n = \sum_{k=n+1}^{\infty} \frac{1}{k!} \quad (407)$$

$$= \frac{1}{(n+1)!} \left(1 + \frac{1}{n+2} + \frac{1}{(n+3)(n+2)} + \dots\right) \quad (408)$$

$$< \frac{1}{(n+1)!} \underbrace{\left(1 + \frac{1}{n+2} + \frac{1}{(n+2)^2} + \dots\right)}_{\text{geometric}} \quad (409)$$

$$= \frac{1}{(n+1)!} \left(\frac{1}{1 - (1/(n+2)!)}\right) \quad (410)$$

$$= \frac{1}{n!n} \cdot \underbrace{\frac{(n+2)n}{(n+1)^2}}_{<1} \quad (411)$$

$$= \frac{1}{n!n} \quad (412)$$

Note that we can combine and split sums since we know that e is convergent. Now suppose that $e = p/q$. Then,

$$0 < q!(e - t_q) < \frac{1}{q} \quad (413)$$

But $q!e$ is an integer and $q!t_q$ is also an integer. So we have $q! \cdot \frac{p}{q}$, an integer, between 0 and 1, which is a contradiction.

Since we have defined some number $e \in \mathbb{R}$, we know that exponential exist, and therefore we the function $x \mapsto e^x$ is well-defined. In fact, it is so important that we have a separate name for it.

Definition 3.8 (Exponential Function)

The **exponential function** is generally referred to as the function $x \mapsto e^x$.

There is a nice series representation.

Theorem 3.18 (Exponential Function as a Series)

We have

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad (414)$$

Proof.

Now that this is done, we can define the trigonometric functions formally as such.

Definition 3.9 (Trigonometric Functions)

We have

$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \quad (415)$$

(416)

$$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots \quad (417)$$

3.7 Exercises**Exercise 3.1 (Math 531 Spring 2025, PS4.6)**

Consider the set of all bounded sequences of real numbers. That is, we consider sequences $\{x_n\}$ for which

$$\sup_{n \in \mathbb{N}} |x_n| \quad (418)$$

exists. For example, the sequence $\{1, 2, 3, \dots\}$ does not belong to the set, but the sequence $\{1, -\frac{1}{2}, \frac{1}{3}, -\frac{1}{4}, \dots\}$ does. Call this set X . Endow it with a metric:

$$d(\{x_n\}, \{y_n\}) = \sup_{n \in \mathbb{N}} |x_n - y_n|. \quad (419)$$

Explain why this is a metric. Make sure to explain why the supremum on the right hand side exists.

Solution.

Exercise 3.2 (Math 531 Spring 2025, PS4.7)

Consider the metric space (X, d) from the previous problem. Is $\overline{B_1(\{0\})}$ a compact set? Here, $\{0\}$ is just the sequence of zeros: $\{0, 0, 0, 0, \dots\}$.

Solution.

Exercise 3.3 (Math 531 Spring 2025, PS6.1)

Give a direct proof that

$$\sum_{n=1}^{\infty} \frac{1}{k^2 + k} \quad (420)$$

converges and find the exact value of the series.

Solution. This is a telescoping series.

$$\sum_{n=1}^{\infty} \frac{1}{k^2 + k} = \sum_{n=1}^{\infty} \frac{1}{k} - \frac{1}{k-1} = 1 - \frac{1}{2} + \frac{1}{2} - \frac{1}{3} + \frac{1}{3} - \frac{1}{4} + \dots \quad (421)$$

and so the n th partial sum is $1 - \frac{1}{n} \rightarrow 1$ as $n \rightarrow \infty$.

Exercise 3.4 (Math 531 Spring 2025, PS6.2)

Suppose that a_n is a sequence of non-negative real numbers and suppose that $\sum_{n=1}^{\infty} a_n$ converges. Prove that there exists a sequence b_n with $\lim_{n \rightarrow \infty} b_n = +\infty$ so that $\sum_{n=1}^{\infty} a_n b_n$ is still convergent.

Solution. Let us take $r_n = \sum_{m=n}^{\infty} a_m$. Since a_n is nonnegative, $r_n = 0 \iff a_m = 0$ for all $m \geq n$. In this case set

$$(b_m) := \begin{cases} 1 & \text{if } m < n \\ m & \text{if } m \geq n \end{cases} \quad (422)$$

and

$$\sum_{n=1}^{\infty} a_n = \sum_{m=1}^{\infty} a_m = \sum_{m=1}^n a_m b_m = \sum_{m=1}^{\infty} a_m b_m < +\infty \quad (423)$$

So we may assume $r_n > 0$. Now define $(b_n) = (\frac{1}{\sqrt{r_n}})$. By the Cauchy criterion, $r_n \rightarrow 0$ as $n \rightarrow +\infty$, so $b_n \rightarrow +\infty$. But we claim that $\frac{a_n}{\sqrt{r_n}} < 2(\sqrt{r_n} - \sqrt{r_{n+1}})$ since

$$\frac{a_n}{\sqrt{r_n}}(\sqrt{r_n} + \sqrt{r_{n+1}}) = a_n + a_n \frac{\sqrt{r_{n+1}}}{\sqrt{r_n}} < 2a_n = 2(r_n - r_{n+1}) \quad (424)$$

Since the series $\sum(\sqrt{r_n} - \sqrt{r_{n+1}})$ converges to $\sqrt{r_1}$, it follows by the comparison test that $\sum \frac{a_n}{\sqrt{r_n}}$ also converges.

Exercise 3.5 (Math 531 Spring 2025, PS6.3)

Prove that $\sum_{n=1}^{\infty} \frac{\sin(n)}{n}$ converges^a. Hint: Use summation by parts and the fact that $\sin(x) = \frac{1}{2i}(\exp(ix) - \exp(-ix))$ where $i = \sqrt{-1}$.

^aHere, we are going to assume that we know a little bit about trigonometric functions!

Solution. We can see that

$$\left| \operatorname{Im} \sum_{n=1}^N e^{in} \right| = \left| \sum_{n=1}^N \sin(n) \right| \leq \left| e^i \frac{1 - e^{iN}}{1 - e^i} \right| \leq \frac{2}{|1 - e^i|} < +\infty \quad (425)$$

and so the partial sums form a bounded sequence, implying that $\sum \frac{\sin(n)}{n}$ is convergent.

Exercise 3.6 (Math 531 Spring 2025, PS6.4)

Fix $c \in (0, \infty)$.

- Assume x_n is a sequence of numbers with $x_n \rightarrow 0$. Prove that

$$c^{x_n} \rightarrow 1. \quad (426)$$

- Deduce that, if $x_n \rightarrow x$, then

$$c^{x_n} \rightarrow c^x, \quad (427)$$

for any $x \in \mathbb{R}$.

Solution. $x_n \rightarrow 0$ as $n \rightarrow \infty$ means that $\forall \delta > 0, \exists N \in \mathbb{N}$ s.t. $|x_n| < \delta$ for all $n \geq N$. Now take $\epsilon > 0$. Then we wish to show that $\exists M \in \mathbb{N}$ s.t. $|1 - c^{x_n}| < \epsilon$ for all $n \geq M$. We choose $\delta = \min\{-\log_c(1 - \epsilon), \log_c(1 + \epsilon)\}$. Then $\exists M$ s.t.

$$x_n \in (-\log_c(1 - \epsilon), \log_c(1 + \epsilon)) \iff |x_n| < \delta \quad (428)$$

for $n \geq M$ by convergence $x_n \rightarrow 0$. Since $f(x) = c^x$ is monotonically increasing, this is equivalent to

$$f(x_n) = c^{x_n} \quad (429)$$

Exercise 3.7 (Math 531 Spring 2025, PS6.5)

For every $z \in \mathbb{C}$, verify that the series

$$E(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!} \quad (430)$$

converges. Prove that for every $z, w \in \mathbb{C}$ we have that $E(z)E(w) = E(z + w)$. Deduce that, for $q \in \mathbb{Q}$, we have that $E(q) = e^q$. Deduce that $E(x) = e^x$, for all $x \in \mathbb{R}$.

Solution.

Exercise 3.8 (Rudin 3.1)

Prove that convergence of $\{x_n\}$ implies convergence of $\{|x_n|\}$. Is the converse true?

Solution. If $\{x_n\}$ converges to x , then for all $\epsilon > 0$, there exists a $N \in \mathbb{N}$ s.t. $|x_n - x| < \epsilon$ if $n > N$. We use the inequality $||x_n| - |x|| \leq |x_n - x|$ to show that then for every $\epsilon > 0$ there exists a $N \in \mathbb{N}$ s.t.

$$||x_n| - |x|| \leq |x_n - x| \leq \epsilon$$

and so $\{|x_n|\}$ converges to $|x|$.

Exercise 3.9 (Rudin 3.2)

Calculate

$$\lim_{n \rightarrow \infty} \sqrt{n^2 + n} - n$$

Solution. We can compute

$$\lim_{n \rightarrow \infty} \sqrt{n^2 + n} - n = \lim_{n \rightarrow \infty} (\sqrt{n^2 + n} - n) \cdot \frac{\sqrt{n^2 + n} + n}{\sqrt{n^2 + n} + n} = \lim_{n \rightarrow \infty} \frac{n}{\sqrt{n^2 + n} + n}$$

where

$$A_n = \frac{n}{\sqrt{n^2 + 2n + 1} + n} \leq \frac{n}{2n + 1} \leq \frac{n}{\sqrt{n^2 + n} + n} \leq \frac{n}{\sqrt{n^2 + n}} = \frac{n}{2n} = \frac{1}{2} = C_n$$

 C_n is ultimately constant. It suffices to prove that A_n limits to $\frac{1}{2}$ by showing that

$$\frac{n}{2n + 1} = \frac{n/n}{(2n + 1)/n} = \frac{1}{2 + \frac{1}{n}}$$

where $\{\frac{1}{n}\}$ is infinitesimal.**Exercise 3.10 (Rudin 3.3)**If $s_1 = \sqrt{2}$ and

$$s_{n+1} = \sqrt{2 + \sqrt{s_n}}$$

for $n = 1, 2, \dots$, prove that $\{s_n\}$ converges and that $s_n < 2$ for $n = 1, 2, \dots$ *Solution.* We can show that $s_n < 2$ by induction. $s_1 = \sqrt{2} < 2$, so the base case is proved. Now, given that $s_n < 2$, $\sqrt{s_n} < 2 \implies 2 + \sqrt{s_n} < 2 + \sqrt{2} < 4 \implies s_{n+1} = \sqrt{2 + \sqrt{s_n}} < 2$ and we are done.**Exercise 3.11 (Rudin 3.4)**Find the upper and lower limits of the sequence $\{s_n\}$ defined by

$$s_1 = 0; \quad s_{2m} = \frac{s_{2m-1}}{2}; \quad s_{2m+1} = \frac{1}{2} + s_{2m}.$$

*Solution.***Exercise 3.12 (Rudin 3.5)**For any two real sequences $\{a_n\}$, $\{b_n\}$, prove that

$$\limsup_{n \rightarrow \infty} (a_n + b_n) \leq \limsup_{n \rightarrow \infty} a_n + \limsup_{n \rightarrow \infty} b_n,$$

provided the sum on the right is not of the form $\infty - \infty$.*Solution.*

Exercise 3.13 (Rudin 3.6)

Investigate the behavior (convergence or divergence) of $\sum a_n$ if

- (a) $a_n = \sqrt{n+1} - \sqrt{n}$;
- (b) $a_n = (\sqrt{n+1} - \sqrt{n})/n$;
- (c) $a_n = (\sqrt[n]{n} - 1)^n$;
- (d) $a_n = \frac{1}{1+z^n}$, for complex values of z .

Solution.

Exercise 3.14 (Rudin 3.7)

Prove that the convergence of $\sum a_n$ implies the convergence of

$$\sum \frac{\sqrt{a_n}}{n},$$

if $a_n \geq 0$.

Solution.

Exercise 3.15 (Rudin 3.8)

If $\sum a_n$ converges, and if $\{b_n\}$ is monotonic and bounded, prove that $\sum a_n b_n$ converges.

Solution.

Exercise 3.16 (Rudin 3.9)

Find the radius of convergence of each of the following power series:

- (a) $\sum n^3 z^n$,
- (b) $\sum \frac{2^n}{n!} z^n$,
- (c) $\sum \frac{2^n}{n^2} z^n$,
- (d) $\sum \frac{n^3}{3^n} z^n$.

Solution.

Exercise 3.17 (Rudin 3.10)

Suppose that the coefficients of the power series $\sum a_n z^n$ are integers, infinitely many of which are distinct from zero. Prove that the radius of convergence is at most 1.

Solution.

Exercise 3.18 (Rudin 3.11)

Suppose $a_n > 0$, $s_n = a_1 + \cdots + a_n$, and $\sum a_n$ diverges.

- (a) Prove that $\sum \frac{a_n}{1+a_n}$ diverges.

(b) Prove that

$$\frac{a_{N+1}}{s_{N+1}} + \cdots + \frac{a_{N+k}}{s_{N+k}} \geq 1 - \frac{s_N}{s_{N+k}}$$

and deduce that $\sum \frac{a_n}{s_n}$ diverges.

(c) Prove that

$$\frac{a_n}{s_n^2} \leq \frac{1}{s_{n-1}} - \frac{1}{s_n}$$

and deduce that $\sum \frac{a_n}{s_n^2}$ converges.

(d) What can be said about

$$\sum \frac{a_n}{1 + na_n} \text{ and } \sum \frac{a_n}{1 + n^2 a_n}?$$

Solution.

Exercise 3.19 (Rudin 3.12)

Suppose $a_n > 0$ and $\sum a_n$ converges. Put

$$r_n = \sum_{m=n}^{\infty} a_m.$$

(a) Prove that

$$\frac{a_m}{r_m} + \cdots + \frac{a_n}{r_n} > 1 - \frac{r_n}{r_m}$$

if $m < n$, and deduce that $\sum \frac{a_n}{r_n}$ diverges.

(b) Prove that

$$\frac{a_n}{\sqrt{r_n}} < 2(\sqrt{r_n} - \sqrt{r_{n+1}})$$

and deduce that $\sum \frac{a_n}{\sqrt{r_n}}$ converges.

Solution.

Exercise 3.20 (Rudin 3.13)

Prove that the Cauchy product of two absolutely convergent series converges absolutely.

Solution.

Exercise 3.21 (Rudin 3.14)

If $\{s_n\}$ is a complex sequence, define its arithmetic means σ_n by

$$\sigma_n = \frac{s_0 + s_1 + \cdots + s_n}{n+1} \quad (n = 0, 1, 2, \dots).$$

- (a) If $\lim s_n = s$, prove that $\lim \sigma_n = s$.
 (b) Construct a sequence $\{s_n\}$ which does not converge, although $\lim \sigma_n = 0$.
 (c) Can it happen that $s_n > 0$ for all n and that $\limsup s_n = \infty$, although $\lim \sigma_n = 0$?
 (d) Put $a_n = s_n - s_{n-1}$, for $n \geq 1$. Show that

$$s_n - \sigma_n = \frac{1}{n+1} \sum_{k=1}^n k a_k.$$

Assume that $\lim(na_n) = 0$ and that $\{\sigma_n\}$ converges. Prove that $\{s_n\}$ converges. [This gives a converse of (a), but under the additional assumption that $na_n \rightarrow 0$.]

- (e) Derive the last conclusion from a weaker hypothesis: Assume $M < \infty$, $|na_n| \leq M$ for all n , and $\lim \sigma_n = \sigma$. Prove that $\lim s_n = \sigma$, by completing the following outline:
 If $m < n$, then

$$s_n - \sigma_n = \frac{m+1}{n-m}(\sigma_n - \sigma_m) + \frac{1}{n-m} \sum_{i=m+1}^n (s_n - s_i).$$

For these i ,

$$|s_n - s_i| \leq \frac{(n-i)M}{i+1} \leq \frac{(n-m-1)M}{m+2}.$$

Fix $\varepsilon > 0$ and associate with each n the integer m that satisfies

$$m \leq \frac{n-\varepsilon}{1+\varepsilon} < m+1.$$

Then $(m+1)/(n-m) \leq 1/\varepsilon$ and $|s_n - s_i| < M\varepsilon$. Hence

$$\limsup_{n \rightarrow \infty} |s_n - \sigma| \leq M\varepsilon.$$

Since ε was arbitrary, $\lim s_n = \sigma$.

Solution.

Exercise 3.22 (Rudin 3.15)

Definition 3.21 can be extended to the case in which the a_n lie in some fixed \mathbb{R}^k . Absolute convergence is defined as convergence of $\sum |\mathbf{a}_n|$. Show that Theorems 3.22, 3.23, 3.25(a), 3.33, 3.34, 3.42, 3.45, 3.47, and 3.55 are true in this more general setting. (Only slight modifications are required in any of the proofs.)

Solution.

Exercise 3.23 (Rudin 3.16)

Fix a positive number α . Choose $x_1 > \sqrt{\alpha}$, and define x_2, x_3, x_4, \dots , by the recursion formula

$$x_{n+1} = \frac{1}{2} \left(x_n + \frac{\alpha}{x_n} \right).$$

- (a) Prove that $\{x_n\}$ decreases monotonically and that $\lim x_n = \sqrt{\alpha}$.

(b) Put $\varepsilon_n = x_n - \sqrt{\alpha}$, and show that

$$\varepsilon_{n+1} = \frac{\varepsilon_n^2}{2x_n} < \frac{\varepsilon_n^2}{2\sqrt{\alpha}}$$

so that, setting $\beta = 2\sqrt{\alpha}$,

$$\varepsilon_{n+1} < \beta \left(\frac{\varepsilon_1}{\beta} \right)^{2^n} \quad (n = 1, 2, 3, \dots).$$

(c) This is a good algorithm for computing square roots, since the recursion formula is simple and the convergence is extremely rapid. For example, if $\alpha = 3$ and $x_1 = 2$, show that $\varepsilon_1/\beta < \frac{1}{10}$ and that therefore

$$\varepsilon_5 < 4 \cdot 10^{-16}, \quad \varepsilon_6 < 4 \cdot 10^{-32}.$$

Solution.

Exercise 3.24 (Rudin 3.17)

Fix $\alpha > 1$. Take $x_1 > \sqrt{\alpha}$, and define

$$x_{n+1} = \frac{\alpha + x_n}{1 + x_n} = x_n + \frac{\alpha - x_n^2}{1 + x_n}.$$

- (a) Prove that $x_1 > x_3 > x_5 > \dots$.
- (b) Prove that $x_2 < x_4 < x_6 < \dots$.
- (c) Prove that $\lim x_n = \sqrt{\alpha}$.
- (d) Compare the rapidity of convergence of this process with the one described in Exercise 16.

Solution.

Exercise 3.25 (Rudin 3.18)

Replace the recursion formula of Exercise 16 by

$$x_{n+1} = \frac{p-1}{p}x_n + \frac{\alpha}{p}x_n^{-p+1}$$

where p is a fixed positive integer, and describe the behavior of the resulting sequences $\{x_n\}$.

Solution.

Exercise 3.26 (Rudin 3.19)

Associate to each sequence $a = \{\alpha_n\}$, in which α_n is 0 or 2, the real number

$$x(a) = \sum_{n=1}^{\infty} \frac{\alpha_n}{3^n}.$$

Prove that the set of all $x(a)$ is precisely the Cantor set described in Sec. 2.44.

Solution.

Exercise 3.27 (Rudin 3.20)

Suppose $\{p_n\}$ is a Cauchy sequence in a metric space X , and some subsequence $\{p_{n_i}\}$ converges to a point $p \in X$. Prove that the full sequence $\{p_n\}$ converges to p .

Solution.

Exercise 3.28 (Rudin 3.21)

Prove the following analogue of Theorem 3.10(b): If $\{E_n\}$ is a sequence of closed nonempty and bounded sets in a complete metric space X , if $E_n \supset E_{n+1}$, and if

$$\lim_{n \rightarrow \infty} \text{diam } E_n = 0,$$

then $\bigcap_{n=1}^{\infty} E_n$ consists of exactly one point.

Solution.

Exercise 3.29 (Rudin 3.22)

Suppose X is a nonempty complete metric space, and $\{G_n\}$ is a sequence of dense open subsets of X . Prove Baire's theorem, namely, that $\bigcap_{n=1}^{\infty} G_n$ is not empty. (In fact, it is dense in X .) Hint: Find a shrinking sequence of neighborhoods E_n such that $E_n \subset G_n$, and apply Exercise 21.

Solution.

Exercise 3.30 (Rudin 3.23)

Suppose $\{p_n\}$ and $\{q_n\}$ are Cauchy sequences in a metric space X . Show that the sequence $\{d(p_n, q_n)\}$ converges. Hint: For any m, n ,

$$d(p_n, q_n) \leq d(p_n, p_m) + d(p_m, q_m) + d(q_m, q_n);$$

it follows that

$$|d(p_n, q_n) - d(p_m, q_m)|$$

is small if m and n are large.

Solution.

Exercise 3.31 (Rudin 3.24)

Let X be a metric space.

(a) Call two Cauchy sequences $\{p_n\}, \{q_n\}$ in X equivalent if

$$\lim_{n \rightarrow \infty} d(p_n, q_n) = 0.$$

Prove that this is an equivalence relation.

- (b) Let X^* be the set of all equivalence classes so obtained. If $P \in X^*, Q \in X^*, \{p_n\} \in P, \{q_n\} \in Q$, define

$$\Delta(P, Q) = \lim_{n \rightarrow \infty} d(p_n, q_n);$$

by Exercise 23, this limit exists. Show that the number $\Delta(P, Q)$ is unchanged if $\{p_n\}$ and $\{q_n\}$ are replaced by equivalent sequences, and hence that Δ is a distance function in X^* .

- (c) Prove that the resulting metric space X^* is complete.
 (d) For each $p \in X$, there is a Cauchy sequence all of whose terms are p ; let P_p be the element of X^* which contains this sequence. Prove that

$$\Delta(P_p, P_q) = d(p, q)$$

for all $p, q \in X$. In other words, the mapping φ defined by $\varphi(p) = P_p$ is an isometry (i.e., a distance-preserving mapping) of X into X^* .

- (e) Prove that $\varphi(X)$ is dense in X^* , and that $\varphi(X) = X^*$ if X is complete. By (d), we may identify X and $\varphi(X)$ and thus regard X as embedded in the complete metric space X^* . We call X^* the completion of X .

Solution.

Exercise 3.32 (Rudin 3.25)

Let X be the metric space whose points are the rational numbers, with the metric $d(x, y) = |x - y|$. What is the completion of this space? (Compare Exercise 24.)

Solution.

4 Limits and Continuity of Functions

We now extend our analysis to real-valued functions over a metric space. The ones that we will be particularly interested in are *continuous functions*. But before this, let's introduce a new notation. Given a metric space X , we will talk about a variable x approaching a particular value $a \in X$, denoted $x \rightarrow a$. But this isn't clear. When we talk about the concept of something approaching another thing, we have two definitions.

1. A *sequence* can approach to its limit, which is a *point*.
2. A *point* can be a limit point of a *set*.

When we write $x \rightarrow a$, we are talking about some indeterminate variable x and a point a , it isn't immediately clear what this means. As we will soon define, this will refer to a neighborhood of a or equivalently to *all* sequences converging to a . So we can think of $x \rightarrow a$ as notation for all sequences $(x_n) \rightarrow a$.

Definition 4.1 (Constant and Ultimately Constant Functions)

Given a real-valued function $f : E \rightarrow \mathbb{R}$ defined on domain $E \subset \mathbb{R}$,

1. f is a **constant function** if $f(x) = A$ for all $x \in E$
2. f is called **ultimately constant** as $x \rightarrow a$ if it is constant in some deleted neighborhood $\dot{U}(a)$, where a is a limit point of E .

Definition 4.2 (Bounded and Ultimately Bounded Functions)

Given a real-valued function $f : E \rightarrow \mathbb{R}$ defined on domain $E \subset \mathbb{R}$,

1. f is **bounded**, **bounded above**, or **bounded below** respectively if there is a number $C \in \mathbb{R}$ such that $|f(x)| < C$, $f(x) < C$, or $C < f(x)$ for all $x \in E$.
2. f is **ultimately bounded**, **ultimately bounded above**, or **ultimately bounded below** as $x \rightarrow a$ if it is bounded, bounded above, or bounded below in some deleted neighborhood $\dot{U}_E(a)$.

Example 4.1 (Unbounded but Ultimately Bounded)

The function

$$f(x) = \sin \frac{1}{x} + x \cos \frac{1}{x} \quad (431)$$

for $x \neq 0$ is not bounded on the domain of definition, but it is ultimately bounded as $x \rightarrow 0$.

4.1 Limits of Functions

Definition 4.3 (Limit of a Function)

Let $f : X \rightarrow Y$ be a map between metric spaces, with $E \subset X$ and $p \in E'$ (note the limit point!). We say $f(x) \rightarrow q$ as $x \rightarrow p$, i.e.

$$\lim_{x \rightarrow p} f(x) = q \quad (432)$$

if it meets the following equivalent conditions.

1. ϵ - δ Definition. If $\forall \epsilon > 0, \exists \delta > 0$ s.t. $0 < d_X(x, p) < \delta \implies d_Y(f(x), q) < \epsilon$.^a




Figure 2: Said in one line, the preimage of any open ball around $y = f(x)$ must contain some open deleted open ball around x .

2. *Sequential Definition.* If for all sequences $(x_n) \rightarrow p$, $f(x_n) \rightarrow q$.




Figure 3: For every sequence that converges to the left, the new sequence mapped through f converges to q . Note that we choose the points x_n to be in the "deleted" neighborhood $E \setminus a$ (neighborhood E with point a removed) to force us to choose a sequence that is not a, a, \dots . That is, it forces us to choose different points for the sequence.

^aNote that the strictly inequality $0 < d_X(x, p)$ is important to ensure that $x \neq p$, since functions can jump at p .

Proof. We prove equivalence.

1. (\rightarrow) . Assume $\lim_{x \rightarrow p} f(x) = q$. Let $(x_n) \in E$ s.t. $x_n \rightarrow p$ with $x_n \neq p$. We wish to show that $f(x_n) \rightarrow q$. Let $\epsilon > 0$. Then $\exists \delta > 0$ s.t. $0 < d_X(x, p) < \delta \implies d_Y(f(x), q) < \epsilon$. Since $\delta > 0$, by definition $\exists N \in \mathbb{N}$ s.t. if $n \geq N$, $d_X(x_n, p) < \delta \implies d_Y(f(x_n), q) < \epsilon$.

Sometimes, the ϵ - δ definition is good, but a lot of the times the sequential definition is good enough and more insightful.

Example 4.2 (Limit of the Signum Function)

The function $\text{sgn}: \mathbb{R} \rightarrow \mathbb{R}$ defined

$$\text{sgn } x = \begin{cases} 1, & x > 0 \\ 0, & x = 0 \\ -1, & x < 0 \end{cases} \quad (433)$$

has no limit as $x \rightarrow 0$.

First, it is ludicrous that the limit would be any number that is not $\{-1, 0, 1\}$. If we assume that $A \notin \{-1, 0, 1\}$, then we can choose any arbitrarily small ϵ -neighborhood of A that does not include the three numbers. Clearly, there doesn't exist any $\delta > 0$ such that the deleted δ -neighborhood of 0 maps to a set completely contained in the ϵ -neighborhood of A . That is,

$$\text{sgn}(\dot{U}_\delta(0)) = \{-1, 1\} \not\subset U_\epsilon(A) \quad (434)$$

It doesn't even intersect the ϵ -neighborhood at all.

1. If $A = 1$, we can construct a ϵ -neighborhood V_A for $\epsilon = \frac{1}{2}$. Clearly, there exists no open neighborhood U_0 of 0 that is entirely mapped to V , since U_0 contains both negative numbers and 0 and hence must be mapped to 0, -1.
2. Similarly, given the $(\epsilon = \frac{1}{2})$ -neighborhood of $A = -1$, there exists no open neighborhood U_0 of 0 that is entirely mapped to it, since U_0 contains both positive numbers and 0 and hence must be mapped to 0, 1.
3. Finally, given the $(\epsilon = \frac{1}{2})$ -neighborhood of $A = 0$, there exists no open neighborhood U_0 of 0 that is entirely mapped to it, since U_0 contains both positive and negative numbers and hence must be mapped to ± 1 .

Therefore, the limit does not exist.

Example 4.3 (Limit of Absolute Value of Signum Function)

We will show that

$$\lim_{x \rightarrow 0} |\operatorname{sgn} x| = 1 \quad (435)$$

We construct a ϵ -neighborhood $U_\epsilon(1)$ around 1. Given this neighborhood, we can imagine choosing the deleted δ -neighborhood $\dot{U}_\delta(0)$ around 0. Since every element in $\dot{U}_\delta(0)$ maps to 1, it is clearly in U_ϵ . In fact, for arbitrarily small $\epsilon > 0$, we can choose **any** $\delta > 0$ since everything in $\mathbb{R} \setminus 0$ maps to 1. We can visualize this in \mathbb{R}^2 as

Theorem 4.1 (Arithmetic on Limits of Functions)

Given two numerical valued functions $f, g : E \subset \mathbb{R} \rightarrow \mathbb{R}$ with a common domain where $g(x) \neq 0$ for all $x \in E$, let

$$\lim_{x \rightarrow a} f(x) = A, \quad \lim_{x \rightarrow a} g(x) = B \quad (436)$$

then,

$$\lim_{x \rightarrow a} (f + g)(x) = A + B$$

$$\lim_{x \rightarrow a} (cf)(x) = cA$$

$$\lim_{x \rightarrow a} (f \cdot g)(x) = A \cdot B$$

$$\lim_{x \rightarrow a} \left(\frac{f}{g} \right)(x) = \frac{A}{B}$$

Proof. Cauchy sequence criterion for a limit immediately proves this.

We end this with a theorem connecting the relationship between a limit of a function as $x \rightarrow a$ and its ultimate behavior as $x \rightarrow a$.

Theorem 4.2 ()

Let $f : E \rightarrow \mathbb{R}$ be a function. Then,

1. f is ultimately the constant A as $x \rightarrow a$ implies that $\lim_{x \rightarrow a} f(x) = A$.
2. $\lim_{x \rightarrow a} f(x)$ implies that f is ultimately bounded as $x \rightarrow a$.

Definition 4.4 (Infinitesimal Function)

A function $f : E \subset \mathbb{R} \rightarrow \mathbb{R}$ is said to be **infinitesimal** as $x \rightarrow a$ if

$$\lim_{x \rightarrow a} f(x) = 0 \quad (437)$$

Lemma 4.1 (Sums, Products of Infinitesimals)

It is clear that if α, β are infinitesimal as $x \rightarrow a$, then

1. $\alpha + \beta$ is infinitesimal as $x \rightarrow a$
2. $\alpha \cdot \beta$ is infinitesimal as $x \rightarrow a$

Furthermore, if α is infinitesimal and β is ultimately bounded as $x \rightarrow a$, then the product $\alpha \cdot \beta$ is infinitesimal as $x \rightarrow a$.

Proof. We prove all three statements.

1. Assume that α and β are infinitesimal as $x \rightarrow a$. Then, let us fix a small $\epsilon > 0$. This means that for every $\frac{\epsilon}{2}$ there exists an open deleted neighborhood $\dot{U}'(a)$ such that its image $\alpha(\dot{U}'(a)) \subset U'_{\epsilon/2}(0) \subset \mathbb{R}$. Additionally, for every $\frac{\epsilon}{2}$ there exists an open deleted neighborhood $\dot{U}''(a)$ such that its image $\beta(\dot{U}''(a)) \subset U'_{\epsilon/2}(0) \subset \mathbb{R}$. Thus, for the deleted neighborhood

$$\dot{U}(a) \subset \dot{U}'(a) \cup \dot{U}''(a) \quad (438)$$

we can see that for all $x \in \dot{U}(a)$,

$$|(\alpha + \beta)(x)| = |\alpha(x) + \beta(x)| \leq |\alpha(x)| + |\beta(x)| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \quad (439)$$

and hence $(\alpha + \beta)(\dot{U}(a)) \subset U_\epsilon(0)$.

2. This case is a special case of assertion 3. That is, every function that has a limit is ultimately bounded.
3. Since $\beta(x)$ is ultimately bounded, this means that there exists a constant M and an open deleted neighborhood $\dot{U}'(a) \subset E$ such that for all $x \in \dot{U}'(a)$, its image is bounded: $|\beta(x)| < M$. Let us fix a small $\epsilon > 0$. Then, by definition of the limit, for every $\frac{\epsilon}{M}$ there exists an open deleted neighborhood $\dot{U}''(a)$ such that its image $\beta(\dot{U}''(a)) \subset U_{\epsilon/M}(0) \subset \mathbb{R}$. Therefore, for the deleted neighborhood

$$\dot{U}(a) \subset \dot{U}'(a) \cup \dot{U}''(a) \quad (440)$$

we can see that for all $x \in \dot{U}(a)$,

$$|(\alpha \cdot \beta)(x)| = |\alpha(x)\beta(x)| = |\alpha(x)||\beta(x)| < \frac{\epsilon}{M} \cdot M = \epsilon \quad (441)$$

Therefore, $(\alpha \cdot \beta)(\dot{U}(a)) \subset U_\epsilon(0)$.

Note that in proving these properties of the limits, we have used the following fact about open deleted neighborhoods around a .

1. $\dot{U}(a)$ is not the empty set.
2. Given open deleted neighborhoods $\dot{U}'(a)$ and $\dot{U}''(a)$, there exists an open deleted neighborhood in the intersections of these neighborhoods.

$$\dot{U}(a) \subset \dot{U}'(a) \cup \dot{U}''(a) \quad (442)$$

Theorem 4.3 (Representation of a Convergent Function as a Shift of its Infinitesimal)

Given a function $f : E \subset \mathbb{R} \rightarrow \mathbb{R}$, its limit exists and

$$\lim_{x \rightarrow a} f(x) = A \quad (443)$$

if and only if f can be represented as

$$f(x) = A + \alpha(x) \quad (444)$$

where α is infinitesimal as $x \rightarrow a$. We can visualize this theorem by thinking of a function f that results from a "shift" of an infinitesimal.




Figure 4: Shift of $f(x) = \frac{1}{2}x \sin(3x) + 2$.

Finally, we reiterate some limit theorems already stated for sequences, but now corresponding to functions. Interpreting the function limit as the Cauchy sequence definition of limits renders the proofs of these theorems trivial.

Theorem 4.4 (Bounds on Limits of Functions)

If the functions $f, g : E \rightarrow \mathbb{R}$ are such that

$$\lim_{x \rightarrow a} f(x) = A < B = \lim_{x \rightarrow a} g(x) \quad (445)$$

then there exists a deleted neighborhood $U_\delta(a)$ in E at each point of which $f(x) < g(x)$.

Theorem 4.5 (Squeeze Theorem for Limits of Functions)

Given the functions $f, g, h : E \subset \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(x) \leq g(x) \leq h(x) \text{ for all } x \in E \quad (446)$$

then,

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = C \implies \lim_{x \rightarrow a} g(x) = C \quad (447)$$

4.2 Asymptotic Behavior of Functions

Definition 4.5 (Little-O Notation)

The function $f : E \rightarrow \mathbb{R}$ is said to be **infinitesimal compared with the function** $g : E \rightarrow \mathbb{R}$ as $x \rightarrow a$, written (by abuse of notation) $f = o(g)$ as $x \rightarrow a$, if

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = 0 \quad (448)$$

or in other words, if f/g is an infinitesimal function as $x \rightarrow a$. Therefore, $f = o(1)$ as $x \rightarrow a$ means that f is infinitesimal as $x \rightarrow a$.^a

^aNote that writing $f = o(g)$ is again, an abuse of notation. $f = o(g)$ is really a shorthand way of writing that f is in the class of functions that is infinitesimal compared with the function g .

Intuitively, $f = o(g)$ means that the ratio between $f(x)$ and $g(x)$ will tend to infinity as $x \rightarrow a$ (this does not mean that f will be infinitely greater than g , however!).

Example 4.4 (Linear vs Quadratic)

For example, looking at the two functions $f(x) = x^2$ and $g(x) = x$, we have

1. $x^2 = o(x)$ as $x \rightarrow 0$ (since $\frac{x^2}{x} = x$ is infinitesimal as $x \rightarrow 0$)
2. $x = o(x^2)$ as $x \rightarrow \infty$ (since $\frac{x}{x^2} = \frac{1}{x}$ is infinitesimal as $x \rightarrow \infty$)

We can visualize $g/f(x)$ tending to infinity within a neighborhood of 0 and $f/g(x)$ tending to infinity within a neighborhood of ∞ .




Figure 5

Definition 4.6 (Orders of Infinitesimals, Infinities)

If $f = o(g)$ and g is infinitesimal as $x \rightarrow a$, then f is an **infinitesimal of higher order than g** as $x \rightarrow a$. Furthermore, if f and g are infinite functions as $x \rightarrow a$ and $f = o(g)$ as $x \rightarrow a$, then g is a **higher order infinity than f** as $x \rightarrow a$.

Definition 4.7 (Big-O Notation)

By abuse of notation, $f = O(g)$ as $x \rightarrow a$ means that

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \infty \quad (449)$$

or in other words, f/g is ultimately bounded as $x \rightarrow a$. In particular, $f = O(1)$ as $x \rightarrow a$ means that f is bounded within a certain neighborhood $U(a)$ of a .

Definition 4.8 (Functions of Same Order)

The functions f and g are of the same order as $x \rightarrow a$, written

$$f \asymp g \text{ as } x \rightarrow a \quad (450)$$

if $f = O(g)$ and $g = O(f)$ as $x \rightarrow a$. Intuitively, this means that the ratio between f and g within some deleted neighborhood of a is finite.

Note that the condition that f and g be of the same order as $x \rightarrow a$ is (by definition of ultimately bounded functions) equivalent to the condition that there exist $c_1, c_2 > 0$ and an open neighborhood $U(a)$ such that the relations

$$c_1|g(x)| \leq |f(x)| \leq c_2|g(x)| \quad (451)$$

is true for $x \in U(a)$.

Definition 4.9 (Asymptotic Equivalence of Functions)

For functions f and g , if

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = 1 \quad (452)$$

we say that f **behaves asymptotically like** g as $x \rightarrow a$, or that f **is equivalent to** g as $x \rightarrow a$, written

$$f \sim g \text{ as } x \rightarrow a \quad (453)$$

Moreover, \sim is an equivalence relation, which means that

1. $f \sim f$ as $x \rightarrow a$
2. $f \sim g$ as $x \rightarrow a \implies g \sim f$ as $x \rightarrow a$
3. $f \sim g$ and $g \sim h$ as $x \rightarrow a \implies f \sim h$ as $x \rightarrow a$

We list a few examples in order to develop some sort of visual intuition for when two functions are asymptotically equivalent.

Example 4.5 (Both Converges at Finite Value to Nonzero Finite Value)

If $f(a) = g(a) \neq 0$, then $f \sim g$ trivially since the ratio of f and g converges to 1 within a neighborhood of a .






Figure 6

Example 4.6 (Both Converges at Finite Value to 0)


When $f(a) = g(a) = 0$, it may be f may be equivalent to g or one function may be infinitesimally smaller than the other.




(a) When $f(x) = \sin x$ and $g(x) = x$, then $f \sim g$ since we see that $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$, and so $\sin x \sim x$ as $x \rightarrow 0$.



(b) When $f(x) = x^2$ and $g(x) = x^3$, then $\lim_{x \rightarrow 0} \frac{x^3}{x^2} = 0$, and so $x^3 \not\sim x^2$. In fact, $x^3 = o(x^2)$.



(c) When $f(x) = x^2$ and $g(x) = x^4$, then $\lim_{x \rightarrow 0} \frac{x^4}{x^2} = 0$, and so $x^4 \not\sim x^2$. In fact, $x^4 = o(x^2)$.



(d) When $f(x), g(x) = x^2, 0.5x^2$, then $\lim_{x \rightarrow 0} \frac{0.5x^2}{x^2} = \frac{1}{2}$. So $0.5x^2 \not\sim x^2$.

Figure 7: Examples of different scenarios.

Example 4.7 (Analyzing at Infinity)

When analyzing the behavior of functions as $x \rightarrow \infty$, we can picture the two graphs of f and g on the plane and "zoom out" to see if the ratio of the values converge to 1. This would mean that as $x \rightarrow \infty$, we should see the graphs overlapping more and more.




Figure 8: taking $f(x) = x^2$ and $g(x) = x^2 + 10x + 100$, we can see that the discrepancy is high around a neighborhood of $x = 0$. But as $x \rightarrow +\infty$, we get $\lim_{x \rightarrow +\infty} \frac{x^2 + 10x + 100}{x^2} = 1$, and so the graphs look like they are overlapping. Notice that even though the absolute difference $|(x^2 + 10x + 100) - x^2| = |10x + 100|$ tends to infinity, this difference increases infinitesimally compared to f and g .

From this, we can see that if $f \sim g$ as $x \rightarrow a$, then their difference

$$f - g = o(g) = o(f) \quad (454)$$

That is, $(f - g)(x)$ is infinitesimal compared to g or f (doesn't matter which one we compare it to). This leads to our next section, where we formalize this concept with absolute and relative errors.

4.2.1 Approximations of Functions

It is useful to note that since the relation $\lim_{x \rightarrow a} \gamma(x) = 1$ is equivalent to

$$\gamma(x) = 1 + \alpha(x), \text{ where } \lim_{x \rightarrow a} \alpha(x) = 0 \quad (455)$$

the relation $f \sim g$ as $x \rightarrow a$ is equivalent to saying that

$$\frac{f(x)}{g(x)} = \gamma(x), \text{ where } \lim_{x \rightarrow a} \gamma(x) = 1 \quad (456)$$


which implies

$$f(x) = g(x) + \alpha(x)g(x) = g(x) + o(g(x)) \text{ as } x \rightarrow a \quad (457)$$

or, symmetrically,

$$g(x) = f(x) + \alpha(x)f(x) = f(x) + o(f(x)) \text{ as } x \rightarrow a \quad (458)$$

This means that f can be exactly represented by another function g , plus another (error) function $o(g(x))$ that is infinitesimal compared to g .

Figure 9: Visualization of asymptotic behavior where $f - g = o(g)$

Note that it is not a sufficient condition that the error function be infinitesimal! The error function $f - g$ must be infinitesimal *compared to g* ! This tells us that not only does the error function decrease infinitesimally, but also is infinitesimal compared to the approximation function we already have, which is in general a much stronger claim. This representation of certain types functions will provide the foundation for differential calculus when we talk about "good" approximations for a function.

Definition 4.10 (Relative Error)

Since $f \sim g$ as $x \rightarrow a$ means that

$$f(x) = g(x) + \alpha(x)g(x) = g(x) + o(g(x)) \quad (459)$$

we can define the **relative error** of g as an approximation of f to be

$$|\alpha(x)| = \left| \frac{f(x) - g(x)}{g(x)} \right| \quad (460)$$

Clearly, since $f \sim g$, the relative error must be infinitesimal as $x \rightarrow a$.

We use the following lemma to check whether two functions are asymptotically equivalent.

Lemma 4.2 ()

$f \sim g$ as $x \rightarrow a$ if and only if the relative error of g is infinitesimal as $x \rightarrow a$.

Example 4.8 ()

We claim that

$$x^2 + x = \left(1 + \frac{1}{x}\right)x^2 \sim x^2 \text{ as } x \rightarrow \infty \quad (461)$$

We see that the absolute error of this approximation $|(x^2 + x) - x^2| = |x|$ tends to infinity, but the relative error $\frac{|x|}{x^2} = \frac{1}{|x|} \rightarrow 0$ as $x \rightarrow \infty$.

Theorem 4.6 (Prime Number Theorem)

Let $\pi(x)$ be the number of prime numbers strictly less than x . Then $\pi \sim \frac{x}{\ln x}$ as $x \rightarrow +\infty$, or more precisely,

$$\pi(x) = \frac{x}{\ln x} + o\left(\frac{x}{\ln x}\right) \text{ as } x \rightarrow +\infty \quad (462)$$

Example 4.9 ()

It is a fact that $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$, so we have $\sin x \sim x$ as $x \rightarrow 0$. So,

$$\sin x = x + o(x) \text{ as } x \rightarrow 0 \quad (463)$$

The following theorem proves useful when computing limits.

Theorem 4.7 ()

If $f \sim \tilde{f}$ as $x \rightarrow a$, then

$$\lim_{x \rightarrow a} f(x)g(x) = \lim_{x \rightarrow a} \tilde{f}(x)g(x) \quad (464)$$

provided one of these limits exist.

Theorem 4.8 (Properties of $o(g)$ and $O(g)$ Functions)

For $x \rightarrow a$,

1. $o(f) + o(f) = o(f)$
2. $o(f)$ is also $O(f)$
3. $o(f) + O(f) = O(f)$
4. $O(f) + O(f) = O(f)$
5. If $g(x) \neq 0$, then

$$\frac{o(f(x))}{g(x)} = o\left(\frac{f(x)}{g(x)}\right), \text{ and } \frac{O(f(x))}{g(x)} = O\left(\frac{f(x)}{g(x)}\right) \quad (465)$$

4.3 Continuous Functions**Definition 4.11 (Continuity of a Function)**

A function f is **continuous at point a** if for any neighborhood $V(f(a))$ of $f(a)$, there is a neighborhood $U(a)$ of a whose image under the mapping f is contained in $V(f(a))$.

Generalizing this, we say that a function is **(globally) continuous** if the preimage of every neighborhood in its codomain is an open set in its domain.

Lemma 4.3 (Existence of Limits of Continuous Functions)

$f : E \rightarrow \mathbb{R}$ is continuous at $a \in E$, where a is a limit point of E if and only if

$$\lim_{x \rightarrow a} f(x) = f(a) \quad (466)$$

Proof. The limit equaling $f(a)$ means that, by definition, for any arbitrarily small deleted neighborhood of $f(a)$, denoted $U_{f(a)} \setminus f(a)$, its preimage will be an open neighborhood of a , which itself will contain an open set.

This also means that we can use the Cauchy limit definition to defined continuity of a function at a point. That is, for any sequence $\{a_n\}$ of point in codomain E which converges to point a , the function f is continuous at a if the corresponding sequence $\{f(a_n)\}$ converges to $f(a)$.

Theorem 4.9 ()

This means that the continuous functions commute with the operation of passing to the limit at a point.

$$\lim_{x \rightarrow a} f(x) = f\left(\lim_{x \rightarrow a} x\right) \quad (467)$$

Lemma 4.4 (Properties of Continuous Functions)

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $g : \mathbb{R}^m \rightarrow \mathbb{R}^p$ with $c \in \mathbb{R}$.

1. f continuous at $x_0 \implies cf$ continuous at x_0 .
2. f, g continuous at $x_0 \implies f + g$ continuous at x_0 .
3. Let $m = 1$. f, g continuous at $x_0 \implies fg$ continuous at x_0 .
4. f continuous at x_0 and $f(x) \neq 0 \forall x \in \mathbb{R}^n \implies 1/f$ continuous at x_0 .
5. If $f(x) = (f_1(x), f_2(x), \dots, f_n(x))$ coordinate-wise, then

$$f \text{ continuous at } x_0 \iff f_1, f_2, \dots, f_m \text{ continuous at } x_0 \quad (468)$$

6. f continuous at x_0 and g continuous at $y_0 = f(x_0) \implies g \circ f$ continuous at x_0 .

Proof. This is an immediate result of the equivalence of a function being continuous at point a and its limit at point a existing.

Theorem 4.10 (Local Properties of Continuous Functions)

Let $f : E \rightarrow \mathbb{R}$ be a function that is continuous at the point $a \in E$. Then,

1. f is bounded in some neighborhood $U(a)$.
2. If $f(a) \neq 0$, then in some neighborhood $U(a)$ all the values of the function have the same sign as $f(a)$.
3. If the function $g : U(a) \subset E \rightarrow \mathbb{R}$ is defined in some neighborhood of a and is continuous at a , then the following functions

$$(f + g)(x) \quad (469)$$

$$(f \cdot g)(x) \quad (470)$$

$$\left(\frac{f}{g}\right)(x) \text{ where } g(a) \neq 0 \quad (471)$$

are also defined in $U(a)$ and continuous at a .

4. If the function $g : Y \rightarrow \mathbb{R}$ is continuous at a point $b \in Y$ and f is such that $f : E \rightarrow Y$, $f(a) = b$, and f is continuous at a , then the composite function

$$g \circ f : E \rightarrow \mathbb{R} \quad (472)$$

is defined on E and continuous at a . This is easy to see because given the open neighborhood of

$g(b)$, we know for a fact that $U_\delta(a)$ maps completely into $U_\epsilon(b)$, and that $U_\epsilon(b)$ maps completely into $U_\kappa(g(b))$ and so the composition of these mappings must mean that $U_\delta(a)$ maps completely into $U_\kappa(g(b))$.

Example 4.10 ()

An algebraic polynomial

$$P(x) = a_0x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_{n-1}x + a_n \quad (473)$$

is a continuous function on \mathbb{R} . Since $f(x) = x$ and $f(x) = c$ are continuous functions, by induction on x , we can multiply them together to find that $f(x) = x^n$ is continuous, which implies that ax^n is continuous, which implies that the sums of these functions are also continuous.

4.3.1 Intermediate and Extreme Value Theorem

Unlike local properties, the global property of a function is a property involving the entire domain of definition of the function.

Theorem 4.11 (Compact Sets to Compact)

If $f : X \rightarrow Y$ is continuous and $K \subset X$ is compact, then $f(K)$ is compact in Y .

Proof.

Corollary 4.1 (Extreme Value Theorem)

A continuous real-valued function over a compact set attains its maximum and minimum.

Theorem 4.12 (Intermediate Value Theorem)

If a function f is continuous on an open interval and assumes values $f(a) = A$, $f(b) = B$, then for any number $C \in (A, B)$, there is a point c between a and b such that $f(c) = C$.




Figure 10: Illustration of a continuous function with a root in the interval $[a, b]$

Proof.

This following proof provides a very simple algorithm for finding the zero of the equation $f(x) = 0$ on an interval whose endpoints has values with opposite signs. Note that the colloquial description of the intermediate value theorem, that is is impossible to pass continuously from positive to negative values without assuming the value 0 along the way), assumes more than they state. That is, this theorem is actually dependent on the domain of definition: that is is a closed interval, or more generally, that it is **connected**.

4.3.2 Inverse Function Theorem

We begin by introducing this intuitive lemma.

Lemma 4.5 ()

A continuous mapping $f : E \rightarrow \mathbb{R}$ of a closed interval $E = [a, b]$ into \mathbb{R} is injective if and only if the function f is strictly monotonic on $[a, b]$.

Furthermore, every strictly monotonic function $f : X \subset \mathbb{R} \rightarrow \mathbb{R}$ (for arbitrary X) has an inverse

$$f^{-1} : f(X) \subset \mathbb{R} \rightarrow \mathbb{R} \quad (474)$$

with the same kind of monotonicity on $f(X)$ that f has on X .


Lemma 4.6 (Criterion for Continuity of a Monotonic Function)

A monotonic function $f : E \rightarrow \mathbb{R}$ defined on a closed interval $E = [a, b]$ is continuous if and only if its set of values $f(E)$ is the closed interval with endpoints $f(a)$ and $f(b)$.

Note that both conditions imply that there are no points of discontinuities in the graph of f .

Theorem 4.13 (Inverse Function Theorem)

A function $f : X \rightarrow \mathbb{R}$ that is strictly monotonic on a set $X \subset \mathbb{R}$ has an inverse $f^{-1} : Y \rightarrow \mathbb{R}$ defined on the set $Y = f(X)$ of values of f . The function $f^{-1} : Y \rightarrow \mathbb{R}$ is monotonic and has the same type of monotonicity on Y that f has on X .



If in addition, X is a closed interval $[a, b]$ and f is continuous on X , then the set $Y = f(X)$ is the closed interval with endpoints $f(a)$ and $f(b)$ and the function $f^{-1} : Y \rightarrow \mathbb{R}$ is continuous on it.

Example 4.11 (Sin and Arcsin)

The function $f(x) = \sin x$ is increasing and continuous on the closed interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$. Hence, the restriction to the closed interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$ has an inverse $x = f^{-1}(y)$, called the **arcsin**, and denoted by $x = \arcsin y$.




Figure 11: This function is defined on the closed interval $[-\sin(-\frac{\pi}{2}), \sin(\frac{\pi}{2})] = [-1, 1]$ and increases continuously from $-\frac{\pi}{2}$ to $\frac{\pi}{2}$.

4.4 Uniform Continuity

Roughly speaking, a function f is uniformly continuous if it is possible to guarantee that $f(x)$ and $f(y)$ be as close to each other as we please by requiring only that x and y be sufficiently close to each other. Intuitively, uniform continuity says that given any two points x, y in the domain where their distance is arbitrarily small (δ apart), we can guarantee that the distance between $f(x), f(y)$ is at maximum some arbitrarily small ϵ .

Definition 4.12 (Uniform Continuity)


A function $f : E \rightarrow \mathbb{R}$ is **uniformly continuous** on a set $E \subset \mathbb{R}$ if for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|f(x_1) - f(x_2)| < \epsilon \quad (475)$$

for all points $x_1, x_2 \in E$ such that $|x_1 - x_2| < \delta$.


Example 4.12 (Uniformly Continuous)

The following visual shows the radical function $f(x) = \sqrt{x}$ defined on \mathbb{R}^+ . We can see that it satisfies uniform continuity because the graph does not escape the top and/or bottom of the $\epsilon \times \delta$ window, no matter where the box is located on the graph. More strictly speaking, no matter what we set the ϵ (how long the box is), uniform continuity says that we can choose a sufficient δ (width of the box) such that the graph does not escape the top/bottom of the window no matter where the window is.



Example 4.13 (Not Uniformly Continuous)


We can clearly see that the function $f(x) = 1/x$ is not uniformly continuous, since the graph escapes the $\epsilon \times \delta$ window at some point (marked in red). More strictly speaking, given any length ϵ of the window, we cannot create a thin-enough δ box that will contain the graph, since as $x \rightarrow 0$, the function becomes unbounded. That is, arbitrarily thin boxes don't help when the slope is arbitrarily steep.




To compare uniform continuity with regular continuity, we can adapt this alternate (yet equivalent interpretation): Let there exist function $f : E \rightarrow \mathbb{R}$. Given any $\epsilon > 0$, we can choose a $\delta > 0$ such that given any point $x \in E$ and $f(x)$, as long as a second point y is δ away from x , then $f(y)$ is ϵ away from $f(x)$. This visualization would lead to there being a $2\epsilon \times 2\delta$ window around point x . Therefore, given a certain $\epsilon > 0$, the way we choose δ is only dependent on ϵ , and so it must be a function of ϵ :

$$\delta = \delta(\epsilon) \quad (476)$$

However, in continuity, there just has to exist *some* δ -neighborhood of x such that its image is contained in the ϵ -neighborhood of $f(x)$.



(a) Uniform continuity means that the box above does not change dimensions no matter where the point is (hence, the name uniform).



(b) In continuity, there are no restrictions on the dimensions of this box. It just has to exist for every point, through a function of ϵ .

Figure 14: Uniform continuity vs continuity.

With this intuition, it is easy to see the result below.

Lemma 4.7 (Uniform Continuity Implies Continuity)

If f is uniformly continuous on the set E , it is continuous at each point of that set. However, the converse is not generally true.

Example 4.14 ()

Let $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = 3x + 7$. Then f is uniformly continuous. Choose $\epsilon > 0$. Let $\delta = \epsilon/3$. Choose $x, y \in \mathbb{R}$ and assume $|x - y| < \delta$. Then,

$$|f(x) - f(y)| = |3x + 7 - 3y - 7| = 3|x - y| < 3\delta = \epsilon \quad (477)$$

Example 4.15 ()

Let $f : (0, 4) \subset \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = x^2$. Then f is uniformly continuous on $(0, 4)$. Choose $\epsilon > 0$. Let $\delta = \epsilon/8$. Choose $x, y \in (0, 4)$ and assume $|x - y| < \delta$. Then,

$$|f(x) - f(y)| = |x^2 - y^2| = (x + y)|x - y| < (4 + 4)|x - y| = 8\delta = \epsilon \quad (478)$$

A natural question one might ask is: under what assumptions is the converse true?

Theorem 4.14 (Cantor's Theorem on Uniform Continuity)

A function that is continuous on a compact set is uniformly continuous on that set.

Proof. Here is a proof I derived myself. The general idea is that with continuity, we can control the δ with an ϵ plus some point z . To decouple this dependence on z , we can restrict our scope to each z that are the center of the balls in the finite cover of x , and then taking the minimum (most restrictive) over them.

We wish to show that $\forall \epsilon > 0, \exists \delta > 0$ s.t.

$$d(x, x') < \delta \implies d(f(x), f(x')) < \epsilon \quad \forall x, x' \in X \quad (479)$$

We begin by constructing an open cover. Fix $\epsilon > 0$. $\forall z \in X, \exists \delta_z > 0$ s.t.

$$d(z, x) < \delta_z \implies d(f(z), f(x)) < \epsilon \quad (480)$$

This gives us an open cover $\{B(\frac{\delta_z}{2}, z)\}_{z \in X}$ of X , and by compactness, let's take a finite cover $\{B(\frac{\delta_{z_i}}{2}, z_i)\}_{i=1}^n$.

Now set $\delta = \min_i \{\delta_{z_i}/2\}$. We hope that when we choose $x, y \in X$ s.t. $d(x, y) < \delta$, we can trivially assume that $x \in B_i$ for some i . By setting the radius to be half, we can also assume that y is also in the same ball as x since

$$d(z_i, y) \leq d(z_i, x) + d(x, y) \leq \frac{\delta_{z_i}}{2} + \min_i \{\delta_{z_i}/2\} \leq \delta_{z_i} \quad (481)$$

This pretty much means that we can use the continuity properties locally within B_i , and so we can bound

$$d(f(x), f(y)) \leq d(f(x), f(z_i)) + d(f(z_i), f(y)) \leq 2\epsilon \quad (482)$$

Proof. Here is another proof recommended by Prof. Kiselev, who mentioned that a sequential proof is what he would go with intuitively. Let's prove by contradiction.^a Assume f is not uniformly continuous. Then $\exists \epsilon_0 > 0$ s.t. $\forall \delta = \frac{1}{n} > 0$, there exists $x_n, y_n \in X$ s.t.

$$d(x, y) < \frac{1}{n} \text{ but } d(f(x_n), f(y_n)) \geq \epsilon_0 \quad (483)$$

Since X is compact, we can take a convergent subsequence (x_{n_k}) of (x_n) , which converges to say z . We can take the same subindex to define the corresponding (y_{n_k}) . We first show that these two subsequences converge to the same value. By triangle inequality, we see that

$$d(z, y_{n_k}) \leq d(z, x_{n_k}) + d(x_{n_k}, y_{n_k}) \quad (484)$$

As we take the limit as $k \rightarrow \infty$, we see that $d(z, x_{n_k}) \rightarrow 0$ by construction of our subsequence, and $d(x_{n_k}, y_{n_k}) < \frac{1}{n_k}$ by construction. Therefore, $d(z, y_{n_k}) \rightarrow 0$ and we establish convergence. By continuity of f , we can see that

$$\lim_{k \rightarrow +\infty} f(x_k) = f(z) = \lim_{k \rightarrow +\infty} f(y_k) \quad (485)$$

However, we have picked x_n, y_n —and thus x_{n_k}, y_{n_k} —such that $d(f(x_{n_k}), f(y_{n_k})) \geq \epsilon_0$, which is a contradiction to the equality above.

^aThanks to Kiselev. This is also Exercise 4.10 in Baby Rudin.

Example 4.16 (Wrong Proof Attempt to Reflect on My Actions)

Here's a wrong proof I had in my first attempt. Fix $\epsilon > 0$, and consider any two points $x, x' \in X$. Take $r = d(x, x') + 1$, and take the finite subcover of the open cover of all r -balls of X , giving us

$$\{B_r(z_i)\}_{i=1}^n \quad (486)$$

Let's focus on one ball, which we can assume contains both x, x' since we can just add it. Since we

know that f is continuous on z_i , we know that $\exists \delta'_i > 0$ s.t.

$$d(x, z_i) < \delta_i \implies d(f(x), f(z_i)) < \frac{\epsilon}{2} \quad (487)$$

$$d(x', z_i) < \delta_i \implies d(f(x'), f(z_i)) < \frac{\epsilon}{2} \quad (488)$$

Therefore, if we unfix x, x' , we can always find a ball $B_r(z_i)$ containing both of them and find a $\delta_i = 2\delta'_i$ such that

$$d(f(x), f(x')) \leq d(f(x), f(z_i)) + d(f(z_i), f(x')) < \epsilon \quad (489)$$

Theorem 4.15 (Uniformly Continuous Function is Linearly Bounded)

If $f : \mathbb{R} \rightarrow \mathbb{R}$ is uniformly continuous, then there exists constants, $a, b \in \mathbb{R}$ such that $|f(x)| \leq a + b|x|$ for all $x \in \mathbb{R}$.

Proof. Since f is uniformly continuous, for every $\epsilon > 0$ there exists a $\delta > 0$ s.t.

$$|x - y| < \delta \implies |f(x) - f(y)| < \epsilon \quad (490)$$

for all $x, y \in \mathbb{R}$. Then, setting $y = 0$ and $\epsilon = 1$, we have some $\delta > 0$ s.t.

$$|x| < \delta \implies |f(0) - f(x)| < 1 \quad (491)$$

Now for any $k \in \mathbb{N}$, take $|x| < k\delta$. Then we can construct a sequence $(x_i = i\frac{|x|}{k})_{i=0}^k$, where $x_0 = 0$ and $x_k = |x|$, and from the triangle inequality followed by uniform convergence, we have

$$|x_i - x_{i+1}| = \frac{|x|}{k} < \frac{k\delta}{k} = \delta \implies |f(0) - f(x)| \leq \sum_{i=0}^{k-1} |f(x_i) - f(x_{i+1})| < \sum_{i=0}^{k-1} 1 = k \quad (492)$$

and so we come to the result

$$|x| < k\delta \implies |f(0) - f(x)| < k \quad (493)$$

Now set $m(x) = \min\{k \in \mathbb{N} \mid |x| < k\delta\}$. It must be nonempty by the Archimedean property, so it's lower bounded by 1.

$$|x| < \delta m(x) \implies \frac{|x|}{\delta} < m(x) \implies m(x) \leq \frac{|x|}{\delta} + 1 \quad (494)$$

where the final implication comes from $m(x)$ being minimum. Therefore,

$$|f(x)| \leq |f(0)| + |f(x) - f(0)| \leq |f(0)| + m(x) \leq |f(0)| + \frac{|x|}{\delta} + 1 \quad (495)$$

and we have found such $a = |f(0)| + 1, b = \frac{1}{\delta}$.

4.5 Lipshitz and Holder Continuity

In both examples, the function satisfied an inequality of form

$$|f(x_1) - f(x_2)| \leq M|x_1 - x_2| \quad (496)$$

this is called the *Lipshitz inequality*. Lipshitz continuity is a strong form of uniform continuity for functions. Intuitively, a Lipshitz continuous function is limited in how fast it can change (by the Lipshitz constant).

Definition 4.13 (Lipshitz Continuous Function)

Given $f : E \subset \mathbb{R} \rightarrow \mathbb{R}$, f is **Lipshitz continuous** if there exists a $M > 0$ —called the *Lipshitz constant*—such that for all $x, y \in E$,

$$|f(x) - f(y)| \leq M|x - y| \quad (497)$$

Note that Lipshitz continuity pops up as a very natural extension of uniform continuity. The inequality above just means that given an ϵ , we can choose a δ such that a linear multiple of δ is always greater than ϵ . This means that Lipshitz continuity is just uniform continuity such that the δ function is linear:

$$\delta = \delta(\epsilon) = \frac{1}{M}\epsilon \quad (498)$$




Figure 15: Relationship between slope M and the ratio of δ to ϵ

Definition 4.14 (Bi-Lipshitz Continuity)

A function $f : E \subset \mathbb{R} \rightarrow \mathbb{R}$ is **Bi-Lipshitz continuous** if there exists constant $M \geq 1$ such that for all real $x, y \in E$,

$$\frac{1}{M}|x - y| \leq |f(x) - f(y)| \leq M|x - y| \quad (499)$$

It immediately follows that for $x \neq y$, $|f(x) - f(y)|$ cannot equal 0, which means that a bilipshitz map is injective. A bilipshitz map is really just Lipshitz map with its inverse also being Lipshitz.

Theorem 4.16 ()

A bilipshitz map f is a homeomorphism onto its image.

Definition 4.15 (Holder Continuity)

Given $f : E \subset \mathbb{R} \rightarrow \mathbb{R}$, f is **α -Holder continuous** if there exists a $C > 0$ such that for all $x, y \in E$,

$$|f(x) - f(y)| \leq C|x - y|^\alpha \quad (500)$$

4.6 Discontinuity

If the function $f : E \rightarrow \mathbb{R}$ is not continuous at a point of E , then this point is called a **point of discontinuity**, or simply a **discontinuity** of f . That is, a is a point of discontinuity of f if for some neighborhood $V(f(a))$ of $f(a)$, there exists no neighborhood of a whose image under the mapping f is contained in $V(f(a))$.

There are three types of discontinuities, ranging from least to most extreme.⁸

Definition 4.16 (Removable Discontinuity)

A **removable discontinuity** is characterized by the fact that the limit $\lim_{x \rightarrow a} f(x) = A$ exists, but $A \neq f(a)$.




Figure 16: A function with a removable discontinuity at $x = 1$. The function is defined as $f(x) = \frac{x^2-1}{x-1}$ for $x \neq 1$ and $f(1) = 1$. The limit of the function as x approaches 1 is 2 (shown by the open circle), but the function value at $x = 1$ is 1 (shown by the filled circle).

This means that we can modify f and define a new function $\tilde{f} : E \rightarrow \mathbb{R}$ as

$$\tilde{f}(x) = \begin{cases} f(x), & x \in E \setminus a \\ A, & x = a \end{cases} \quad (501)$$

which would be continuous on E .

Definition 4.17 (Jump/Step Discontinuity, of First Kind)

A **discontinuity of first kind**, also known as a jump/step discontinuity, is characterized by both the left and right-hand limits

$$\lim_{x \rightarrow a-0} f(x) \text{ and } \lim_{x \rightarrow a+0} f(x) \quad (502)$$

existing, but at least one of them is not equal to the value $f(a)$ that the function assumes at a .




Figure 17: A function with a step discontinuity at $x = 0$. The function is defined as $f(x) = 1 + 0.25x^2$ for $x < 0$ and $f(x) = 2 + 0.25x^2$ for $x \geq 0$. The limit from the left $\lim_{x \rightarrow 0-} f(x) = 1$ is shown by the open circle, while the function value at $x = 0$ is $f(0) = 2$ shown by the filled circle. The dashed line highlights the jump in value.


⁸Note that strictly speaking, a removable discontinuity is really a discontinuity of first kind, but in this context we distinguish them.

Definition 4.18 (Essential Discontinuity, of Second Kind)


A **discontinuity of second kind**, also known as an essential discontinuity, is characterized by at least one of the two limits

$$\lim_{x \rightarrow a-0} f(x) \text{ and } \lim_{x \rightarrow a+0} f(x) \quad (503)$$

not existing.



(a) The function $f(x) = \frac{1}{x}$ has an infinite discontinuity at $x = 0$



(b) The function $f(x) = |1.5 \sin(\frac{1}{x})|$ has an oscillatory discontinuity at $x = 0$

Figure 18: Examples of discontinuities of the second kind, where the limit does not exist as x approaches the point of discontinuity

Example 4.17 (Dirichlet Function)

The Dirichlet function, defined

$$\mathcal{D}(x) = \begin{cases} 1, & \text{if } x \in \mathbb{Q} \\ 0, & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases} \quad (504)$$

is discontinuous at every point, and obviously all of its discontinuities are of second kind, since in every interval there are both rational and irrational numbers and therefore there exists no limit at any point $a \in \mathbb{R}$.

More specifically, given any point $a \in \mathbb{R}$, assume that a is rational. We can set $\epsilon = 0.1$ -neighborhood around the value 1, but no matter how small we let δ , the interval $(a - \delta, a + \delta)$ will contain both rationals and irrationals, meaning that it will map to $\{0, 1\}$ always, which is not fully contained in $(0.9, 1.1)$.

Here is a slightly more interesting example.

Example 4.18 (Riemann Function)

Let the Riemann function \mathcal{R} be defined

$$\mathcal{R}(x) = \begin{cases} \frac{1}{n}, & \text{if } x = \frac{m}{n} \in \mathbb{Q}, \text{ where } \gcd(m, n) = 1 \\ 0, & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases} \quad (505)$$

We first note that for any point $a \in \mathbb{R}$, any bounded neighborhood $U(a)$ of it, and any number $N \in \mathbb{N}$, the neighborhood $U(a)$ contains only a finite number of rational numbers $\frac{m}{n}$, where $n < N$. By shrinking the neighborhood, we can assume that the denominators of all rational numbers in the neighborhood are larger than N , since rationals with larger denominators have smaller gaps between

them. Thus, at any point $x \in U(a) \setminus a$, we have

$$|\mathcal{R}(x)| < \frac{1}{N} \quad (506)$$

and therefore $\lim_{x \rightarrow a} \mathcal{R}(x) = 0$ at any point $a \in \mathbb{R} \setminus \mathbb{Q}$. Hence, the Riemann function is continuous at any irrational number.

4.7 Exercises

Exercise 4.1 (Math 531 Spring 2025, PS6.6)

Let X be a complete metric space. Suppose that $f : X \rightarrow X$ is such that $d(f(x), f(y)) \leq \frac{1}{2}d(x, y)$ for all $x, y \in X$.

- (a) Prove that f is continuous
- (b) Pick any $x_0 \in X$. Define a sequence of points $\{x_n\}$ in X by:

$$x_{n+1} = f(x_n). \quad (507)$$

Prove that $\{x_n\}$ converges. Hint: use a previous homework assignment.

- (c) Denote the limit of $\{x_n\}$ by x . Prove that $f(x) = x$.
- (d) Prove that if $f(y) = y$ and $f(x) = x$ then $x = y$.

Solution.

Exercise 4.2 (Math 531 Spring 2025, PS6.7)

Does there exist a continuous function $f : [0, 1] \rightarrow \mathbb{R}$ that is onto? If so, construct one from scratch. If not, prove such a function cannot exist.

Solution.

Exercise 4.3 (Math 531 Spring 2025, PS7.1)

If a function $f : [0, 1] \rightarrow \mathbb{R}$ is continuous, it is uniformly continuous. This means that, given $\epsilon > 0$, there exists $\delta(\epsilon) > 0$ so that

$$d(x, y) < \delta(\epsilon) \Rightarrow d(f(x), f(y)) < \epsilon. \quad (508)$$

- Prove that if $f(x) = 1$ for all x , we can take $\delta(\epsilon) = +\infty$.
- Prove that if $f(x) = Mx$, we can take $\delta(\epsilon) = \frac{\epsilon}{M}$.
- Prove that if $f(x) = \sqrt{x}$, we can take $\delta(\epsilon) = \frac{\epsilon^2}{4}$.
- Prove that if $f(x) = \frac{x}{x+1}$, we can take $\delta(\epsilon) = \epsilon$.
- Prove that if $f(x) = x^N$, for some $N \in \mathbb{N}$, then we can take $\delta(\epsilon) = \frac{\epsilon}{N}$.

You cannot use differentiation for Problem 1. Do everything from scratch.

Solution.

Exercise 4.4 (Math 531 Spring 2025, PS7.2)

Let X be a general metric space. A function $f : X \rightarrow Y$ is called compact if the image of every closed ball $B_r(x)$ is compact in Y . Prove that any continuous $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is compact. Give an example of a discontinuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ that is compact.

Solution.

Exercise 4.5 (Math 531 Spring 2025, PS7.3)

Prove that if $f : \mathbb{R} \rightarrow \mathbb{R}$ sends connected sets to connected sets and compact sets to compact sets, then it is continuous. Hint: Assume that $f : \mathbb{R} \rightarrow \mathbb{R}$ sends every connected set to a connected set. Assume also that f is discontinuous at some $x \in \mathbb{R}$. Find a compact set K so that $f(K)$ is not compact.

Solution.

Exercise 4.6 (Math 531 Spring 2025, PS7.4)

- Let X be a general metric space and assume $K \subset X$ is compact. Let $f : K \rightarrow K$ and assume that

$$d(f(x), f(y)) < d(x, y) \quad (509)$$

for all $x \neq y \in K$. Show that there exists $x \in K$ with $f(x) = x$.

- Find a function $f : [0, \infty) \rightarrow [0, \infty)$ with the property that $|f(x) - f(y)| < |x - y|$, for all non-equal $x, y \in [0, \infty)$, but for which there is no point $x \in [0, \infty)$ for which $f(x) = x$.

Hint for the first part: define the function $g : K \rightarrow \mathbb{R}$ by $g(x) = d(f(x), x)$. First prove that g is continuous. Then deduce that g must attain its minimum at some x^* . Then show $g(f(x^*)) < g(x^*)$ unless $f(x^*) = x^*$. Conclude that $f(x^*) = x^*$.

Solution.

Exercise 4.7 (Math 531 Spring 2025, PS7.6)

A function $f : X \rightarrow X$ is said to be Hölder continuous of degree α for some $\alpha \in (0, \infty)$ if there exists $M > 0$ so that

$$d(f(x), f(y)) \leq C \cdot d(x, y)^\alpha. \quad (510)$$

Prove that if f is Hölder continuous of degree $\alpha > 0$, then f is continuous. Give an example of a function on \mathbb{R} that is Hölder continuous of degree $\frac{1}{2}$ but not of degree 1. Prove that continuously differentiable functions $[0, 1]$ are Hölder continuous of degree 1. Prove that the only functions on \mathbb{R} that are Hölder continuous of degree larger than 1 are constants.

Solution.

5 Differentiation of Single-Variable Functions

In general, there are two ways that we define a derivative: as a limit and with infinitesimals. In a standard analysis course we use limits, but in differential equations physicists tend to use the language of infinitesimals. This is where our introduction of the hyperreals in smooth infinitesimal analysis (SIA) will be useful.

Definition 5.1 (Derivative Using Limits)

Given $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$, fix $x \in [a, b]$, and define the **difference quotient** as the following function of y .^a

$$\frac{f(x) - f(y)}{x - y} \quad (511)$$

If the following limit exists, we define its value—called the **derivative of f at x** —as

$$f'(x) := \lim_{y \rightarrow x} \frac{f(x) - f(y)}{x - y} \quad (512)$$

and say f is **differentiable** at x . If f is differentiable at x for every $x \in E$, then f is said to be **differentiable over E** .

^aOften, textbooks introduce the difference quotient as $\frac{f(x+h)-f(x)}{h}$. These two are equivalent definitions since the following two different quotients $\phi(y) = \frac{f(x)-f(y)}{x-y}$ and $\gamma(h) = \frac{f(x+h)-f(x)}{h}$ are related in the sense that $\phi(y) = \gamma(y-x)$. So the following two limits exist simultaneously (or fail to exist simultaneously). If they do both exist, then $\lim_{y \rightarrow x} \phi(y) = \lim_{y \rightarrow x} \gamma(y-x) = \lim_{y \rightarrow 0} \gamma(y) = \lim_{h \rightarrow 0} \gamma(h)$ where the only nontrivial equality is the second equality, which is true (should be shown).

So if f is differentiable, we can just treat it as a new function $g(x) = f'(x)$. An immediate consequence of the derivative is the following lemma, which states the existence of some error function φ that is “small” in a sense.

Lemma 5.1 (Fundamental Increment Lemma)

Suppose the derivative of $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ at x exists. Then, there exists a function $\varphi : (a, b) \subset \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(x+h) = f(x) + f'(x)h + \varphi(h)h \quad (513)$$

for sufficiently small nonzero h , and

$$\lim_{h \rightarrow 0} \varphi(h) = 0 \quad (514)$$

Proof. We can define

$$\varphi(h) = \frac{f(x+h) - f(x)}{h} - f'(x) \quad (515)$$

Is the converse is true? I think so. This gives us a new language to work with.

Definition 5.2 (Little-O Notation)

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be function and $g(h)$ is a monotonically increasing function with $g(0) = 0$. We say that $f(h)$ is $o(g(h))$ if

$$\lim_{h \rightarrow 0} \frac{f(h)}{g(|h|)} = 0 \quad (516)$$

More generally, $o(g(h))$ is a *class* of functions.


So, the φ in the fundamental increment lemma is really just describing an $o(h)$ function. This is a more powerful language since we can use little-O notation to extend to other classes of functions.

Theorem 5.1 (Differentiability with Little-o Notation)

A function $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ is differentiable at x if and only if there is a number $f'(x)$ s.t.

$$f(x+h) - f(x) = f'(x)h + o(h) \quad (517)$$

where by abuse of notation, we write $+o(h)$ to denote adding a function in $\varphi(h) \in o(h)$.^a



^aWe also use the *increment notation* to write $\Delta f(x) := f(x+h) - f(x)$ and $\Delta x := h$ to denote the difference in the input and outputs of f , leading to another familiar form: $\Delta f(x) = f'(x)\Delta x + o(h)$.

Proof.

In other words, the difference between the increment of the function and the value of the function $df(x)$ in h is an infinitesimal of higher order than h . For this reason, we say that the derivative is the *principal linear part of the increment of the function*.

In particular, if $f(x) := x$, then we have $f'(x) := 1$ and

$$dx(h) = 1 \cdot h = h \quad (518)$$

Substituting this equality into $df(x)(h) = f'(x)h$, we get

$$df(x)(h) = f'(x) dx(h) \quad (519)$$

or without the input parameter h ,

$$df(x) = f'(x) dx \quad (520)$$

Note that this is an equality between two functions of h . From this, we obtain the familiar **Leibniz notation** of the derivative:

$$\frac{df(x)(h)}{dx(h)} = f'(x) \iff \frac{df(x)}{dx} = f'(x) \quad (521)$$

That is, the function $\frac{df(x)}{dx}$, which is the ratio of the functions $df(x)$ and dx , is constant and equals $f'(x)$.

Let us try to construct successive approximations to an arbitrary function $f : E \rightarrow \mathbb{R}$ at a given limit point x_0 . That is, we find a function g such that

$$f = g + o(g) \quad (522)$$

Depending on what g is, we can construct better approximations of f .

Example 5.1 (Constant Approximation)

The 0th order approximation is when g is a constant. That is, $g := c_0$ for some $c_0 \in \mathbb{R}$. This means

$$f(x) = c_0 + o(c_0) = c_0 + o(1) \text{ as } x \rightarrow x_0 \quad (523)$$


More precisely, we want this difference $f(x) - c_0$ to be $o(1)$ as $x \rightarrow x_0$, which means that it is simply infinitesimal. Visualizing this, we can see that given a constant approximation (labeled in blue) to a function at x_0 , its error term (labeled in green) is in fact, infinitesimal. All this boils down to the fact that

$$\lim_{x \rightarrow x_0} f(x) = c_0 \quad (524)$$

If the function is continuous at x_0 , then

$$\lim_{x \rightarrow x_0} f(x) = f(x_0) \quad (525)$$

and naturally $c_0 = f(x_0)$. Both the continuous (left) and noncontinuous case (right) is shown, but in most cases, we will assume continuity.

**Example 5.2 (Linear Approximation)**

The 1st order approximation is a linear function that approximates f as

$$f(x) = c_0 + c_1(x - x_0) + o(x - x_0) \text{ as } x \rightarrow x_0 \quad (526)$$

Following the previous logic, assuming f continuous means that $c_0 = f(x_0)$. Furthermore, as $x \rightarrow x_0$

$$f(x) = c_0 + c_1(x - x_0) + o(x - x_0) \implies c_1 = \frac{f(x) - c_0 - o(x - x_0)}{x - x_0} \quad (527)$$

$$\implies c_1 = \frac{f(x) - c_0}{x - x_0} - \frac{o(x - x_0)}{x - x_0} \quad (528)$$

$$\implies c_1 = \frac{f(x) - c_0}{x - x_0} - o(1) \quad (529)$$

$$\implies c_1 = \lim_{x \rightarrow x_0} \frac{f(x) - c_0}{x - x_0} = f'(x_0) \quad (530)$$

But this just means that $f'(x_0) = c_1$. Note that before, we have proved the equivalence of the existence of a derivative at x_0 with differentiability at x_0 (which itself means that there exists a linear approximation $df(x)(h)$ that is a function of h). Here, we have created a linear approximation with respect to $x = x_0 + h$, rather than h (shifted the function).

Therefore, the function

$$\alpha(x) = f(x_0) + f'(x_0)(x - x_0) \quad (531)$$

provides the best linear approximation to the function f in a neighborhood of x_0 in the sense that for any other function $\beta(x)$ of the form

$$\beta(x) = c_0 + c_1(x - x_0) \quad (532)$$

we have $f(x) - \beta(x) \neq o(x - x_0)$ as $x \rightarrow x_0$. The graph of the function α is the straight line

$$y - f(x_0) = f'(x_0)(x - x_0) \quad (533)$$

This leads to the definition of our familiar tangent line.

Definition 5.3 (Tangent Line)


If a function $f : E \rightarrow \mathbb{R}$ is differentiable at a point $x_0 \in E$, the line defined by

$$y - f(x_0) = f'(x_0)(x - x_0) \quad (534)$$

is called the **tangent** to the graph of f at the point $(x_0, f(x_0))$.

Definition 5.4 (Tangent Space)

Given function $f : E \rightarrow \mathbb{R}$ and a point $x_0 \in E$, the increment of the argument $h = x - x_0$ can be regarded as a vector attached to the point x_0 and defining the transition from x_0 to $x_0 + h$. h is called a **tangent vector**, and the set of all such vectors as $T_{x_0}\mathbb{R}$. Similarly, we denote $T_{y_0}\mathbb{R}$ the set of all displacement vectors from the point y_0 along the y -axis.



Then, we can see that the differential is a mapping

$$df(x_0) : T_{x_0}\mathbb{R} \rightarrow T_{f(x_0)}\mathbb{R} \quad (535)$$

Note that there are two functions to pay attention to here:

1. The true increment of f , defined $h \mapsto f(x_0 + h) - f(x_0) = \Delta f(x_0; h)$ (labeled in green).
2. The differential $h \mapsto f'(x_0)h = df(x_0)(h)$, which gives the increment of the tangent to the graph for increment h in the argument (labeled in red).

Example 5.3 ()

Let $f(x) = \sin x$. Then we will show that $f'(x) = \cos x$.

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin(x)}{h} &= \lim_{h \rightarrow 0} \frac{2 \sin\left(\frac{h}{2}\right) \cos\left(x + \frac{h}{2}\right)}{h} \\ &= \lim_{h \rightarrow 0} \cos\left(x + \frac{h}{2}\right) \cdot \lim_{h \rightarrow 0} \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} = \cos(x) \end{aligned}$$

Here, we have used the theorem on the limit of a product, the continuity of the function $\cos(x)$, the equivalence $\sin t \sim t$ as $t \rightarrow 0$, and the theorem on the limit of a composite function.

Example 5.4 ()

We will show that $\cos'(x) = -\sin(x)$.

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{\cos(x+h) - \cos(x)}{h} &= \lim_{h \rightarrow 0} \frac{-2 \sin\left(\frac{h}{2}\right) \sin\left(x + \frac{h}{2}\right)}{h} \\ &= - \lim_{h \rightarrow 0} \sin\left(x + \frac{h}{2}\right) \cdot \lim_{h \rightarrow 0} \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} = -\sin(x) \end{aligned}$$

Lemma 5.2 (Differentiability Implies Continuity)

If f is differentiable at x , it is continuous at x .

Proof. If f is differentiable at x , then the derivative

$$f'(x) = \lim_{y \rightarrow x} \frac{f(x) - f(y)}{x - y} \quad (536)$$

exists. Therefore,

$$0 = f'(x) \cdot 0 = f'(x) \left(\lim_{y \rightarrow x} x - y \right) \quad (537)$$

$$= \lim_{y \rightarrow x} \frac{f(x) - f(y)}{x - y} \cdot \lim_{y \rightarrow x} x - y \quad (538)$$

$$= \lim_{y \rightarrow x} \frac{f(x) - f(y)}{x - y} (x - y) \quad (539)$$

$$= \lim_{y \rightarrow x} f(x) - f(y) \quad (540)$$

which implies that $f(x) = \lim_{y \rightarrow x} f(y)$, and hence f is continuous at x .

Therefore, we can see that the set of differentiable functions is a subset of the set of continuous functions.

5.1 Rules of Differentiation

Lemma 5.3 (Arithmetic)

If f and g are differentiable at x , then

1. $f + g$ is differentiable at x with

$$(f + g)'(x) = f'(x) + g'(x) \quad (541)$$

2. fg is differentiable at x with

$$(fg)' = f'(x)g(x) + f(x)g'(x) \quad (542)$$

3. f/g is differentiable at x with

$$\left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2} \quad (543)$$

Proof. The proof for addition is pretty trivial, so we will prove for multiplication and division. For products, let's not take the quotient just yet.

$$(fg)(x) - (fg)(y) = f(x)g(x) - f(y)g(y) \quad (544)$$

We know something about $f(x) - f(y)$ and $g(x) - g(y)$, so try to put it into this form.

$$(f(x) - f(y))g(x) + f(y)(g(x) - g(y)) \quad (545)$$

Therefore,

$$\frac{(fg)(x) - (fg)(y)}{x - y} = \underbrace{\frac{f(x) - f(y)}{x - y}}_{\text{exists}} g(x) + f(y) \underbrace{\frac{g(x) - g(y)}{x - y}}_{\text{exists}} \quad (546)$$

So by taking limits, f is continuous so $f(y) \rightarrow x$ as $y \rightarrow x$, and we finally have

$$f'(x)g(x) + f(x)g'(x) \quad (547)$$

For the quotient rule, it suffices to show from the product rule that $(1/g)'(x) = -\frac{g'(x)}{g(x)^2}$.

In proving properties of differentiability, it is useful to observe that

$$\lim_{y \rightarrow x} \frac{f(x) - f(y)}{x - y} \iff f(x) - f(y) = (x - y)(f'(x) + E) \quad (548)$$

for some function E where $\lim_{y \rightarrow x} E(x) = 0$. This is known as Taylor's formula. This is similar to the decomposition of sequences into a constant plus an infinitesimal sequence.

From this, we can find the derivative of polynomials.

Corollary 5.1 (Polynomial Derivatives)

The following are true.

1. The derivative of a constant function is 0
2. The derivative of the identity function $f(x) = x$ is 1.
3. The derivative of $f(x) = x^n$ is nx^{n-1} .
4. The derivative of a polynomial $f(x) = a_n x^n + \dots + a_0$ can then be found.

Proof.

Theorem 5.2 (Chain Rule)

Let $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function, fix $x \in [a, b]$, and assume differentiable $g : I \rightarrow \mathbb{R}$ where $f(x) \in I$. Then, $h = g \circ f$ is differentiable at x with the derivative

$$h'(x) = g'(f(x))f'(x) \quad (549)$$

Proof. We have

$$g(f(x)) - g(f(y)) = (f(x) - f(y))(g'(f(x)) + E_{f(y) \rightarrow f(x)}) \quad (550)$$

Now we divide by $x - y$.

$$\frac{g(f(x)) - g(f(y))}{x - y} = \frac{f(x) - f(y)}{x - y} \cdot (g'(f(x)) + E_{f(y) \rightarrow f(x)}) \quad (551)$$

Now if $x \rightarrow y$, then f is continuous, which implies $f(y) \rightarrow f(x)$ and so $E \rightarrow 0$, and so by taking this limit, the above evaluates to

$$f'(x) \cdot g'(f(x)) \quad (552)$$

We could have also done

$$\frac{g(f(x)) - g(f(y))}{x - y} = \frac{g(f(x)) - g(f(y))}{f(x) - f(y)} \cdot \frac{f(x) - f(y)}{x - y} \rightarrow g'(f(x)) \cdot f'(x) \quad (553)$$

Theorem 5.3 (Differentiation of Inverse Functions over \mathbb{R})

Let $E_1, E_2 \subset \mathbb{R}$, and $f : E_1 \rightarrow E_2$ and $f^{-1} : E_2 \rightarrow E_1$ be mutually inverse and continuous at points $x_0 \in E_1$ and $f(x_0) = y_0 \in E_2$. If f is differentiable at x_0 and $f'(x_0) \neq 0$, then f^{-1} also differentiable at the point y_0 , and

$$(f^{-1})^{-1}(y_0) = (f'(x_0))^{-1} \iff df^{-1}(y_0) = (df(x_0))^{-1} \quad (554)$$




Figure 19: Relationship between a function f and its inverse f^{-1} , showing how their derivatives are related

Note that if we knew in advance that f^{-1} was differentiable at y_0 (which is a stronger hypothesis), we can find immediately by the identity

$$(f^{-1} \circ f)(x) = x \quad (555)$$

and the theorem on the differentiation of a composite function that

$$(f^{-1})'(y_0) \cdot f'(x_0) = 1 \quad (556)$$

Note that if the hypothesis was satisfied, but $f'(x_0) = 0$, then f^{-1} would not be differentiable since it would have an undefined differential.




Figure 20: A function through three points and its inverse relation

5.2 Theorems of Differentiable Functions

Theorem 5.4 (Local Extrema of Differentiable Functions Have Vanishing Derivative)

Let $f : [a, b] \rightarrow \mathbb{R}$ and assume f has a local maximum at $c \in (a, b)$ with f differentiable at c . Then $f'(c) = 0$.

Proof. Let us pick two sequences—a left one and a right one—that converges to x from either side.

1. If $x > c$ (with x sufficiently close to c), then $f(c) \geq f(x)$ and so

$$\frac{f(c) - f(x)}{c - x} \leq 0 \implies f'(c) = \lim_{c-x} \frac{f(c) - f(x)}{c - x} \leq 0 \quad (557)$$

2. If $x < c$, then $f(c) \geq f(x)$ and so

$$\frac{f(c) - f(x)}{c - x} \geq 0 \implies f'(c) = \lim_{c-x} \frac{f(c) - f(x)}{c - x} \geq 0 \quad (558)$$

So $0 \leq f'(c) \leq 0 \implies f'(c) = 0$.

Note that it is generally not true that $f'(c) = 0$ if $c = a$ or $c = b$, i.e. at the endpoints.

Theorem 5.5 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is differentiable on (a, b) . Then, if $f(a) = f(b)$, then there exists a $c \in (a, b)$ such that $f'(c) = 0$.

Proof. Since f is continuous on $[a, b]$, it has to attain its global max and min values somewhere in $[a, b]$. If either is in (a, b) , then the derivative is 0. If max and min are attained on $\{a, b\}$, then since $f(a) = f(b)$, this implies that $f(x) = f(a)$ for all $x \in [a, b]$, which implies $f'(x) = 0$.

Theorem 5.6 (Mean Value Theorem)

Assume $f : [a, b] \rightarrow \mathbb{R}$ is differentiable. Then there exists a $c \in (a, b)$ for which

$$f'(c) = \frac{f(b) - f(a)}{b - a} \iff f'(c)(b - a) = f(b) - f(a) \quad (559)$$

Proof. Just use Rolle's on

$$g(x) = f(x) - \left[\frac{f(b) - f(a)}{b - a}(x - a) + f(a) \right] \quad (560)$$

which satisfies $g(a) = g(b) = 0$, and so there must exist some $c \in (a, b)$ such that $g'(c) = 0$, i.e.

$$f'(c) - \frac{f(b) - f(a)}{b - a} = 0 \implies f'(c) = \frac{f(b) - f(a)}{b - a} \quad (561)$$

Geometrically, this means that there exists a tangent line somewhere at $\zeta \in (a, b)$ that is parallel the secant line connecting the two points $(a, f(a))$ and $(b, f(b))$.

Some remarks:

1. Physically, if x is interpreted as time and $f(b) - f(a)$ as the amount of displacement over the time $b - a$ of a particle moving along the line, this theorem says that the velocity $f'(x)$ of the particle at some time $\zeta \in (a, b)$ is such that if the particle had moved with constant velocity $f'(\zeta)$ over the whole time interval, it would have been displaced by the same amount $f(b) - f(a)$. We call $f'(\zeta)$ the **average velocity** over the time interval $[a, b]$.
2. Note that the Mean Value Theorem is important in that it connects the increment of a function over a finite interval with the derivative of the function on that interval. Up to now, we have characterized only the local (infinitesimal) increment of a function in terms of the derivative or differential at a given point. MVT connects the increment of a function over a **finite** interval with the derivative of the function.

The MVT actually leads to multiple useful corollaries.

Theorem 5.7 (Derivative of a Monotonic Function)

Given function $f : [a, b] \rightarrow \mathbb{R}$ that is differentiable on (a, b) ,

$$\begin{aligned} f'(x) &> 0 \implies f \text{ is increasing} \\ f'(x) &\geq 0 \iff f \text{ is nondecreasing} \\ f'(x) &:= 0 \iff f \text{ is constant} \\ f'(x) &\leq 0 \iff f \text{ is nonincreasing} \\ f'(x) &< 0 \implies f \text{ is decreasing} \end{aligned}$$

Note the one-sided direction for the strict inequalities.^a The reverse implication is a bit weaker.

$$\begin{aligned} f \text{ is increasing} &\implies f'(x) \geq 0 \\ f \text{ is decreasing} &\implies f'(x) \leq 0 \end{aligned}$$

^aThink of the function $f(x) = x^3$, which is strictly increasing, but has derivative $f'(0) = 0$ at $x = 0$.

Proof. If $x_1 < x_2$ are two points of the interval, then the MVT

$$f(x_2) - f(x_1) = f'(\zeta)(x_2 - x_1) \quad (562)$$

shows that the sign of the left hand side must equal that of the right.

Corollary 5.2 (Derivative of a Constant Function)

A function that is continuous on a closed interval $[a, b]$ is constant on it if and only if its derivative equals 0 at every point of the interval $[a, b]$ or the open interval (a, b) .

Therefore, if the derivatives $f'_1(x)$ and $f'_2(x)$ of two functions $f_1(x)$ and $f_2(x)$ are equal on some interval (that is, $f'_1(x) = f'_2(x)$ on the interval), then the difference

$$(f_1 - f_2)(x) = f_1(x) - f_2(x) \quad (563)$$

is constant.

Proof. Given constant function f , the MVT equation

$$0 = f(x_2) - f(x_1) = f'(\zeta)(x_2 - x_1) \quad (564)$$

implies that $f'(\zeta) = 0$ for all $x_1, x_2 \in E$. It follows that by the arithmetic properties of the derivative, given two functions f_1, f_2 with the same derivative on an interval, the derivative of their difference $(f_1 - f_2)' = 0$, and therefore must be constant on that interval.

Theorem 5.8 (IVT For Derivatives)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a real and differentiable function and suppose $f'(a) < \lambda < f'(b)$. Then there exists $x \in (a, b)$ such that $f'(x) = \lambda$.^a

^ai.e. f doesn't have to be continuous, but it must have a middle value.

Proof. Let $g(x) = f(x) - \lambda x$. Then g is differentiable with $g'(x) = f'(x) - \lambda$. But this implies that

1. $g'(a) = f'(a) - \lambda < 0$, which implies that

$$\frac{g(t_1) - g(a)}{t_1 - a} < 0 \implies g(t_1) - g(a) < 0 \implies g(t_1) < g(a) \quad (565)$$

for some $t_1 > a$ sufficiently close to a .

2. $g'(b) = f'(b) - \lambda > 0$, which implies that

$$\frac{g(t_2) - g(b)}{t_2 - b} > 0 \implies g(t_2) - g(b) > 0 \implies g(t_2) > g(b) \quad (566)$$

for some $t_2 < b$ sufficiently close to b .

By the mean value theorem there exists $x \in (a, b)$ s.t. $g'(x) = 0 \implies f'(x) = \lambda$.

Corollary 5.3 (Derivatives Cannot Have Jump Discontinuities)

You can't have a jump discontinuity^a for derivatives.

^aAlso called a discontinuity of the first kind.

That is, the derivative of a differentiable function cannot “jump,” so it’s like the IVT of derivatives. However, it may as well have discontinuities of the second kind.

Example 5.5 (Derivative Might Jump if Not Differentiable)

A non-example is $f(x) = |x|$. It is not differentiable over $[-1, 1]$, and so we see a jump in the derivative.

The following theorem is a useful generalization of Lagrange’s theorem.

Theorem 5.9 (Cauchy’s Finite-Increment Theorem)

Let $x = x(t)$ and $y = y(t)$ be functions that are continuous on a closed interval $[\alpha, \beta]$ and differentiable on the open interval (α, β) . Then, there exists a point $\tau \in [\alpha, \beta]$ such that

$$x'(\tau)(y(\beta) - y(\alpha)) = y'(\tau)(x(\beta) - x(\alpha)) \quad (567)$$

If in addition $x'(t) \neq 0$ for each $t \in (\alpha, \beta)$, then $x(\alpha) \neq x(\beta)$ and we have the equality

$$\frac{y(\beta) - y(\alpha)}{x(\beta) - x(\alpha)} = \frac{y'(\tau)}{x'(\tau)} \quad (568)$$

5.3 Extrema and Concavity

Similarly, we can connect the concepts of extrema and derivatives.

Theorem 5.10 (First Derivative Test)


Let function $f : E \rightarrow \mathbb{R}$ be defined in a neighborhood $U(x_0)$ of point x_0 , which is continuous at x_0 and differentiable in $\mathring{U}(x_0)$, a deleted neighborhood of x_0 . (Note that this is broader hypothesis than just assuming that f be differentiable at x_0 .) Let

$$\mathring{U}^-(x_0) := \{x \in U(x_0) \mid x < x_0\}, \quad \mathring{U}^+(x_0) := \{x \in U(x_0) \mid x > x_0\} \quad (569)$$

That is, $\mathring{U}^-(x_0)$ is the left portion of $\mathring{U}(x_0)$ and $\mathring{U}^+(x_0)$ is the right portion of $\mathring{U}(x_0)$. Then,

1. $(x_0, f(x_0))$ is strict local minimum if $f'(x) < 0$ in $\mathring{U}^-(x_0)$ and $f'(x) > 0$ in $\mathring{U}^+(x_0)$.
2. $(x_0, f(x_0))$ is strict local maximum if $f'(x) > 0$ in $\mathring{U}^-(x_0)$ and $f'(x) < 0$ in $\mathring{U}^+(x_0)$.
3. $(x_0, f(x_0))$ has no extremum at x_0 if $f'(x) > 0$ in both $\mathring{U}^-(x_0), \mathring{U}^+(x_0)$, or if $f'(x) < 0$ in both $\mathring{U}^-(x_0), \mathring{U}^+(x_0)$.

Note that if there is a discontinuity at a point x_0 , then this theorem does not apply. For example, $(x_0, f(x_0))$ in the graph below is a local minimum, even though the derivatives to the left of x_0 are positive and those to the right of x_0 are negative (within neighborhood $U(x_0)$). Similarly, $(x_0, g(x_0))$ is a local maximum, even though the derivative to the left and to the right of x_0 are both positive.


Theorem 5.11 (2nd, n th Derivative Test)

Let function $f : E \rightarrow \mathbb{R}$ be defined on a neighborhood $U(x_0)$ of x_0 has derivatives of order up to n inclusive at x_0 . If its derivatives up to the $(n-1)$ th order vanishes

$$f'(x_0) = f''(x_0) = \dots = f^{(n-1)}(x_0) = 0 \quad (570)$$

but the n th derivative at x_0 does **not** vanish

$$f^{(n)}(x_0) \neq 0 \quad (571)$$

then

1. n is odd \implies there is no local extremum at x_0
2. n is even \implies there is a local extremum at x_0
 - (a) $f^{(n)}(x_0) > 0 \implies$ it is a strict local minimum
 - (b) $f^{(n)}(x_0) < 0 \implies$ it is a strict local maximum

Definition 5.5 (Convex, Concave Functions)

A function $f : (a, b) \rightarrow \mathbb{R}$ defined on an open interval $(a, b) \subset \mathbb{R}$ is **convex** if the inequality

$$f(\alpha_1 x_1 + \alpha_2 x_2) \leq \alpha_1 f(x_1) + \alpha_2 f(x_2) \quad (572)$$

holds and **concave**, or **convex upward**, if the inequality

$$f(\alpha_1 x_1 + \alpha_2 x_2) \geq \alpha_1 f(x_1) + \alpha_2 f(x_2) \quad (573)$$

holds for all pairs of points $x_1, x_2 \in (a, b)$ and any numbers $\alpha_1, \alpha_2 \geq 0$ such that $\alpha_1 + \alpha_2 = 1$. If this inequality is strict whenever $x_1 \neq x_2$ and $\alpha_1 \alpha_2 \neq 0$, the function is said to be **strictly convex** and **strictly concave**, respectively.

Note that using induction on the number of points, we get a primitive form of Jensen's inequality.

Lemma 5.4 (Jensen's Inequality)

If $f : (a, b) \rightarrow \mathbb{R}$ is a convex function, x_1, \dots, x_n are points of (a, b) , and $\alpha_1, \dots, \alpha_n$ are nonnegative numbers such that $\alpha_1 + \dots + \alpha_n = 1$, then

$$f(\alpha_1 x_1 + \dots + \alpha_n x_n) \leq \alpha_1 f(x_1) + \dots + \alpha_n f(x_n) \quad (574)$$

The following is also another equivalent condition for a function to be convex over (a, b) .

Theorem 5.12 ()

A function $f : (a, b) \rightarrow \mathbb{R}$ that is differentiable on the open interval (a, b) is convex on (a, b) if and only if its graph contains no points below any tangent drawn to it.

Theorem 5.13 (2nd Derivatives of Convex Functions)

Given a function $f : (a, b) \rightarrow \mathbb{R}$ that is differentiable in its domain,

1. f is convex $\iff f'$ is nondecreasing on $(a, b) \iff f'' \geq 0$ on (a, b)
2. f is strictly convex $\iff f'$ is increasing on $(a, b) \iff f'' > 0$ on (a, b)
3. f is concave $\iff f'$ is nonincreasing on $(a, b) \iff f'' \leq 0$ on (a, b)
4. f is strictly concave $\iff f'$ is decreasing on $(a, b) \iff f'' < 0$ on (a, b)

Definition 5.6 (Inflection Point)

Let $f : E \rightarrow \mathbb{R}$ be a function defined and differentiable on a neighborhood $U(x_0)$. If the function is convex downward (resp. upward) on the set $\mathring{U}^-(x_0) = \{x \in U(x_0) \mid x < x_0\}$ and convex upward (resp. downward) on $\mathring{U}^+(x_0) = \{x \in U(x_0) \mid x > x_0\}$, then the point

$$(x_0, f(x_0)) \quad (575)$$

is called a **inflection point of the graph**.




Figure 21: Curve with changing concavity and inflection point at π

5.4 Theorems of Continuously Differentiable Functions

Now continuously differentiable functions are called *smooth functions*, denoted $f \in C^1([a, b])$.

Theorem 5.14 (C^1 Implies Lipschitz)

A continuously differentiable function is Lipschitz continuous.

Proof.

Example 5.6 ()

Suppose f is twice-differentiable on \mathbb{R} and that M_0, M_1, M_2 are the least upper bounds of $|f(x)|$, $|f'(x)|$, and $|f''(x)|$. Then $M_1^2 \leq 4M_0M_2$.

Theorem 5.15 (L'Hopital's Rule)

Suppose f, g are continuously differentiable functions with $f(c) = g(c) = 0$ and $g'(c) \neq 0$. Then,

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)} \quad (576)$$

Proof. We have

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{g(x) - g(c)} \quad (577)$$

$$= \lim_{x \rightarrow c} \frac{\frac{f(x) - f(c)}{x - c}}{\frac{g(x) - g(c)}{x - c}} \quad (578)$$

$$= \frac{\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}}{\lim_{x \rightarrow c} \frac{g(x) - g(c)}{x - c}} \quad (579)$$

$$= \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)} \quad (580)$$

Example 5.7 ()

Let $f(x) = \sin x$ and $g(x) = -0.5x$. Then, the function

$$h(x) = \frac{f(x)}{g(x)} = \frac{\sin x}{-0.5x} \quad (581)$$

is clearly undefined at $x = 0$. However, we can solve the limit using L'Hopital's rule to get

$$\lim_{x \rightarrow 0} \frac{\sin x}{-0.5x} = \lim_{x \rightarrow 0} \frac{\cos x}{-0.5} = -2 \quad (582)$$

Therefore, $h : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ can be completed to continuous function on all of \mathbb{R} by defining the extension:

$$H(x) := \begin{cases} h(x), & x \neq 0 \\ -2, & x = 0 \end{cases} \quad (583)$$

5.5 Higher Order Derivatives

Note that the mean value theorem states that given differentiable $f : [a, b] \rightarrow \mathbb{R}$, there exists a $c \in (a, b)$ s.t.

$$f(b) = f(a) + f'(c)(b - a) \quad (584)$$

which is like a first order approximation. We would like to attain a second order approximation using the fact that f is twice differentiable. To do this, recall how we proved the MVT. We subtracted a linear function from $f(x)$ to get a new function $g(x)$ satisfying $g(a) = g(b) = 0$. We will do the same here.

Theorem 5.16 (Taylor's Theorem of 2nd Order)

If f is twice differentiable on $[a, b]$, then

$$f(b) = f(a) + f'(a)(b - a) + \frac{f''(c)}{2}(b - a)^2 \quad (585)$$

for some $c \in (a, b)$. This is like a mean value theorem for the second order, where only the final term is dependent on c .

Proof. Let us define the function

$$g(x) := f(x) - f(a) - f'(a)(x - a) - M(x - a)^2 \quad (586)$$

where M was chosen such that $g(b) = 0$. Now notice that

$$g(a) = f(a) - f(a) - 0 - 0 = 0 \quad (587)$$

$$g'(a) = f'(a) - f'(a) - 0 = 0 \quad (588)$$

and so by using Rolle's theorem on g , there exists a $c_1 \in (a, b)$ s.t.

$$g'(c_1)(b - a) = g(b) - g(a) = 0 - 0 = 0 \implies g'(c_1) = 0 \quad (589)$$

therefore, we can use Rolle's theorem again on g' and claim there exists a $c \in (a, c_1)$ s.t.

$$g''(c)(c_1 - a) = g'(c_1) - g'(a) = 0 - 0 = 0 \implies g''(c) = 0 \quad (590)$$

This gives us all we need. By taking the double derivative of g , we get

$$0 = g''(c) = f''(c) - 2M \implies M = \frac{f''(c)}{2} \quad (591)$$

and substituting this in gives

$$0 = g(b) = f(b) - f(a) - f'(a)(b - a) - \frac{f''(c)}{2}(b - a)^2 \quad (592)$$

We can continue this process to get a n th order approximation.

Theorem 5.17 (Taylor's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$, is n th differentiable over $[a, b]$ and $f^{(n+1)}$ exists over (a, b) . Then there exists a $c \in (a, b)$ s.t.

$$f(b) = \left(\sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (b - a)^k \right) + \frac{f^{(n+1)}(c)}{(n+1)!} (b - a)^{n+1} \quad (593)$$

Where

$$\varepsilon_f := \quad (594)$$

Proof. We do the exact same process. Let us define

$$P(x) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k \quad (595)$$

and set

$$g(x) := f(x) - P(x) - M(x-a)^{n+1} \quad (596)$$

where M was chosen such that $g(b) = 0$. Now notice that evaluating on g gives us

$$g(a) = g'(a) = g''(a) = \dots = g^{(n)}(a) = 0 \quad (597)$$

So by using Rolle's theorem on g , there exists a $c_1 \in (a, b)$ s.t. $g'(c_1) = 0$. Therefore we can use Rolle's theorem on g' to show there exists a $c_2 \in (a, c_1)$. We keep doing this until we show that there exists a $c \in (a, b)$ s.t. $g^{(n)}(c) = 0$. With this, we can directly evaluating the n th derivative of g to find

$$0 = g^{(n)}(c) = f^{(n)}(c) - n!M \implies M = \frac{f^{(n)}(c)}{n!} \quad (598)$$

We can continue this pattern to get a quadratic approximation of f in the form

$$f(x) = c_0 + c_1(x-x_0) + c_2(x-x_0)^2 + o((x-x_0)^2) \text{ as } x \rightarrow x_0 \quad (599)$$

As we have done in the previous subsection, we can derive (assuming continuity of f) $c_0 = f(x_0)$, $c_1 = f'(x_0)$. To derive what c_2 should be, we see that the equation above implies

$$c_2 = \frac{f(x) - c_0 - c_1(x-x_0) - o((x-x_0)^2)}{(x-x_0)^2} = \frac{f(x) - c_0 - c_1(x-x_0)}{(x-x_0)^2} - o(1) \quad (600)$$

which means

$$c_2 = \lim_{x \rightarrow x_0} \frac{f(x) - c_0 - c_1(x-x_0)}{(x-x_0)^2} \quad (601)$$

Extending this, if we are seeking a polynomial $P_n(x_0; x) = c_0 + c_1(x-x_0) + \dots + c_n(x-x_0)^n$ such that

$$f(x) = c_0 + c_1(x-x_0) + \dots + c_n(x-x_0)^n + o((x-x_0)^n) \text{ as } x \rightarrow x_0 \quad (602)$$

we would find

$$c_0 = \lim_{x \rightarrow x_0} f(x) \quad (603)$$

$$c_1 = \lim_{x \rightarrow x_0} \frac{f(x) - c_0}{x - x_0} \quad (604)$$

$$c_2 = \lim_{x \rightarrow x_0} \frac{f(x) - c_0 - c_1(x-x_0)}{(x-x_0)^2} \quad (605)$$

$$\dots = \dots \quad (606)$$


$$c_n = \lim_{x \rightarrow x_0} \frac{f(x) - (c_0 + \dots + c_{n-1}(x-x_0)^{n-1})}{(x-x_0)^n} \quad (607)$$

We formalize the order of these approximations by analyzing their error bound.

Definition 5.7 (nth Order Contact)

If $f, g : E \rightarrow \mathbb{R}$ are continuous at point x_0 and $(f - g)(x) = o((x - x_0)^n)$ as $x \rightarrow x_0$, then we say that f and g have ***nth order contact at x_0*** , or more precisely, ***contact of order at least n*** .

The following visual shows approximations g of an arbitrary function f that have 0th (left), 1st (middle), and 2nd (right) order contact at x_0 .

**Lemma 5.5 (Leibniz' Formula)**

Let $u(x)$ and $v(x)$ be functions having derivatives up to order n inclusive on a common set E . Then,

$$(uv)^{(n)} = \sum_{m=0}^n \binom{n}{m} u^{(n-m)} v^{(m)} \quad (608)$$

This means that given a polynomial $P_n(x) = c_0 + c_1(x - x_0) + \dots + c_n(x - x_0)^n$, then

$$P_n(x_0) = 0 \quad (609)$$

$$P'_n(x_0) = 1!c_1 \quad (610)$$

$$P''_n(x_0) = 2!c_2 \quad (611)$$

$$\dots = \dots \quad (612)$$

$$P_n^{(n)}(x_0) = n!c_n \quad (613)$$

$$P_n^{(k)}(x_0) = 0 \text{ for } k > n \quad (614)$$

and thus the polynomial $P_n(x)$ can be written as

$$P_n(x) = P_n^{(0)}(x_0) + \frac{1}{1!}P_n^{(1)}(x_0)(x - x_0) + \frac{1}{2!}P_n^{(2)}(x_0)(x - x_0)^2 + \dots + \frac{1}{n!}P_n^{(n)}(x_0)(x - x_0)^n \quad (615)$$

5.5.1 Taylor's Formula

From the following results one may deduce that the more derivatives of two functions coincide (including the derivative of the 0th order) at a point, the better these functions approximate each other in a neighborhood of that point. Using Leibniz's rule, approximations up to a certain degree at a point can be expressed as a polynomial

$$P_n(x_0; x) = P_n(x_0) + \frac{P'_n(x_0)}{1!}(x - x_0) + \dots + \frac{P_n^{(n)}(x_0)}{n!}(x - x_0)^n \quad (616)$$

where each coefficient of the polynomial

Definition 5.8 (Taylor Polynomial)

If a function $f : E \rightarrow \mathbb{R}$ has derivatives of all orders $n \in \mathbb{N}$ at a point x_0 , the unique series

$$P_n(x_0; x) = f(x_0) + \frac{f'(x_0)}{1!}(x - x_0) + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n \quad (617)$$

is the **Taylor polynomial of order n of $f(x)$ at x_0** . We can see that the derivatives of f and P_n coincide up to order n .

Definition 5.9 (Analytic Functions)

We cannot assume that the Taylor series of an infinitely differentiable function converges to the function f within a neighborhood $U(x_0)$, nor can we assume that it converges at all! These types of "nice" functions that have a Taylor approximation within the neighborhood of x_0 are called **analytic functions** and can be written in the form

$$f(x) = f(x_0) + \frac{f'(x_0)}{1!}(x - x_0) + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n + r_n(x_0; x) \quad (618)$$

where r is called the **remainder term**.

Example 5.8 (Infinitely Differentiable, Non-Analytic Function)

A example of a non-analytic function is

$$f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases} \quad (619)$$

which looks like the following.





Figure 22: Graph of the function $y = e^{-\frac{1}{x^2}}$. This function equals 0 at $x = 0$ and approaches 1 as $|x|$ approaches infinity. One can verify that the derivative $f^{(k)}(0) = 0$ for all k and hence the Taylor series is identically equal to 0, while $f(x) \neq 0$ if $x \neq 0$.

The relationship between these different conditions is nicely summarized in the figure.



The following lemma proves why Taylor Polynomials are considered a "good" approximations to analytic functions.

Lemma 5.6 (Infinitesimality of Functions with Vanishing Derivative up to Order n)

Given a function $\varphi : E \rightarrow \mathbb{R}$ defined on a closed interval E with endpoint x_0 , let its derivatives vanish up to order n at x_0 . That is

$$\varphi(x_0) = \varphi'(x_0) = \dots = \varphi^{(n)}(x_0) = 0 \quad (620)$$

Then, $\varphi = o((x - x_0)^n)$ as $x \rightarrow x_0$.

Proof. We prove by induction. For $n = 1$, the definition of differentiability states that

$$\varphi(x) = \varphi(x_0) + \varphi'(x - x_0) + o(x - x_0) \text{ as } x \rightarrow x_0 \quad (621)$$

and so we have proved that

$$\varphi(x_0) = \varphi'(x_0) = 0 \implies \varphi(x) = o(x - x_0) \text{ as } x \rightarrow x_0 \quad (622)$$

Now, suppose this assertion has been proved for order $n = k - 1 \geq 1$. That is, we have shown that

$$\varphi(x_0) = \dots = \varphi^{(k-1)}(x_0) = 0 \implies \varphi = o((x - x_0)^{k-1}) \text{ as } x \rightarrow x_0 \quad (623)$$

Then we must show that this is valid for order $n = k \geq 2$. Assume that

$$\varphi(x_0) = \varphi'(x_0) = \dots = \varphi^{(k)}(x_0) = 0 \quad (624)$$

We can see that this is equivalent to

$$(\varphi')'(x_0) = (\varphi')^{(2)}(x_0) = \dots = (\varphi')^{(k-1)}(x_0) = 0 \quad (625)$$

and therefore by the induction assumption, we have

$$\varphi' = o((x - x_0)^{k-1}) \text{ as } x \rightarrow x_0 \quad (626)$$

which means that we can put it in form

$$\varphi(x) = \alpha(x)(x - x_0)^{k-1} \text{ so that } \lim_{x \rightarrow x_0} \varphi(x) = \lim_{x \rightarrow x_0} \alpha(x) = 0 \quad (627)$$

From the mean value theorem and substituting what we have above, we get

$$\varphi(x) = \varphi(x) - \varphi(x_0) = \varphi'(\zeta)(x - x_0) \quad (628)$$

$$= \varphi(\zeta)(\zeta - x_0)^{k-1}(x - x_0) \quad (629)$$

where $\zeta \in (x_0, x)$. However, this implies that $|\zeta - x_0| < |x - x_0|$, and thus, as $x \rightarrow x_0$, $\zeta \rightarrow x_0$, which then makes $\alpha(\zeta) \rightarrow 0$. Since

$$|\varphi(x)| \leq |\alpha(\zeta)||x - x_0|^{k-1}|x - x_0| = |\alpha(\zeta)||x - x_0|^k \quad (630)$$

This means that $\varphi(x)$ is bounded by function $|\alpha(\zeta)||x - x_0|^k$, which is $o((x - x_0)^k)$, and so

$$\varphi = o((x - x_0)^k) \text{ as } x \rightarrow x_0 \quad (631)$$

By induction, this works for all orders n .

Theorem 5.18 (Peano's Form of the Remainder)

Given analytic function $f : E \rightarrow \mathbb{R}$, a point $x_0 \in E$, and its n th order Taylor polynomial $P_n(x_0; x)$ around x_0 , P_n is a "good" approximation of f in the fact that its error term is $o((x - x_0)^n)$. That is,

$$f(x) = P_n(x_0; x) + o((x - x_0)^n) \text{ as } x \rightarrow x_0 \quad (632)$$

This equation where $r_n(x; x_0) = o((x - x_0)^n)$ is called the **Peano's form of the remainder**.

Proof. Since the Taylor polynomial $P_n(x_0; x)$ is constructed from the requirement that its derivatives up to order n inclusive must coincide with the corresponding derivatives of f at x_0 , it follows that

$$r_n(x_0; x_0) := f^{(k)}(x_0) - P_n^{(k)}(x_0; x_0) = 0 \text{ for } k = 0, 1, \dots, n \quad (633)$$

Using the previous lemma, at this means that $r_n(x; x_0) = o((x - x_0)^n)$ as $x \rightarrow x_0$.

Theorem 5.19 (Lagrange Form of the Remainder)

If $f : E \rightarrow \mathbb{R}$ has derivatives of order $n + 1$ on the open interval with endpoints x_0 and x , then

$$f(x) = f(x_0) + \frac{f'(x_0)}{1!}(x - x_0) + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n + r_n(x; x_0) \quad (634)$$

where

$$r_n(x; x_0) = \frac{f^{(n+1)}(\zeta)}{(n+1)!}(x - x_0)^{n+1} \quad (635)$$

This form is called **Taylor's formula with the Lagrange form of the remainder**. Furthermore, this form says that if $f^{(n+1)}(x)$ is bounded in a neighborhood of x_0 , it also implies the formula

$$f(x) = f(x_0) + \frac{f'(x_0)}{1!}(x - x_0) + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n + O((x - x_0)^{n+1}) \quad (636)$$

Therefore, we can use this boundedness of $f^{(n+1)}$ to find the maximum error bound

$$|r_n(x; x_0)| \quad (637)$$

of $P_n(x; x_0)$.

Proof. It is a direct result from the lemma. This is actually a generalization of the mean value theorem but for higher orders.

Corollary 5.4 (Table of Asymptotic Formulas for Elementary Functions)

We write the Maclaurin series (Taylor series around $x = 0$) for elementary functions. Note that these

error terms are $O(x^{n+1})$ (bounded compared to x^{n+1}) and $o(x^n)$ (infinitesimal compared to x^n).

$$e^x = 1 + \frac{1}{1!}x + \frac{1}{2!}x^2 + \dots + \frac{1}{n!}x^n + O(x^{n+1}) \quad (638)$$

$$\cos x = 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \dots + \frac{(-1)^n}{(2n)!}x^{2n} + O(x^{2n+2}) \quad (639)$$

$$\sin x = x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \dots + \frac{(-1)^n}{(2n+1)!}x^{2n+1} + O(x^{2n+3}) \quad (640)$$

$$\cosh x = 1 + \frac{1}{2!}x^2 + \frac{1}{4!}x^4 + \dots + \frac{1}{(2n)!}x^{2n} + O(x^{2n+2}) \quad (641)$$

$$\sinh x = x + \frac{1}{3!}x^3 + \frac{1}{5!}x^5 + \dots + \frac{1}{(2n+1)!}x^{2n+1} + O(x^{2n+3}) \quad (642)$$

$$\ln(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \dots + \frac{(-1)^n}{n}x^n + O(x^{n+1}) \quad (643)$$

$$(1+x)^\alpha = 1 + \frac{\alpha}{1!}x + \frac{\alpha(\alpha-1)}{2!}x^2 + \dots + \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!}x^n + O(x^{n+1}) \quad (644)$$

5.6 Exercises

Exercise 5.1 (Math 531 Spring 2025, PS7.5)

Coming back to Problem 1 above, assume that f is differentiable on $[0, 1]$ and that $|f'(x)| \leq M$ for every $x \in [0, 1]$. Prove that we can take $\delta(\epsilon) = \frac{\epsilon}{M}$.

Solution.

Exercise 5.2 (Math 531 Spring 2025, PS7.7)

Suppose that $f : [0, 2] \rightarrow [0, 2]$ is twice differentiable. Suppose $f(0) = 0$, $f(1) = 1$, and $f(2) = 2$. Prove that there exists $c \in (0, 2)$ so that $f''(c) = 0$.

Solution.

Exercise 5.3 (Math 531 Spring 2025, PS7.8)

Suppose that f is differentiable on $[0, 1]$ satisfies:

$$f'(x) = f(x). \quad (645)$$

- Prove that f is automatically infinitely differentiable.
- Let $M = \max\{f(x) : x \in [0, 1]\}$. Why does M exist? Similarly, let $M(\epsilon) = \max\{|f(x)| : x \in [0, \epsilon]\}$.
- Prove that for all $x \in [0, \epsilon]$, we have that

$$|f(x)| \leq M(\epsilon)x + |f(0)|. \quad (646)$$

- Deduce that if $f(0) = 0$, we have that

$$M\left(\frac{1}{2}\right) = 0. \quad (647)$$

Similarly, deduce that $M(c) = 0$ for all $c \in [0, 1]$.

- What you have just proved is that if $f'(x) = f(x)$ while $f(0) = 0$, it follows that $f(x) = 0$ for all x .
- Assume that $E : \mathbb{R} \rightarrow [0, \infty)$ is differentiable and

$$|E'(t)| \leq CE(t). \quad (648)$$

Prove that if $E(0) = 0$, then $E(t) = 0$ for all t .

Solution.

Exercise 5.4 (Math 531 Spring 2025, PS8.1)

Prove that if $f : [0, 1] \rightarrow \mathbb{R}$ is differentiable and $f' > 0$ on $[0, 1]$, then f is strictly increasing on $[0, 1]$.

Solution.

Exercise 5.5 (Math 531 Spring 2025, PS8.2)

Explain why if $x(t)$ represents the position of a particle at time t , $x'(t)$ is called the velocity of the particle, $x''(t)$ is called its acceleration, and $x'''(t)$ is called its jerk.

Solution.

Exercise 5.6 (Math 531 Spring 2025, PS8.3)

A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is said to be T periodic if $f(t + T) = f(t)$, for all $t \in \mathbb{R}$. Now, given a function \tilde{f} defined on $[0, T)$, we can always extend \tilde{f} to be T periodic in the following way. First, we can write

$$\mathbb{R} = \bigcup_{n \in \mathbb{Z}} [nT, (n+1)T). \quad (649)$$

Then define for $t \in [nT, (n+1)T)$:

$$f(t) = \tilde{f}(t - nT). \quad (650)$$

- Show that f defined this way is T periodic.
- Suppose \tilde{f} is continuous on $[0, T)$. Does this mean that its extension f will also be continuous on \mathbb{R} ? What condition do you have to add?
- Suppose \tilde{f} is continuously differentiable on $[0, T)$. What conditions do you have to put on \tilde{f} to ensure that f is continuously differentiable? How about k -times continuously differentiable?

Solution.

Exercise 5.7 (Math 531 Spring 2025, PS8.4)

Assume $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable and $|f'(x)| \leq \frac{1}{1+x^2}$. Prove that

$$\lim_{x \rightarrow \infty} f(x) \quad (651)$$

exists. You are not allowed to use integration. Hint: How do you prove convergence without knowing what the limit is?

Solution.

Exercise 5.8 (Math 531 Spring 2025, PS8.5)

In a previous homework assignment, we defined the function

$$E(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!}. \quad (652)$$

We proved it is continuous, satisfies $E(z+w) = E(z)E(w)$ for all $z, w \in \mathbb{C}$, and then deduced that for all $x \in \mathbb{R}$, we have that $E(x) = e^x$.

- Prove that $E'(0) = 1$. Hint: write $\frac{H(x)-H(0)}{x} = \frac{1}{x} \sum_{n=1}^{\infty} \frac{x^n}{n!} = \sum_{n=1}^{\infty} \frac{x^{n-1}}{n!}$. Then prove that

$$\lim_{x \rightarrow 0} \sum_{n=1}^{\infty} \frac{x^{n-1}}{n!} = 1. \quad (653)$$

- By studying the difference quotient, prove that $E'(x) = E(x)$ for all $x \in \mathbb{R}$. This is much easier than the preceding point.
- Prove that $\lim_{x \rightarrow \infty} E(x) = \infty$ and $\lim_{x \rightarrow -\infty} E(x) = 0$.
- Prove that $E : (-\infty, \infty) \rightarrow (0, \infty)$ is 1-1 and onto.
- Let $L = E^{-1} : (0, \infty) \rightarrow (-\infty, \infty)$. Prove that

$$L'(t) = \frac{1}{t}, \quad (654)$$

for all $t \in (0, \infty)$.

Solution.

Exercise 5.9 (Math 531 Spring 2025, PS8.6)

For each $k \in \mathbb{N}$, consider $\sum_{j=1}^N j^k$. It turns out that this can be expressed as a polynomial of degree $k+1$ in N . For example,

$$\sum_{j=1}^N j^0 = N, \quad (655)$$

is a polynomial of degree 1 in N . Similarly,

$$\sum_{j=1}^N j = \frac{N(N+1)}{2} = \frac{N^2}{2} + \frac{N}{2}, \quad (656)$$

is a polynomial of degree 2 in N . If we write:

$$\sum_{j=1}^N j^k = a_0 + a_1 N + a_2 N^2 + \dots + a_{k+1} N^{k+1}, \quad (657)$$

what is the value of a_{k+1} ? Hint: divide by N^{k+1} and take the limit as $N \rightarrow \infty$.

Solution.

Exercise 5.10 (Math 531 Spring 2025, PS9.3)

Assume we have a twice differentiable function $f : [0, \infty) \rightarrow \mathbb{R}$. Assume that $f''(x) > 0$ for all $x \in [0, \infty)$, while $f'(0) \geq 0$. Prove that $\lim_{x \rightarrow \infty} f(x) = +\infty$.

Solution.

Exercise 5.11 (Math 531 Spring 2025, PS9.4)

We proved that the function $E : \mathbb{C} \rightarrow \mathbb{C}$ defined by:

$$E(z) = \sum_{k=0}^{\infty} \frac{z^k}{k!} \quad (658)$$

satisfies $E(z+w) = E(z)E(w)$. Let us now investigate the real and imaginary parts of $E(it)$, where $t \in \mathbb{R}$. Let us call the real part $C(t)$ and the imaginary part $S(t)$.

- Prove that $E(\bar{z}) = \overline{E(z)}$, for every $z \in \mathbb{C}$.
- Deduce that $|E(it)| = 1$ for all $t \in \mathbb{R}$ and thus:

$$C(t)^2 + S(t)^2 = 1, \quad (659)$$

for all $t \in \mathbb{R}$.

- Prove that $C(-t) = C(t)$ for all t and that $S(-t) = -S(t)$ for all t .
- Prove that $C(0) = 1$ and $S(0) = 0$, while $C'(t) = -S(t)$ and $S'(t) = C(t)$, for all $t \in \mathbb{R}$.
- Deduce that $C''(t) = -C(t)$ and prove that there must be some $t > 0$ for which $C(t) = 0$. (Hint: Use Problem 3)
- Prove that there is a smallest $t_* > 0$ for which $C(t_*) = 0$.
- Define $\pi = 2t_*$ so that $C(\frac{\pi}{2}) = 0$. Since S is increasing on $[0, t_*]$, deduce that $S(\frac{\pi}{2}) = 1$.
- Now use the formula $E(z+w) = E(z)E(w)$ to deduce that

$$C(t+2\pi) = C(t), \quad S(t+2\pi) = S(t), \quad (660)$$

for all $t \in \mathbb{R}$.

- It is now reasonable to unveil that C and S are none other but our old friends: \cos and \sin .

Solution.

6 Riemann Integration

We would like to define an integral. We do this by defining the Riemann sums for a particular partition, which is some number in \mathbb{R} . By taking lower and upper approximations and then taking the supremum and infimum of them, respectively, we can determine what it means for a function to be Riemann integrable.

Definition 6.1 (Partition)

Let $[a, b]$ be an interval. A **partition** P of $[a, b]$ is a set of $P = \{x_0, \dots, x_n\}$ (note that this is finite!) s.t.

$$a = x_0 \leq x_1 \leq \dots \leq x_{n-1} \leq x_n = b \quad (661)$$

with $\Delta x_i = [x_{i-1}, x_i]$ for $i = 1, \dots, n$.

Note that the natural way to define the Riemann integral is as the limit of the finite Riemann sums as partitions gets finer and finer. But we must be careful in saying what “finer” means. It is not simply as the number of partitions $n \rightarrow \infty$, since this may lead to multiple subsequential values of convergence by increasing the partition within different subsets of $[a, b]$.




Figure 23: Upper (top row) and lower (bottom row) Riemann sums with refinement of partition. In the upper row, the rightmost rectangle remains fixed while other rectangles become thinner. In the lower row, the leftmost rectangle remains fixed while other rectangles become thinner.

An alternative way is to have the partitions all converge “uniformly” as in the maximum length of an interval in a partition must go to 0.




Figure 24: Approximating an integral with increasingly fine partitions

A cleaner way it to simply look at the set of all partitions along with the set of the corresponding upper and lower Riemann sums, and then hope that they behave nicely with each other. This is the approach we will take.

Definition 6.2 (Riemann Sums with Respect to Partition)

Let P be a partition of $[a, b]$ and $f : [a, b] \rightarrow \mathbb{R}$ be bounded. Then, the **upper and lower Riemann sum** is defined

$$U(P, f) = \sum_{i=1}^n M_i \Delta x_i, \quad L(P, f) = \sum_{i=1}^n m_i \Delta x_i \quad (662)$$

where

$$M_i = \sup_{\Delta x_i} f(x), \quad m_i = \inf_{\Delta x_i} f(x) \quad (663)$$




Figure 25

Definition 6.3 (Riemann Integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. Then, the **upper and lower Riemann integrals** of $f(x)$ are defined

$$\int_a^b f := \inf_P U(P, f), \quad \overline{\int_a^b} f := \sup_P L(P, f) \quad (664)$$

If the upper and lower Riemann integrals are equal, then f is said to be **Riemann integrable** over $[a, b]$, denoted $f \in \mathcal{R}([a, b])$.^a

^aWhere $\mathcal{R}(X)$ is the set of all Riemann integrable functions over X .

In some textbooks, we also define a *partition with distinguished points* which simply is a partition P along with some set of ξ_i 's that land in each interval. This allows for extra degrees of freedom for choosing points, but it is generally not needed here.




Figure 26: Riemann sum approximation using sample points ξ_i within each subinterval. This is known as a Riemann sum of a partition with distinguished points. The Riemann sum is a mapping that takes in a partition with distinguished points $p = (P, \xi)$ on the closed interval $[a, b]$ and outputs a number representing the total area of the Riemann sums.

6.1 Axiomatic Properties of the Integral

The most important properties of integrable functions is linearity, additivity, and monotonicity.

Theorem 6.1 (Linearity of the Riemann Integral)

The set of Riemann integrable functions $\mathcal{R}[a, b]$ over closed interval $[a, b]$ is a vector space. That is, given $f, g \in \mathcal{R}[a, b]$ and $c \in \mathbb{R}$, then

1. $(f + g) \in \mathcal{R}[a, b]$

2. $(cf) \in \mathcal{R}[a, b]$

which makes $\mathcal{R}([a, b])$ into a \mathbb{R} -vector space.

Proof. We prove the following properties of a vector space.

1. If $c \in \mathbb{R}$ and $f \in \mathcal{R}$, then we wish to show that $cf \in \mathcal{R}$ and $\int cf = c \int f$.

- (a) If $c > 0$, then $U(cf, P) = cU(f, P)$, and $L(cf, P) = cL(f, P)$.

- (b) If $c < 0$, then $U(cf, P) = cL(f, P)$, and $L(cf, P) = cU(f, P)$.

So, for all $\epsilon > 0$, we can find P s.t.

$$U(f, P) - L(f, P) < \frac{\epsilon}{c} \implies U(cf, P) - L(cf, P) < \epsilon \quad (665)$$

and so $cf \in \mathcal{R}$

2. If $f_1, f_2 \in \mathcal{R}$, then

$$\text{osc}_E(f_1 + f_2) \leq \text{osc}_E(f_1) + \text{osc}_E(f_2) \text{ since } \begin{cases} \sup_E(f_1 + f_2) \leq \sup_E(f_1) + \sup_E(f_2) \\ \inf_E(f_1 + f_2) \geq \inf_E(f_1) + \inf_E(f_2) \end{cases} \quad (666)$$

for all $E \subset [a, b]$, which implies that $f_1 + f_2 \in \mathcal{R}$.

Lemma 6.1 (Restrictions of Integrable Functions)

The restriction of f in any $[c, d] \subset [a, b]$, denoted $f|_{[c, d]}$, is in $\mathcal{R}[c, d]$

Proof.

Theorem 6.2 (Additivity of Riemann Integral)

We have $\int_a^c + \int_c^b = \int_a^b$.

Proof. Let P be a partition of $[a, b]$. If $c \in P$, then we can view $P = P_1 \cup P_2$. If $c \notin P$, consider $P \cup \{c\}$. Then we have

$$U(f, P) = U(f, P_1) + U(f, P_2) \quad (667)$$

$$U(f, P) = L(f, P_1) + L(f, P_2) \quad (668)$$

So $f \in \mathcal{R}([a, c])$, $f \in \mathcal{R}([c, b])$.

Theorem 6.3 (Monotonicity of the Riemann Integral)

If $a \leq b$, $f_1, f_2 \in \mathcal{R}[a, b]$, and $f_1(x) \leq f_2(x)$ for every $x \in [a, b]$, then

$$\int_a^b f_1 \leq \int_a^b f_2 \quad (669)$$

This immediately implies that given constants m, M such that $m \leq f(x) \leq M$ at each $x \in [a, b]$, we have

$$m \cdot (b - a) \leq \int_a^b f(x) \leq M \cdot (b - a) \quad (670)$$

In particular, if $0 \leq f(x)$ on $[a, b]$, then

$$0 \leq \int_a^b f(x) \quad (671)$$

Theorem 6.4 ()

For fixed $a, b \in \mathbb{R}$ with $a < b$, $f \mapsto \int_a^b f$ is a linear map on $\mathcal{R}([a, b])$, i.e. a dual vector.

Proof. Removing the a, b for convenience, we first show that $\int f_1 + f_2 = \int f_1 + \int f_2$. Let $\epsilon > 0$. Then there exists P_i s.t.

$$U(f_i, P_i) < \int f_i + \epsilon \quad (672)$$

for $i = 1, 2$. Define $P = P_1 \cup P_2$ as the common refinement. Then

$$U(f_i, P) < \int f_i + \epsilon \quad (673)$$

and so

$$\int f_1 + f_2 \leq U(f_1 + f_2, P) \leq U(f_1, P) + U(f_2, P) \quad (674)$$

$$\leq 2\epsilon + \int f_1 + \int f_2 \quad (675)$$

which implies $\int f_1 + f_2 \leq \int f_1 + \int f_2$. To prove the other way, we see that

$$\int (-f_1) + (-f_2) \leq \int (-f_1) + \int (-f_2) \quad (676)$$

and so

$$-\int f_1 + f_2 \leq -\left(\int f_1 + \int f_2\right) \implies \int f_1 + f_2 \geq \int f_1 + \int f_2 \quad (677)$$

For scalar multiplication, we can do similarly.

6.2 Conditions for Riemann Integrability

Great, so we've defined Riemann integrable functions, but it's hard to determine whether a function is Riemann integrable—and if so—what the value of the integral is. We will determine the first problem by talking about sufficient conditions for Riemann integrability, and then introduce the fundamental theorem of calculus to address computability.

Definition 6.4 (Refinement)

P^* is a **refinement** of P if $P \subset P^*$. If P_1, P_2 are two partitions, then their **common refinement** $P^* = P_1 \cup P_2$.




Figure 27: Partitions P' and P'' with their common refinement \tilde{P}

Lemma 6.2 (Fundamental Lemma)

If P^* is a refinement of P and $f : [a, b] \rightarrow \mathbb{R}$ is bounded, then

$$L(P, f) \leq L(P^*, f) \leq U(P^*, f) \leq U(P, f) \quad (678)$$

Proof. By induction on the number of points we add to P to get P^* , we might as well assume that $P^* = P \cup \{x_*\}$. So,

$$P = \{a = x_0, x_1, \dots, x_{n-1}, x_n\} \quad (679)$$

$$P^* = \{a = x_0, x_1, \dots, x_{i-1}, x_i, x_*, x_{i+1}, \dots, x_{n-1}, x_n\} \quad (680)$$

$$(681)$$

Now let's compute $L(f, P^*) - L(f, P)$. Since the only intervals affected are $[x_i, x_{i+1}]$, we have

$$L(f, P^*) - L(f, P) = \inf_{[x_i, x_*]} f(x)(x_* - x_i) + \inf_{[x_*, x_{i+1}]} f(x)(x_{i+1} - x_*) - \inf_{[x_i, x_{i+1}]} f(x)(x_{i+1} - x_i) \quad (682)$$

$$= \underbrace{\left(\inf_{[x_i, x_*]} f(x) - \inf_{[x_*, x_{i+1}]} f(x) \right)}_{>0} (x_* - x_i) + \underbrace{\left(\inf_{[x_*, x_{i+1}]} f(x) - \inf_{[x_i, x_{i+1}]} f(x) \right)}_{>0} (x_{i+1} - x_*) \quad (683)$$

which is therefore greater than 0.

Theorem 6.5 (Lower and Upper Integrals as Bounds of Each Other)

We claim

$$\int_a^b f \leq \overline{\int_a^b f} \quad (684)$$

Proof. Given P_1, P_2 partitions, let $P^* = P_1 \cup P_2$ be their common refinement. Then, from the theorem above,

$$L(P_2, f) \leq L(P^*, f) \leq U(P^*, f) \leq U(P_1, f) \quad (685)$$

So taking the supremum over all partitions P_2 and fixing P_1 gives

$$\int_a^b f = \sup_{P_2} L(P_2, f) \leq \sup_{P_2} U(P_1, f) = U(P_1, f) \quad (686)$$

Then taking the infimum over all partitions P_1 gives us

$$\int_a^b f(x) = \inf_{P_1} \int_a^b f(x) \leq \inf_{P_1} U(P_1, f) = \overline{\int_a^b f(x)} \quad (687)$$

where we note that the infimum does not affect the terms that do not depend on P_1 .

We have seen some bounds of the upper and lower integrals, and defined the Riemann integral. However, checking Riemann integrability is quite tedious, since we have to take the supremum and infimum over all possible partitions. The following theorem is extremely useful as it only requires us to find *one* partition given some ϵ . This is because that the Riemann integral, as complicated as the formula is, is still a limit of a function. That means that we can apply the Cauchy criterion to it to determine convergence.

Theorem 6.6 (Cauchy Criterion for Riemann Integrability)

$f \in \mathcal{R}$ iff $\forall \epsilon > 0$, there exists partition P such that $U(P, f) - L(P, f) < \epsilon$.

Proof. We prove bidirectionally. The reverse implication is easy, but for the forward direction you must use refinements.

1. (\leftarrow). Pick any partition P . Since

$$L(f, P) \leq \int_a^b f \leq \overline{\int_a^b f} \leq U(f, P) \quad (688)$$

This implies that

$$0 \leq \overline{\int_a^b f} - \underline{\int_a^b f} \leq U(f, P) - L(f, P) < \epsilon \quad (689)$$

and since any nonnegative number less than any positive number must be 0 (since there are no infinitesimals in \mathbb{R}), the LHS is 0, and the result is proven.

2. (\rightarrow). f is Riemann integrable, so

$$\overline{\int_a^b f} = \underline{\int_a^b f} \iff \inf_P U(f, P) = \sup_Q L(f, Q) \quad (690)$$

for partitions P, Q . So we can find P that gets really close to the infimum and same for Q close to the supremum, i.e. there exists a P, Q , such that

$$U(f, P) < \overline{\int_a^b f} + \frac{\epsilon}{2}, \quad L(f, Q) > \underline{\int_a^b f} - \frac{\epsilon}{2} \quad (691)$$

Now take the common refinement $P^* = P \cup Q$, and so by the fundamental lemma,

$$\underline{\int_a^b f} - \frac{\epsilon}{2} < L(f, Q) \leq L(f, P^*) \leq U(f, P^*) \leq U(f, P) < \overline{\int_a^b f} + \frac{\epsilon}{2} \quad (692)$$

which implies that $0 \leq U(f, P^*) - L(f, P^*) < \epsilon$.

Note that a necessary condition of f being Riemann integrable is that f is bounded. In fact it is defined that way. You may know that a sufficient condition of integrability is that it is continuous, but we can prove something slightly weaker.

Definition 6.5 (Oscillation)

Given an interval I , the **oscillation** of f on I is defined

$$\text{osc}_I(f) := \sup_I(f) - \inf_I(f) \quad (693)$$

Intuitively, a function f is Riemann integrable if we can make $U(f, P) - L(f, P)$ as small as we wish. This is the case if we can find a sufficiently refined partition P such that the oscillation on f on each interval is small.

Lemma 6.3 (Functions with Vanishing Oscillations are Riemann Integrable)

Let f be bounded on a closed interval $[a, b]$. If, for every $\epsilon > 0$, there exists a partition P such that

$$\sum_{i=0}^{n-1} \text{osc}_{[x_i, x_{i+1}]} f < \epsilon \quad (694)$$

then f is Riemann integrable.

Proof. Given $\epsilon > 0$, choose $\epsilon/(b-a)$. By assumption we can find a partition P in which the total

oscillation is bounded above by $\epsilon/(b-a)$. Therefore,

$$U(P, f) - L(P, f) = \sum_{i=0}^{n-1} \sup_{[x_i, x_{i+1}]} f(x) \Delta x_i - \sum_{i=0}^{n-1} \inf_{[x_i, x_{i+1}]} f(x) \Delta x_i \quad (695)$$

$$= \sum_{i=0}^{n-1} \left(\sup_{[x_i, x_{i+1}]} f(x) - \inf_{[x_i, x_{i+1}]} f(x) \right) \Delta x_i \quad (696)$$

$$< \sum_{i=0}^{n-1} \operatorname{osc}_{[x_i, x_{i+1}]} f \Delta x_i \quad (697)$$

$$\leq \sum_{i=0}^{n-1} \frac{\epsilon}{b-a} \Delta x_i \quad (698)$$

$$= \frac{\epsilon}{b-a} \sum_{i=0}^{n-1} \Delta x_i \quad (699)$$

$$= \frac{\epsilon}{b-a} (b-a) = \epsilon \quad (700)$$

Here is a classic example of a non-integrable function.

Example 6.1 (Non-Integrability of the Dirichlet Function)

The Dirichlet function

$$\mathcal{D}(x) := \begin{cases} 1, & \text{for } x \in \mathbb{Q} \\ 0, & \text{for } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases} \quad (701)$$

on the interval $[0, 1]$ is not integrable on that interval. For any partition P of $[0, 1]$ we can find in each interval Δ_i both a rational point ξ'_i and an irrational point ξ''_i . Then, we can see that the lower and upper Riemann sums do not necessarily converge to each other since

$$\sigma(f; P, \xi') = \sum_{i=1}^n 1 \cdot \Delta x_i = 1 \text{ while } \sigma(f; P, \xi'') = \sum_{i=1}^n 0 \cdot \Delta x_i = 0 \quad (702)$$

as $\lambda(P) \rightarrow 0$.

Example 6.2 ()

Is there a function f that is discontinuous on a dense set of $[0, 1]$ but still Riemann integrable?

With this, we can use the uniform continuity of continuous functions over a compact set to place a bound on the oscillation of each subinterval—and thus a bound on the oscillation of the whole interval.

Theorem 6.7 (Continuous Functions are Riemann Integrable)

f continuous on $[a, b] \implies f$ is Riemann integrable on $[a, b]$.

Proof. If f is continuous, then by EVT it is bounded and uniformly continuous. Therefore we can take the evenly-partitioned intervals of $[a, b]$ and by uniform continuity, the oscillation tends to 0, and we are done.

Perhaps more explicitly, we wish to show that for all $\epsilon > 0$, there exists partition P s.t. $U(P, f) -$

$L(P, f) < \epsilon$. Now let $\epsilon > 0$, and since it's uniformly continuous, take $\delta > 0$ s.t.

$$|x - y| < \delta \implies |f(x) - f(y)| < \frac{\epsilon}{2(b-a)} \quad (703)$$

Let $N \in \mathbb{N}$ be so large that $\frac{b-a}{N} < \delta$. Now consider the partition of $[a, b]$ given by $x_i = a + \frac{b-a}{N}i$ for $0 \leq i < N$. Intuitively, we want these subintervals to be so small that f will not deviate too widely. So it better be the case that $\frac{b-a}{N} < \delta$. So, we have

$$U(P, f) - L(P, f) = \sum_{i=1}^n \sup_{[x_i, x_{i+1}]} f(x) \Delta x_i - \sum_{i=1}^n \inf_{[x_i, x_{i+1}]} f(x) \Delta x_i \quad (704)$$

$$= \sum_{i=1}^n \left(\sup_{[x_i, x_{i+1}]} f(x) - \inf_{[x_i, x_{i+1}]} f(x) \right) \Delta x_i \quad (705)$$

$$< \sum_{i=0}^{N-1} \frac{\epsilon}{2(b-a)} \Delta x_i \quad (706)$$

$$= \frac{\epsilon}{2(b-a)} \cdot (b-a) < \frac{\epsilon}{2} < \epsilon \quad (707)$$

We can actually make a stronger claim.

Corollary 6.1 (Integrability of Discontinuous Functions)

If a bounded function f on a closed interval $[a, b]$ is continuous everywhere except at a finite set of points, then $f \in \mathcal{R}[a, b]$.

Corollary 6.2 (Integrability of Monotonic Functions)

A bounded monotonic function on a closed interval is integrable on that interval.

Theorem 6.8 (Continuous Compositions of Integrable Functions are Integrable)

Let $f \in \mathcal{R}([a, b])$. Assume $\phi : \mathbb{R} \rightarrow \mathbb{R}$ is continuous. Then $\phi \circ f \in \mathcal{R}([a, b])$.

Proof. Since $f \in \mathcal{R}([a, b])$ is bounded, let $|f(x)| \leq M$ for all $x \in [a, b]$ for some $M \geq 0$. Now let $K = \sup_{t \in [-M, M]} \phi(t)$, which exists since $[-M, M]$ is compact and ϕ is continuous. ϕ is also uniformly continuous on $[-M, M]$.

Now let $\epsilon > 0$. Then there exists a $\delta > 0$ s.t. $|t - s| < \delta \implies |\phi(t) - \phi(s)| < \epsilon$. Consequently,

$$|f(x) - f(y)| < \delta \implies |\phi(f(x)) - \phi(f(y))| < \epsilon \quad (708)$$

Since $f \in \mathcal{R}([a, b])$, we can find a partition P of $[a, b]$ s.t.

$$U(f, P) - L(f, P) < \delta^2 \implies \sum_{i=1}^{n-1} \left(\sup_{[x_i, x_{i+1}]} f - \inf_{[x_i, x_{i+1}]} f \right) \Delta x_i < \delta^2 \quad (709)$$

Let

$$A = \{i \mid \sup_{[x_i, x_{i+1}]} f - \inf_{[x_i, x_{i+1}]} f < \delta\} \quad (710)$$

$$B = \{i \mid \sup_{[x_i, x_{i+1}]} f - \inf_{[x_i, x_{i+1}]} f \geq \delta\} \quad (711)$$

Colloquially, we can think of A as the “good” intervals with small oscillations, and B as the “bad” intervals with larger oscillations. So,

$$\sum_{i \in B} \Delta x_i = \frac{1}{\delta} \sum_{i \in B} \delta \Delta x_i \leq \frac{1}{\delta} \sum_{i \in B} \text{osc}_{[x_i, x_{i+1}]} \Delta x_i < \frac{1}{\delta} \delta^2 = \delta \quad (712)$$

Now, compute

$$U(\phi(f), P) - L(\phi(f), P) = \sum_i \text{osc}_{[x_i, x_{i+1}]}(\phi(f)) \Delta x_i \quad (713)$$

$$= \sum_{i \in A} \text{osc}_{[x_i, x_{i+1}]}(\phi(f)) \Delta x_i + \sum_{i \in B} \text{osc}_{[x_i, x_{i+1}]}(\phi(f)) \Delta x_i \quad (714)$$

In the good sets, if $f(x)$ ’s are within δ of each other, the oscillation by uniform continuity implies $\text{osc}(\phi(f)) < \epsilon$. In the bad set, we have $\text{osc}_{[x_i, x_{i+1}]}(\phi(f)) < 2K$, so the above can be bounded by

$$'' \leq \epsilon \sum_{i \in A} \Delta x_i + \sum_{i \in B} 2K \Delta x_i \quad (715)$$

$$\leq \epsilon(b - a) + 2K\delta \quad (716)$$

$$< \epsilon(b - a + 2K) \quad (717)$$

where the penultimate step is due to $\sum_{i \in B} \Delta x_i < \delta$.

However, contrary to intuition, f, g both integrable does not imply that $g \circ f$ is integrable. We present a counterexample.

Example 6.3 (Composition of Integrable Functions May Not be Integrable)

Consider the functions

$$|\text{sgn}|(x) := \begin{cases} 1 & x \neq 0 \\ 0 & x = 0 \end{cases} \quad (718)$$

and the Riemann function

$$\mathcal{R}(x) := \begin{cases} \frac{1}{n} & x = \frac{m}{n} \in \mathbb{Q}, \gcd(m, n) = 1 \\ 0 & x \in \mathbb{R} \setminus \mathbb{Q} \end{cases} \quad (719)$$

We can see that \mathcal{R} is continuous at all irrational points and discontinuous at all rational points except 0, meaning that it is integrable (\mathbb{Q} has measure zero). Then, the composition of these two functions is precisely the Dirichlet function

$$\mathcal{D}(x) = |\text{sgn}| \circ \mathcal{R} \quad (720)$$

which is not integrable.

6.3 Further Properties of the Riemann Integral

Theorem 6.9 (Compositions?)

Theorem 6.10 (Products of Riemann Integrable Functions are Riemann Integrable)

If $f, g \in \mathcal{R}[a, b]$, then $fg \in \mathcal{R}[a, b]$.

Proof. A nice trick is that

$$fg = \frac{1}{4}((f+g)^2 - (f-g)^2) \quad (721)$$

which is in $\mathcal{R}([a, b])$ since the sum, difference, and squaring functions are all continuous, and hence the composition $\phi(f, g)$ is Riemann integrable.

Theorem 6.11 (Triangle Inequality for Riemann Integrals)

If $f \in \mathcal{R}[a, b]$, then $|f| \in \mathcal{R}[a, b]$. Furthermore,

$$\left| \int_a^b f \right| \leq \int_a^b |f| \quad (722)$$

Proof. $\phi(x) = |x|$ is continuous, so $\phi(f) \in \mathcal{R}$. Note that if $f \geq 0$, then $\int_a^b f \geq 0$. Consider $|f| - f$ and $|f| + f$, both ≥ 0 . They are integrable as the image of f composed with continuous functions. So we have

$$\int |f| + f \geq 0 \implies \int |f| \geq - \int f \quad (723)$$

$$\int |f| - f \geq 0 \implies \int |f| \geq \int f \quad (724)$$

and so taking the maximum of the right hand side gives $\int |f| \geq |\int f|$.

This will later allow us to define inner products on function spaces.

Example 6.4 ()

Consider the space $X = C([a, b])$. Define $d : X \times X \rightarrow \mathbb{R}_0^+$ as

$$d(f, g) := \int_a^b |f - g| \quad (725)$$

Then d is a metric. Note that in $\mathcal{R}([a, b])$, it is *not* a metric since $d(f, g) = 0 \not\Rightarrow f = g$. Consider two functions that are different in 1 point.

So, we can define a vector space.

Theorem 6.12 (Mean Value Theorem of the Integral)

Given $f \in \mathcal{R}[a, b]$, with

$$m = \inf_{x \in [a, b]} f(x), \quad M = \sup_{x \in [a, b]} f(x) \quad (726)$$

Then

1. there exists a number $\mu \in [m, M]$ such that

$$\int_a^b f = \mu \cdot (b - a) \quad (727)$$

2. Furthermore, if $f \in C[a, b]$, it there exists a point $\xi \in [a, b]$ such that

$$\int_a^b f = f(\xi)(b - a) \quad (728)$$

Theorem 6.13 (Bonnet's Formula)

If $f, g \in \mathcal{R}[a, b]$ and g is a monotonic function on $[a, b]$, then there exists a point $\xi \in [a, b]$ such that

$$\int_a^b (f \cdot g)(x) dx = g(a) \int_a^\xi f(x) dx + g(b) \int_\xi^b f(x) dx \quad (729)$$

6.4 The Fundamental Theorem of Calculus

Let $f \in \mathcal{R}[a, b]$, and let us choose an $x \in [a, b]$ in order to construct the function

$$F(x) := \int_a^x f(t) dt \quad (730)$$

which is called an integral with a variable upper limit. By doing this, we can “upgrade” a Riemann integrable function f to a continuous function F .

Theorem 6.14 (First Fundamental Theorem of Calculus)

Define $F : [a, b] \rightarrow \mathbb{R}$ by

$$F(x) := \int_a^x f(t) dt \quad (731)$$

Then

1. F is continuous.
2. If F is continuous at x_0 , then $F'(x_0) = f(x_0)$.




Figure 28: This theorem amazingly tells us that the rate at which the integral F is increasing at x (represented by the increasing area under the curve of f) is equal to the value of f at the point x itself!

Proof. Listed.

1. Since $f \in \mathcal{R}([a, b])$, let $M = \sup_{x \in [a, b]} |f(x)| < +\infty$. WLOG let $x, y \in [a, b]$ with $x < y$. Then, we can use the “trick” by writing the difference of F as an integral, which follows from linearity of the integral over an interval. So, we have

$$|F(x) - F(y)| = \left| \int_x^y f(t) dt \right| \leq \int_x^y |f(t)| dt \quad (732)$$

$$\leq \int_x^y M dt = M|y - x| \quad (733)$$

So given $\epsilon > 0$, we can take $\delta = \epsilon/M$ and F is continuous.

2. Now let's claim

$$\lim_{h \rightarrow 0} \frac{1}{h} (F(x_0 + h) - F(x_0) - f(x_0)h) = 0 \iff F'(x_0) = f(x_0) \quad (734)$$

since if the limit exists, we can add $f(x_0)$ to both sides. The term in the limit is

$$\frac{1}{h} \left| \int_a^{x_0+h} f(t) dt - \int_a^{x_0} f(t) dt - f(x_0)h \right| \leq \frac{1}{h} \left| \int_{x_0}^{x_0+h} f(t) dt - hf(x_0) \right| \quad (735)$$

Now we do a trick that is simple but powerful. Notice that $hf(x_0) = \int_{x_0}^{x_0+h} f(x_0) dt$, so we can join it with the integral.^a So,

$$'' = \frac{1}{h} \left| \int_{x_0}^{x_0+h} f(t) - f(x_0) dt \right| \quad (736)$$

$$\leq \frac{1}{h} \int_{x_0}^{x_0+h} |f(t) - f(x_0)| dt \quad (737)$$

$$\leq \frac{1}{h} \int_{x_0}^{x_0+h} \sup_{t \in [x_0, x_0+h]} |f(t) - f(x_0)| dt \quad (738)$$

Note that the supremum term in the integral is just a number, so evaluating it and taking the limit as $h \rightarrow 0$ gives

$$\sup_{t \in [x_0, x_0+h]} |f(t) - f(x_0)| \rightarrow 0 \text{ as } h \rightarrow 0 \quad (739)$$

since f is continuous at x_0 .

^aElgindi talked about how simple tricks can go a long way, e.g. the guy who was a master of Cauchy-Schwarz inequality.

Corollary 6.3 ()

Every bounded function $f : [a, b] \rightarrow \mathbb{R}$ on the closed interval $[a, b]$ and has only a finite number of points of discontinuity has a primitive, and every primitive of f on $[a, b]$ has the form

$$F(x) := \int_a^x f(t) dt + c \quad (740)$$

where c is a constant.

Theorem 6.15 (Second Fundamental Theorem of Calculus)

Let f be a real-valued function on a closed interval $[a, b]$ with F any primitive of f on $[a, b]$. If f is Riemann-integrable (i.e. f bounded with finite points of Lebesgue measure zero) on $[a, b]$, then

$$\int_a^b f = F(b) - F(a) \quad (741)$$




Figure 29: Graphical illustration of the Fundamental Theorem of Calculus, showing how the definite integral equals the difference of antiderivative values.

Proof. We already know that a bounded function on a closed interval having a finite number of discontinuities is integrable, and by the corollary, we are guaranteed an existence of a primitive $F(x)$ of the function f on $[a, b]$ with the form

$$F(x) := \int_a^x f(t) dt + c \quad (742)$$

Setting $x = a$, we find that $c = F(a)$, and so

$$F(x) := \int_a^x f(t) dt + F(a) \quad (743)$$

Evaluating F at $x = b$ gives

$$\int_a^b f(t) dt = F(b) - F(a) \quad (744)$$

Theorem 6.16 (Integral Form of the Remainder)

If $f : E \rightarrow \mathbb{R}$ has continuous derivatives up to order n on the closed interval $[a, x]$, then Taylor's formula holds

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \dots + \frac{f^{(n-1)}(a)}{(n-1)!}(x-a)^{n-1} + r_{n-1}(a; x) \quad (745)$$

where

$$r_{n-1}(a; x) = \frac{1}{(n-1)!} \int_a^x f^{(n)}(t)(x-t)^{n-1} dt \quad (746)$$

This form is called **Taylor's formula with the integral form of the remainder**.

Proof. Using the 2nd fundamental theorem and the definite integration by parts formula, we can carry out the following chain of transformations, assuming continuity and differentiability when needed.

$$\begin{aligned}
 f(x) - f(a) &= \int_a^x f'(t) dt \\
 &= - \int_a^x f'(t)(x-t)' dt \\
 &= -f'(t)(x-t) \Big|_a^x + \int_a^x f''(t)(x-t) dt \\
 &= f'(a)(x-a) - \frac{1}{2} \int_a^x f''(t)((x-t)^2)' dt \\
 &= f'(a)(x-a) - \frac{1}{2} f''(t)(x-t)^2 \Big|_a^x + \frac{1}{2} \int_a^x f'''(t)(x-t)^2 dt \\
 &= f'(a)(x-a) + \frac{1}{2} f''(a)(x-a)^2 - \frac{1}{2 \cdot 3} \int_a^x f'''(t)((x-t)^3)' dt \\
 &= \dots \\
 &= f'(a)(x-a) + \dots + \frac{1}{(n-1)!} f^{(n-1)}(a)(x-a)^{n-1} + r_{n-1}(a; x)
 \end{aligned}$$

where $r_{n-1}(a; x)$ is given by the integral formula mentioned.

6.5 Change of Variable

We now show and prove the method what we call “u-substitution” for definite integration. We first start off with a change of variable induced by a monotonic continuous function, and then generalize it for C^1 functions.

Theorem 6.17 (Change of Variable for Monotonic Functions)

If

1. $f \in \mathcal{R}([a, b])$ and
2. φ is continuous and strictly increasing on $[a, b]$,

then

$$\int_a^b f(\varphi(x)) \cdot \varphi'(x) dx = \int_{\varphi(a)}^{\varphi(b)} f(u) du \quad (747)$$

Proof.

Theorem 6.18 (Change of Variable)

If

1. $f \in C([a, b])^a$, and
2. φ is continuously differentiable on $[a, b]$,

then

$$\int_a^b f(\varphi(x)) \cdot \varphi'(x) dx = \int_{\varphi(a)}^{\varphi(b)} f(u) du \quad (748)$$

^aNote that we strengthen our assumption to continuous, not just Riemann integrable f !

Proof.

Note that we have made the additional assumption that continuity is needed.

Example 6.5 (Counterexample when f is Not Continuous)

If f is Riemann integrable, ...

6.6 Integration by Parts

Now a direct application of the fundamental theorem of calculus is the integration by parts. By the product rule of differentiation, we have

$$(u \cdot v)'(x) = (u' \cdot v)(x) + (u \cdot v')(x) \quad (749)$$

where by hypothesis, $u' \cdot v, u \cdot v'$ are continuous and hence integrable on $[a, b]$. Using the linearity of the integral and the 2nd fundamental theorem of calculus, we get

$$(u \cdot v)(x)|_a^b = \int_a^b (u' \cdot v) + \int_a^b (u \cdot v') \quad (750)$$

Theorem 6.19 (Integration by Parts)

Suppose $F, G : [a, b] \rightarrow \mathbb{R}$ are differentiable, with $F' = f, G' = g \in \mathcal{R}([a, b])$. Then

$$\int_a^b Fg = (FG)|_a^b - \int_a^b fG \quad (751)$$

Proof.

6.7 Improper Integrals

Note that by definition of the Riemann integral, it makes sense to talk about integrability over *bounded* functions over *closed and bounded intervals*. This is in a sense quite restrictive, since we cannot integrate over “singularities” where either the interval or the function is unbounded. We develop the tools of improper integration to deal with this problem; there are two types of improper integrals.

Definition 6.6 (Improper Integral of Unbounded Interval)

Let f be bounded on its domain.

1. If it is defined on $[a, +\infty)$ and is integrable on every closed interval $[a, b] \subset [a, +\infty)$, then we define

$$\int_a^{+\infty} f := \lim_{b \rightarrow +\infty} \int_a^b f \quad (752)$$

2. If it is defined on $(-\infty, b]$ and is integrable on every closed interval $[a, b] \subset (-\infty, b]$, then we define

$$\int_{-\infty}^b f := \lim_{a \rightarrow -\infty} \int_a^b f \quad (753)$$

3. If it is defined on \mathbb{R} and is integrable on every closed interval $[a, b] \subset \mathbb{R}$, then we define^a

$$\int_{-\infty}^{+\infty} f := \lim_{b \rightarrow +\infty} \lim_{a \rightarrow -\infty} \int_a^b f \quad (754)$$

If the limit exists, then we say that the integral **converges** and **diverges** otherwise.

^aSmall technicality is that we have to prove that this is independent of the order in which we take the limits.

Example 6.6 (Gaussian Integral)

The integral

$$\int_{-\infty}^{+\infty} e^{-x^2} dx = \sqrt{\pi} \quad (755)$$

Definition 6.7 (Improper Integral of Unbounded Function)

Let $f : [a, b] \rightarrow \mathbb{R}$ be unbounded at some point ω in its domain and is integrable on any closed interval $[c, d] \subset [a, b]$ not containing ω .

1. If $\omega = a$, then we define

$$\int_a^b f := \lim_{c \rightarrow a^+} \int_c^b f \quad (756)$$

2. If $\omega = b$, then we define

$$\int_a^b f := \lim_{d \rightarrow b^-} \int_a^d f \quad (757)$$

3. If $\omega \in (a, b)$, then we define

$$\int_a^b f = \lim_{d \rightarrow s^+} \int_a^d f + \lim_{c \rightarrow s^-} \int_c^b f \quad (758)$$

Lemma 6.4 (Properties of the Improper Integral)

Suppose f, g are functions defined on interval $[a, \omega)$ (without loss of generality, we let ω be the upper limit of integration) and integrable on every closed interval $[a, b] \subset [a, \omega)$. Suppose the improper integrals

$$\int_a^\omega f \text{ and } \int_a^\omega g \quad (759)$$

are well-defined.

1. For any $\lambda_1, \lambda_2 \in \mathbb{R}$ the function $(\lambda_1 f + \lambda_2 g)(x)$ is integrable in the improper sense on $[a, \omega)$ and

$$\int_a^\omega \lambda_1 f + \lambda_2 g = \lambda_1 \int_a^\omega f + \lambda_2 \int_a^\omega g \quad (760)$$

2. For any $c \in [a, \omega)$,

$$\int_a^\omega f = \int_a^c f + \int_c^\omega f \quad (761)$$

3. If $\varphi : [\alpha, \gamma) \rightarrow [a, \omega)$ is a smooth strictly monotonic mapping with $\varphi(\alpha) = a$ and $\varphi(\beta) \rightarrow \omega$ as $\beta \rightarrow \gamma^-$, then the improper integral of the function $t \mapsto (f \circ \varphi)(t)\varphi'(t)$ over $[\alpha, \gamma)$ exists and

$$\int_a^\omega f(x) dx = \int_\alpha^\gamma (f \circ \varphi)(t)\varphi'(t) dt \quad (762)$$

Note that by definition, an improper integral $\int_a^\omega f := \lim_{b \rightarrow \omega} \int_a^b f$ is a limit of the function $F(b) := \int_a^b f$ as $b \rightarrow \omega$. This means that we can use the Cauchy criterion to determine the convergence of this limit, and hence, existence of this improper integral.

Theorem 6.20 (Cauchy Criterion for Convergence of an Improper Integral)

If the function $x \mapsto f(x)$ is defined on the interval $[a, \omega)$ and integrable on every closed interval $[a, b] \subset [a, \omega)$, then the integral

$$\int_a^\omega f \quad (763)$$

converges if and only if for every $\epsilon > 0$ there exists $B \in [a, \omega)$ such that the relation

$$\left| \int_{b_1}^{b_2} f \right| < \epsilon \quad (764)$$

holds for any $b_1, b_2 \in [a, \omega)$ satisfying $B < b_1$ and $B < b_2$.

Proof. We have

$$\int_{b_1}^{b_2} f = \int_a^{b_2} f - \int_a^{b_1} f = F(b_2) - F(b_1) \quad (765)$$

and therefore the condition is simply the Cauchy criterion for the existence of a limit for the function $\mathcal{F}(b)$ as $b \rightarrow \omega$.

Definition 6.8 (Absolute Convergence of an Improper Integral)

The improper integral

$$\int_a^\omega f \quad (766)$$

converges absolutely if the integral

$$\int_a^\omega |f| \quad (767)$$

converges. Clearly, the inequality

$$\left| \int_{b_1}^{b_2} f \right| \leq \left| \int_{b_1}^{b_2} |f| \right| \quad (768)$$

implies that if an improper integral converges absolutely, then it converges.

This study of absolute convergence reduces to the study of convergence of integrals of nonnegative functions. The following lemma is useful in determining convergence of such functions.

Lemma 6.5 ()

Let there be a function f defined on interval $[a, \omega)$ that is also integrable over every closed interval $[a, b] \subset [a, \omega)$. If $f(x) \geq 0$ on $[a, \omega)$, then the improper integral

$$\int_a^\omega f \quad (769)$$

exists if and only if the function

$$\mathcal{F}(b) := \int_a^b f \quad (770)$$

is bounded on $[a, \omega)$.

Proof. It is clear that

$$\int_a^\omega f = \lim_{b \rightarrow \omega} \mathcal{F}(b) \quad (771)$$

If $f(x) \geq 0$, then the function $\mathcal{F}(b)$ is nondecreasing on $[a, \omega)$ and therefore has a limit as $b \rightarrow \omega$ only if it is bounded (since every monotonically increasing sequence that is bounded always converges).

This leads to the familiar integral test for convergence of a series.

Theorem 6.21 (Integral Test for Convergence of a Series)

If the function $x \mapsto f(x)$ is defined on the interval $[1, +\infty)$, nonnegative, nonincreasing, and integrable on each closed interval $[1, b] \subset [1, +\infty)$, then the series

$$\sum_{n=1}^{\infty} f(n) = f(1) + f(2) + \dots \quad (772)$$

and the integral

$$\int_a^{+\infty} f \quad (773)$$

either both converge or both diverge.

We can use the comparison test analogue to determine convergence of improper integrals.

Theorem 6.22 (Comparison Test for Convergence of Improper Integrals)

Suppose the functions $f(x), g(x)$ are defined on the interval $[a, \omega)$ and integrable on any closed interval $[a, b] \subset [a, \omega)$. If

$$0 \leq f \leq g \quad (774)$$

on $[a, \omega)$, then

$$\int_a^\omega g \text{ converges} \implies \int_a^\omega f \text{ converges} \quad (775)$$

and the inequality

$$\int_a^\omega f \leq \int_a^\omega g \quad (776)$$

holds. Also,

$$\int_a^\omega f \text{ diverges} \implies \int_a^\omega g \text{ diverges} \quad (777)$$

6.8 Integration over Paths and Rectifiable Curves

Definition 6.9 (Integration For Vector Valued Functions)

A function $f : [a, b] \rightarrow \mathbb{R}^d$ is Riemann integrable if $f = (f_1, \dots, f_d)$ and each component $f_i : [a, b] \rightarrow \mathbb{R}$ is in $\mathcal{R}([a, b])$. The integral is defined

$$\int_a^b f = \left(\int_a^b f_1, \dots, \int_a^b f_d \right) \quad (778)$$

Now since the codomain is \mathbb{R}^d , we can use the Euclidean norm $|v| := (\sum_i v_i^2)^{1/2}$ on it.

Theorem 6.23 ()

If $f \in \mathcal{R}([a, b], \mathbb{R}^d)$, then $|f| \in \mathcal{R}([a, b], \mathbb{R}^d)$ and

$$\left| \int_a^b f \right| \leq \int_a^b |f| \quad (779)$$

Proof. If $f \in \mathcal{R}([a, b], \mathbb{R}^d)$, then $f_i \in \mathcal{R}([a, b])$, and so

$$|f| = \sqrt{f_1^2 + \dots + f_d^2} \in \mathcal{R} \quad (780)$$

since $x \mapsto x^2$ and $x \mapsto \sqrt{x}$ are continuous. Now consider the vector $v = \int_a^b f$. Then

$$|v| = \left| \int_a^b f \right| \implies |v|^2 = \sum_{j=1}^d v_j^2 = \sum_{j=1}^d v_j \int_a^b f_j \quad (781)$$

$$= \int_a^b \sum_{j=1}^d v_j f_j \quad (782)$$

$$= \int_a^b \sum_{j=1}^d v_j f_j \quad (783)$$

$$= \int_a^b \langle v, f(t) \rangle dt \quad (784)$$

$$\leq \int_a^b |v| |f(t)| dt \quad (785)$$

and so

$$|v|^2 \leq |v| \cdot \int_a^b |f(t)| dt \implies |v| \leq \int_a^b |f(t)| dt \quad (786)$$

Definition 6.10 (Curve)

A **curve** is a function $\gamma : [0, 1] \rightarrow \mathbb{R}^d$.

1. If $\gamma(0) = \gamma(1)$, then it is a **closed curve**.
2. If γ is injective, then it is called a **simple curve**.

Curves are usually continuous but does not have to be.

Example 6.7 ()

The curve can have different parameterizations and/or image. For example, the two are different curves with the image in $S^1 \subset \mathbb{R}^2$.

$$\gamma(t) = (\cos(2\pi t), \sin(2\pi t)) \quad (787)$$

$$\tilde{\gamma}(t) = (\cos(4\pi t), \sin(4\pi t)) \quad (788)$$

Definition 6.11 (Length of a Curve)

Given a curve $\gamma : [0, 1] \rightarrow \mathbb{R}^d$ and partition P of $[0, 1]$, let

$$\Lambda(\gamma, P) = \sum_{i=1}^N |\gamma(x_i) - \gamma(x_{i-1})| \quad (789)$$

i.e. the sum of the straight line distances between the curves. The **length** of the curve is defined as

$$\Lambda(\gamma) := \sup_P \Lambda(\gamma, P) \quad (790)$$

If the length is finite, then we call this a **rectifiable curve**.

Example 6.8 ()

Consider the curve given by

$$\gamma(t) = \left(t, t \sin \frac{1}{t} \right) \quad (791)$$

γ is continuous but $\gamma(t) < +\infty$.

For most continuous curves, this is not finite, but there is a sufficient condition for it to be finite.

Theorem 6.24 (C^1 Curves are Rectifiable)

If $\gamma : [0, 1] \rightarrow \mathbb{R}^d$ is continuously differentiable, then γ is rectifiable, and

$$\Lambda(\gamma) = \int_0^1 |\gamma'(t)| dt \quad (792)$$

Proof. Since $\gamma'(t)$ is continuous, then $|\gamma'(t)|$ is continuous and $|\gamma'(t)|$ is Riemann integrable. Now is P is any partition of $[0, 1]$, then

$$\Lambda(x, P) = \sum_{i=1}^n |\gamma(t_i) - \gamma(t_{i-1})| = \sum_{i=1}^n \left| \int_{t_{i-1}}^{t_i} \gamma'(s) ds \right| \quad (\text{Fund. Thm. of Calc.})$$

$$\leq \sum_{i=1}^n \int_{t_{i-1}}^{t_i} |\gamma'(s)| ds \quad (793)$$

$$= \int_{t_0}^{t_n} |\gamma'(s)| ds \quad (794)$$

So we've proved one inequality. Now we prove the other. Let $\epsilon > 0$ be given. Then since $\gamma'(t)$ is continuous on compact $[0, 1]$, it must be uniformly continuous on $[0, 1]$. So $\exists \delta > 0$ s.t.

$$|s - t| < \delta \implies |\gamma'(s) - \gamma'(t)| < \epsilon \quad (795)$$

Now take a partition P of $[0, 1]$ s.t. $|t_i - t_{i-1}| < \delta$ for each $1 \leq i \leq N$




Figure 30: We can visualize this by partitioning the interval $[a, b]$ into the intervals Δ_i , each with point $\xi_i \in \Delta_i$. This would partition the path to $\Gamma(\Delta_i)$, each with points $\Gamma(\xi_i)$, and at each point $\Gamma(\xi_i)$, we can imagine the velocity vector of the curve. By taking the magnitude of this vector $\Gamma'(\xi_i)$, we multiply it by the length of the interval Δx_i to get one rectangle, creating an approximation for one partition of the path.

Corollary 6.4 (Length of the Graph of a C^1 Function)

An immediate result of this formula is the formula for the length of a graph of a function $f : [a, b] \rightarrow \mathbb{R}$ in \mathbb{R}^2 , by looking at the parameterization $t \mapsto (t, f(t))$.

$$\Lambda(\gamma) = \int_a^b \sqrt{1 + (f'(t))^2} dt \quad (796)$$

The question on the effect of parameterization on the integral now arises.

Definition 6.12 (Admissible Change of Parameter)

The path $\tilde{\Gamma} : [\alpha, \beta] \rightarrow \mathbb{R}^3$ is obtained from $\Gamma : [a, b] \rightarrow \mathbb{R}^3$ by an **admissible change of parameter** if there exists a smooth mapping

$$T : [\alpha, \beta] \rightarrow [a, b] \quad (797)$$

such that $T(\alpha) = a, T(\beta) = b, T'(\tau) > 0$ (that is, the reparameterization T is monotonic) on $[\alpha, \beta]$, and

$$\tilde{\Gamma} = \Gamma \circ T \quad (798)$$

The series of mappings can be represented with the following commutative diagram, where $I_{\alpha, \beta} = [\alpha, \beta] \subset \mathbb{R}$ and $I_{a, b} = [a, b] \subset \mathbb{R}$.

$$\begin{array}{ccc} I_{\alpha, \beta} & \xrightarrow{T} & I_{a, b} \\ & \searrow \tilde{\Gamma} & \downarrow \Gamma \\ & & \mathbb{R}^3 \end{array} \quad (799)$$




Figure 31: Note that the points are labeled 0, 1, 2, 3, 4, 5 do not represent numerical values, but rather the order in which the points are parameterized. We can see from this ordering that T is monotonic.

Theorem 6.25 (Invariance of Arclength Integral under Admissible Change of Parameters)

If a smooth path $\tilde{\Gamma} : [\alpha, \beta] \rightarrow \mathbb{R}^3$ is obtained from a smooth path $\Gamma : [a, b] \rightarrow \mathbb{R}^3$ by an admissible change of parameter, then the lengths of the two paths are equal. That is, a

$$\int_a^b |\Gamma'(t)| dt = \int_\alpha^\beta |\tilde{\Gamma}'(t)| dt := \int_\alpha^\beta |(\Gamma \circ T)'(t)| dt \quad (800)$$

7 Sequences of Functions

Let us slightly extend the notion of limits of sequences with functions.

Definition 7.1 (Pointwise Convergence)

Let $E \subset \mathbb{R}$ and $f, f_n : E \rightarrow \mathbb{R}$. We say that $f_n \rightarrow f$ **pointwise** if

$$\lim_{n \rightarrow \infty} f_n(x) = f(x) \text{ for all } x \in E \quad (801)$$

The next question to ask is whether properties of functions are preserved under the limit operations. For example, if the f_n 's are continuous, differentiable, or Riemann integrable, is the same true for the limit function f ? What are the relations between f'_n and f' or $\int f_n$ and $\int f$? To say that a function f is continuous at a point x means

$$\lim_{t \rightarrow x} f(t) = f(x) \quad (802)$$

For functions, the analogous result is whether

$$\lim_{t \rightarrow x} \lim_{n \rightarrow \infty} f_n(t) = \lim_{n \rightarrow \infty} \lim_{t \rightarrow x} f_n(t) \quad (803)$$

Is this true? Let's look at a few examples.

Example 7.1 (Double Sequence May be Swappable)

Consider the double sequence $(\frac{1}{m+n})_{m,n \in \mathbb{N}}$. We can compute the limit as both $n, m \rightarrow +\infty$ in many ways.

1. We can first set $m \rightarrow +\infty$, then $n \rightarrow +\infty$.
2. We can first set $n \rightarrow +\infty$, then $m \rightarrow +\infty$.
3. We might want to take n twice as slow as m .

All of these converge to the same value of 0, so there is no problem.

Example 7.2 ()

Let $f_n(x) = x/n$. Then $f_n \rightarrow 0$ pointwise, where 0 is the 0 function. This is true since for every fixed x , we can set n so large that $x/n < \epsilon$ for any ϵ .

In this case, we are considering a double sequence in \mathbb{R} . However, if we fix one value, then it becomes a sequence of functions, and we already have established that classes of functions form a vector space. In general, interchanging limits are not allowed.

Example 7.3 (Cannot Exchange Limits Under Pointwise Convergence)

Consider the slightly different sequence $(\frac{m}{m+n})_{m,n}$.

1. If $m \gg n$, i.e. take the sequence of values $(10^k, k)$, then this will approach 1.
2. If $n \gg m$, i.e take the sequence of values $(k, 10^k)$, then this will approach 0.
3. In intermediate cases, you can in fact get any number between 0 and 1.

Therefore, in general, the limits are not equal. Even worse, the properties of these functions are violated. To get equality, we need to make a stronger assumption than simple existence of limits.

Example 7.4 (Pointwise Limit of Continuous Functions is not Continuous)

Let $f_n(x) : [0, 1] \rightarrow \mathbb{R}$ defined $f_n(x) = x^n$. Then,

$$0 \leq x < 1 \implies x^n \rightarrow 0 \text{ as } n \rightarrow \infty \quad (804)$$

$$x = 1 \implies x^n \rightarrow 1 \text{ as } n \rightarrow \infty \quad (805)$$

So,

$$f_n \rightarrow f^*(x) := \begin{cases} 0 & \text{if } x < 1 \\ 1 & \text{if } x = 1 \end{cases} \quad (806)$$

Note that all f_n are continuous but f^* is discontinuous since

$$\lim_{x \rightarrow 1} \lim_{n \rightarrow \infty} f_n(x) = 0 \neq 1 = \lim_{n \rightarrow \infty} \lim_{x \rightarrow 1} f_n(x) \quad (807)$$

Example 7.5 (Pointwise Series Sum of Continuous Functions is not Continuous)

Consider

$$f_n(x) = \frac{x^2}{(1+x^2)^n} \quad (808)$$

and let

$$f(x) = \sum_{n=0}^{\infty} f_n(x) = \sum_{n=0}^{\infty} \frac{x^2}{(1+x^2)^n} \quad (809)$$

Since $f_n(0) = 0$, we have $f(0) = 0$. For $x \neq 0$, the series is a convergent geometric series with sum $1 + x^2$. Therefore,

$$f(x) = \begin{cases} 0 & \text{if } x = 0 \\ 1 + x^2 & \text{if } x \neq 0 \end{cases} \quad (810)$$

Therefore the sum of a convergent series of continuous functions may not be continuous.

Example 7.6 (Cannot Exchange Integrals and Limits Under Pointwise Convergence)

Consider the function $f_n : [0, 1] \rightarrow \mathbb{R}$ defined by $f_n(0) = f_n(1/n) = 0$ and $f_n(1/2n) = 2n$, with everything else linearly interpolated. Then $f_n \rightarrow 0$ since it is constantly 0 at 0 and for every $x > 0$, there exists $1/N < x$ and so $f_n(x) = 0$ for all $n \geq N$. However, $\int_0^1 f_n(x) dx = 1$, so

$$\lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx \neq \int_0^1 \lim_{n \rightarrow \infty} f_n(x) dx \quad (811)$$

So integration is not continuous with respect to the topology induced by pointwise convergence.

The problem is not in the construction of the limits or the integral, but with the pointwise convergence. With pointwise convergence,

1. we cannot exchange two limits, which implies limit of derivatives may not equal to the derivative of the limit
2. limits do not preserve continuity
3. cannot exchange integrals and limits
4. cannot exchange sums

Example 7.7 ()

Review.

$$\left(1 - \frac{1}{n}\right)^n \rightarrow e^{-1} \quad (812)$$

So now the motivation is to find properties of these sequences of functions that do allow these manipulations.

7.1 Uniform Convergence

The problem seems to be that certain points may converge at a much slower rate than others. Therefore, we want to impose some sort of uniform condition, which says that the values of the function at every point must converge.

Definition 7.2 (Uniform Convergence)

Given $f_n : E \rightarrow \mathbb{R}$ of bounded functions, (f_n) is said to **converge uniformly** to a bounded function $f : E \rightarrow \mathbb{R}$ if $\forall \epsilon > 0$, there exists $N \in \mathbb{N}$ s.t.

$$n \geq N \implies |f_n(x) - f(x)| < \epsilon \text{ for all } x \in E \quad (813)$$

the “for all $x \in E$ ” is the uniform part, which is similar to uniform continuity.

Theorem 7.1 (Uniform Convergence iff Supremum of Difference Converges to 0)

This is an immediate consequence of the definition.

$$f_n \rightarrow f \text{ uniformly on } E \iff \lim_{n \rightarrow \infty} \sup_{x \in E} |f_n(x) - f(x)| = 0 \quad (814)$$

Proof. We can prove this with a bunch of iff statements. Let $f_n \rightarrow f$ uniformly, then this is equivalent to $\forall \epsilon > 0$, $\exists N \in \mathbb{N}$ s.t. $n \geq N$ implies

$$|f_n(x) - f(x)| < \epsilon \quad \forall x \in E \quad (815)$$

$$\iff M_n \sup_{x \in E} |f_n(x) - f(x)| \leq \epsilon \quad (816)$$

$$\iff M_n \rightarrow 0 \text{ as } n \rightarrow \infty \quad (817)$$

which implies that $M_n \rightarrow 0$ as $n \rightarrow \infty$.

Generally, to prove uniform convergence, you will need to find that $|f_n(x) - f(x)|$ is bounded by something that is independent of x , and it goes to 0 as $n \rightarrow \infty$. Here is an equivalent condition.

Definition 7.3 (Uniformly Cauchy)

A sequence $f_n : E \rightarrow \mathbb{R}$ is called **uniformly Cauchy** if $\forall \epsilon > 0$, there exists $N \in \mathbb{N}$ s.t. $\forall n, m \geq N$,

$$|f_n(x) - f_m(x)| < \epsilon \text{ for all } x \in E \quad (818)$$

Lemma 7.1 (Cauchy Criterion of Uniform Convergence)

(f_n) converges uniformly iff (f_n) is uniformly Cauchy.

Proof. We prove bidirectionally.

1. (\rightarrow) . Suppose (f_n) converges uniformly on E , and let f be the limit function. Then there exists $N \in \mathbb{N}$ s.t.

$$n \geq N \implies |f_n(x) - f(x)| < \frac{\epsilon}{2} \text{ for all } x \in E \quad (819)$$

Therefore,

$$n, m \geq N \implies |f_n(x) - f_m(x)| \leq |f_n(x) - f(x)| + |f(x) - f_m(x)| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \quad (820)$$

for all $x \in E$.

2. (\leftarrow) . Suppose the uniform Cauchy criterion holds. For every $x \in E$, the sequence $(f_n(x))_n$ converges as a Cauchy sequence in \mathbb{R} . Call this limit $f(x)$. Thus the sequence (f_n) converges pointwise to f .

Now let $\epsilon > 0$. Then, by uniform Cauchy, $\exists N \in \mathbb{N}$ s.t.

$$n, m \geq N \implies |f_n(x) - f_m(x)| < \epsilon \quad \forall x \in E \quad (821)$$

Now, if we fix n^a , we know that

$$m \geq N \implies |f_n(x) - f_m(x)| < \epsilon \quad (822)$$

Now we can take the limit as $m \rightarrow \infty$, and therefore the limit of the LHS must be bounded by that of the RHS.

$$\lim_{m \rightarrow \infty} |f_n(x) - f_m(x)| = |f_n(x) - f(x)| \leq \epsilon \quad (823)$$

Now this is true for all $n \geq N$, which basically is the definition of convergence.^b

^aThis is a classic trick used to convert Cauchy convergence to regular convergence. What we want to do is that since given some x_n , the rest of the points x_m must also be close to x_n , so the limit of the x_m must also be close to x_n , which is basically the definition of convergence.

^bIf you are dissatisfied with the \leq , just set $\epsilon/2$ to get strictly less than.

The wrong proof (that uniformly Cauchy implies uniform convergence) that I had in my first attempt was that I tried to directly bound the distances using some variant of the triangle inequality, like

$$|f_n(x) - f_m(x)| \leq |f_n(x) - f(x)| + |f(x) - f_m(x)| \quad (824)$$

However, this doesn't really help since this is an *upper bound*. Perhaps we can use the reverse triangle inequality.

$$||f_n(x)| - |f_m(x)|| \leq |f_n(x) - f_m(x)| \quad (825)$$

but again, this inequality lacks $f(x)$ entirely.

Example 7.8 ()

We have uniform convergence

$$\frac{\sin(e^n x)}{n} \rightarrow 0 \text{ since } \left| \frac{\sin(e^n x)}{n} \right| \leq \frac{1}{n} \rightarrow 0 \quad (826)$$

Example 7.9 ()

$f_n(x) = x^n$ does not converge uniformly to *any* function in $[0, 1]$. It suffices to find a sequence $(y_n) \subset [0, 1]$ s.t. $|f_n(y_n) - f(y_n)| \not\rightarrow 0$ as $n \rightarrow \infty$. Take $y_n = 1 - \frac{1}{n}$. Then $f_n(y_n) \rightarrow \frac{1}{e} \neq 0$, but $f(y_n) \rightarrow 1$.

Example 7.10 ()

The triangle functions do not converge uniformly since f_n is unbounded, i.e. $f_n(\frac{1}{2n}) = 2n$, while $f_n(\frac{1}{n}) = 0$. So we can construct two sequences that

$$(1/2n) \rightarrow \infty, (1/n) \rightarrow 0 \text{ as } n \rightarrow \infty \quad (827)$$

Lemma 7.2 (Uniform Convergence Implies Pointwise Convergence)

Uniform convergence implies pointwise convergence.

Proof. Say $f_n \rightarrow f$ uniformly. Then for every $\epsilon > 0$, there exists a $N \in \mathbb{N}$ s.t.

$$n \geq N \implies |f_n(x) - f(x)| < \epsilon \text{ for all } x \in E \quad (828)$$

So just fix a point x , take any $\epsilon > 0$, and we have our δ due to uniform convergence.

Theorem 7.2 (Weierstrass M-Test)

Suppose (f_n) is a sequence of functions on E , and suppose

$$|f_n(x)| \leq M_n \text{ for all } x \in E \quad (829)$$

for each $n \in \mathbb{N}$. Then if $\sum_n M_n$ converges, $\sum_n f_n$ converges uniformly.

Proof. If $\sum_n M_n$ converges, then for arbitrary $\epsilon > 0$, we have

$$\left| \sum_{i=n}^m f_i(x) \right| \leq \sum_{i=n}^m M_i \leq \epsilon \quad (830)$$

provided $n, m \in \mathbb{N}$ are large enough. Therefore this is uniformly Cauchy, and so uniformly convergent.

Here is my wrong first attempt for a proof. Assume that $\sum_{k=1}^{\infty} M_k$ converges, and we wish to show that $\forall \epsilon > 0, \exists N \in \mathbb{N}$ s.t.

$$n \geq N \implies \left| \underbrace{\sum_{k=1}^{\infty} f_k(x) - \sum_{k=1}^n f_k(x)}_{\text{not allowed}} \right| = \left| \sum_{k=n+1}^{\infty} f_k(x) \right| < \epsilon \quad (831)$$

The problem is that in our statement, we are assuming that we can actually subtract a finite term $\sum_{k=1}^n f_k(x)$ from a potentially divergent one: $\sum_{k=1}^{\infty} f_k(x)$. Therefore, we are assuming that this is convergent in the first place! This is exactly why we want to work with Cauchy sequences, which doesn't carry this kind of assumption, and so the sum from $i = n$ to m is guaranteed to be finite.

After this, the rest of the steps are fine, since I use the fact that absolutely convergent series are also

convergent, and so

$$\left| \sum_{k=n+1}^{\infty} f_k(x) \right| \leq \sum_{k=n+1}^{\infty} |f_k(x)| \leq \sum_{k=n+1}^{\infty} M_k \quad (832)$$

Example 7.11 (Converse of Weierstrass M-test Not True)

The converse is clearly not true, since we can just take any uniformly convergent bounded sequence and set $M_n \rightarrow +\infty$ as $n \rightarrow \infty$.

The final theorem is a significant one. So far, we uniform convergence to prove that the limit of some sequence satisfies some property. In here, it's sort of the opposite.

Theorem 7.3 (Dini's Theorem)

Suppose K is compact, and

1. (f_n) is a sequence of continuous functions on K .
2. $f_n \rightarrow f$ pointwise on K .
3. f is continuous.^a
4. $f_n \leq f_{n+1}$ for all $n \in \mathbb{N}$.

Then $f_n \rightarrow f$ uniformly on K .

^aNote that the pointwise limit of continuous functions may not be continuous!

Proof. For convenience, let's define $g_n := f - f_n \geq 0$, which is continuous on K . It suffices to prove that $g_n \rightarrow 0$ uniformly; that is, $\forall \epsilon > 0, \exists N \in \mathbb{N}$ s.t.

$$n \geq N \implies |g_n(x)| < \epsilon \quad (833)$$

Fix $\epsilon > 0$, and from compactness, we should already be thinking of trying to construct an open cover of K . Let

$$E_n := \{x \in K \mid g_n(x) < \epsilon\} \quad (834)$$

Note three things.

1. First, E_n is open as the preimage of continuous g_n .
2. Second g_n is decreasing, so $g_n \geq g_{n+1}$, and so

$$E_n \subset E_{n+1} \quad \forall n \in \mathbb{N} \quad (835)$$

3. Third, $g_n \rightarrow 0$ pointwise, so at some point the E_n 's must cover all of K .^a

$$\bigcup_{n=1}^{\infty} E_n = K \quad (836)$$

We have constructed an open cover, and now we take the fact that K is compact to get a finite subcover $\{E_{n_k}\}_k$. But since the E_n 's are nested, we can set $N = \max_k \{n_k\}$, which implies $E_N = K$. Therefore, as long as $n \geq N$, g_n is small enough so that it will map all of $E_N = K$ to a value less than ϵ . We have

$$n \geq N \implies g_n(x) = |g_n(x)| < \epsilon \quad (837)$$

^aRemember that this is for *fixed* ϵ ! This final point is not true if ϵ is not fixed.

Let's reflect on this proof and how we used the theorem's assumptions. First, monotonicity of f_n was needed to make sure that E_n was increasing. Second, compactness of K was needed to extract a finite subcover that we can exploit.

7.2 Limits of Uniformly Convergent Functions

Now we will formalize and prove the manipulations that are unlocked by uniform convergence.

Theorem 7.4 (Limits are Swappable)

Suppose $f_n \rightarrow f$ uniformly over a set E of a metric space. Let $x \in E'$, and suppose that $\lim_{t \rightarrow x} f_n(t) = A_n$ for all n . Then,

1. (A_n) converges, and
2. we have

$$\lim_{t \rightarrow x} f(t) = \lim_{n \rightarrow \infty} A_n \iff \lim_{t \rightarrow x} \lim_{n \rightarrow \infty} f_n(t) = \lim_{n \rightarrow \infty} \lim_{t \rightarrow x} f_n(t) \quad (838)$$

Proof. Let $\epsilon > 0$, then by uniform continuity, there exists a $N \in \mathbb{N}$ s.t.

$$n, m \geq N \implies |f_n(t) - f_m(t)| < \epsilon \quad (839)$$

Now let $t \rightarrow x$, and since the limit exists, we have

$$n, m \geq N \implies |A_n - A_m| < \epsilon \quad (840)$$

and so (A_n) is by definition a Cauchy sequence in \mathbb{R} . Say that $A_n \rightarrow A$.

Now we wish to prove that $\lim_{t \rightarrow x} f(t) = A$. Take $\epsilon > 0$, and we wish to show that there exists some $\delta > 0$ s.t.

$$|t - x| < \delta \implies |f(t) - A| < \epsilon \quad (841)$$

Note that by the triangle inequality

$$|f(t) - A| \leq |f(t) - f_n(t)| + |f_n(t) - A_n| + |A_n - A| \quad (842)$$

Therefore, we can take $\frac{\epsilon}{3} > 0$.

1. By uniform convergence of $f_n \rightarrow f$, there exists a $N_1 \in \mathbb{N}$ s.t.

$$n \geq N_1 \implies |f_n(x) - f(x)| < \frac{\epsilon}{3} \text{ for all } x \in E \quad (843)$$

2. By convergence of $A_n \rightarrow A$, there exists a $N_2 \in \mathbb{N}$ s.t.

$$n \geq N_2 \implies |A_n - A| < \frac{\epsilon}{3} \quad (844)$$

3. Therefore, choose $N = \max\{N_1, N_2\}$, and for this N , by convergence of $f_n(t) \rightarrow A_n$ we can choose a δ s.t.

$$|x - t| < \delta \implies |f_n(t) - A_n| < \frac{\epsilon}{3} \quad (845)$$

This essentially bounds the three values, and so by choosing the δ in (3), we get

$$|t - x| < \delta \implies |f(t) - A| \leq \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon \quad (846)$$

Corollary 7.1 (Uniform Limits of Continuous Functions are Continuous)

If (f_n) is a sequence of continuous functions on E , and if $f_n \rightarrow f$ uniformly on E , then f is continuous on E .

Proof. Using the sequential definition of continuity, we claim that $\lim_{t \rightarrow x} f(t) = f(x)$. We know from the previous theorem that

$$\lim_{t \rightarrow x} \underbrace{\lim_{n \rightarrow \infty} f_n(t)}_{=f(t)} = \lim_{n \rightarrow \infty} \underbrace{\lim_{t \rightarrow x} f_n(t)}_{=f_n(x)} \quad (847)$$

where the LHS follows from (f_n) being uniformly convergent, which implies pointwise convergence, and the RHS follows from f_n being continuous.

However, the converse is not true. A sequence of continuous functions may converge to a continuous function though not uniformly. Now let's move onto integration.

Theorem 7.5 (Limits of Integrals are Integrals of Uniform Limits)

Let (f_n) be a sequence of Riemann integrable functions on $[a, b]$. If $f_n \rightarrow f$ uniformly on $[a, b]$, then $f \in \mathcal{R}([a, b])$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n(x) dx = \int_a^b f(x) dx = \int_a^b \lim_{n \rightarrow \infty} f_n(x) dx \quad (848)$$

Proof. Let $\epsilon_n = \sup |f_n(x) - f(x)|$ with the supremum taken over $a \leq x \leq b$. Then,

$$f_n - \epsilon_n \leq f \leq f_n + \epsilon_n \quad (849)$$

so the upper and lower integrals of f satisfy

$$\int_a^b (f_n - \epsilon_n) dx \leq \int_a^b f(x) dx \leq \int_a^b f(x) dx \leq \int_a^b (f_n + \epsilon_n) dx \quad (850)$$

Hence, we have

$$0 \leq \int_a^b f(x) dx - \int_a^b f_n(x) dx \leq 2\epsilon_n(b-a) \quad (851)$$

and taking $n \rightarrow \infty$ sets $\epsilon_n \rightarrow 0$.

Corollary 7.2 (Series Function May be Integrated Term by Term)

If $f_n \in \mathcal{R}([a, b])$, and f is defined

$$f(x) := \sum_{n=1}^{\infty} f_n(x) \quad (852)$$

with the series converging uniformly on $[a, b]$, then

$$\int_a^b f(x) dx = \sum_{n=1}^{\infty} \int_a^b f_n(x) dx \quad (853)$$

Now we move onto differentiation.

Theorem 7.6 (Limits of Derivatives)

Suppose (f_n) is a sequence of differentiable functions on $[a, b]$. If

1. there exists some $x_0 \in [a, b]$ such that $(f_n(x_0))_n$ converges, and
2. $(f'_n) \rightarrow f'^a$ on $[a, b]$,

Then (f_n) converges uniformly to a function f , with

$$f'(x) = \lim_{n \rightarrow \infty} f'_n(x) \quad (854)$$

^aAt this point, f' is just notation since f isn't even defined.

Proof. Choose $N \in \mathbb{N}$ s.t. for $n, m \geq N$, the following hold

$$|f_n(x_0) - f_m(x_0)| < \frac{\epsilon}{2}, \quad |f'_n(t) - f'_m(t)| < \frac{\epsilon}{2(b-a)} \quad (855)$$

which is possible due to convergence of $f_n(x_0)$ and uniform convergence of the derivative. Then applying the mean value theorem to the function $(f_n - f_m)$ gives

$$|f_n(x) - f_m(x) - f_n(t) + f_m(t)| = (f_n - f_m)'(c)|x - t| \quad (856)$$

for any $x, t \in [a, b]$ and $c \in (x, t)$. However, $(f_n - f_m)'$ is bounded, so

$$|f_n(x) - f_m(x) - f_n(t) + f_m(t)| \leq \frac{|x - t|\epsilon}{2(b-a)} \leq \frac{\epsilon}{2} \quad (857)$$

and therefore we can use the triangle inequality to get

$$|f_n(x) - f_m(x)| \leq |f_n(x) - f_m(x) - f_n(x_0) + f_m(x_0)| + |f_n(x_0) - f_m(x_0)| \quad (858)$$

$$\leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \quad (859)$$

and so f_n is uniformly Cauchy $\iff (f_n)$ converges uniformly.

To show the equality of the limit, we fix a point $x \in [a, b]$ and define

$$\phi_n(t) = \frac{f_n(t) - f_n(x)}{t - x}, \quad \phi(t) = \frac{f(t) - f(x)}{t - x} \quad (860)$$

for $t \in (a, b), t \neq x$. Then,

$$\lim_{t \rightarrow x} \phi_n(t) = f'_n(x) \quad (861)$$

and we can see from the above inequality that

$$|\phi_n(t) - \phi(t)| \leq \frac{\epsilon}{2(b-a)} \quad (862)$$

for $n, m \geq N$. So we can conclude

$$\phi(t) = \lim_{n \rightarrow \infty} \phi_n(t) \quad (863)$$

Proof. Here is an alternative shorter proof of the equality (but not uniform convergence of f_n !) where we assume $f_n \in C^1([a, b])$. Let's call the limit of f'_n to be g to avoid confusion. We know that since f'_n is continuous, it's integrable and by the fundamental theorem of calculus we have

$$f_n(x) - f_n(0) = \int_0^x f'_n(t) dt \quad (864)$$

Since $f'_n \rightarrow g$ uniformly and $f'_n \in C^0$, g is continuous and so we can define the function

$$f(x) := f(0) + \int_0^x g(t) dt \quad (865)$$

But we can see that by uniform convergence, we can swap integrals

$$f(x) - f(0) = \int_0^x g(t) dt \quad (866)$$

$$= \int_0^x \lim_{n \rightarrow \infty} f'_n(t) dt \quad (867)$$

$$= \lim_{n \rightarrow \infty} \int_0^x f'_n(t) dt \quad (868)$$

$$= \lim_{n \rightarrow \infty} f_n(x) - f_n(0) \quad (869)$$

which implies that $f(x) - f(0) = \lim_{n \rightarrow \infty} f_n(x) - f_n(0)$ and so $f(x) = \lim_{n \rightarrow \infty} f_n(x)$. But since f' is continuous, the function f defined above is differentiable, with derivative $f'(x)$ to be whatever function is in the integral, i.e. $g(x)$. So

$$f'(x) = g(x) = \lim_{n \rightarrow \infty} f'_n(x) \quad (870)$$

7.3 Equicontinuous Families

Note that uniform convergence may not be met due to some counterexamples. In general, there are 3 ways that uniform convergence can fail to happen.

1. *Concentration.* Note that x^n as $n \rightarrow \infty$ almost converges except at one point.
2. *Translation.* Consider $f_n(x) = \sin(x - n)$. Then by increasing n we are shifting it to $+\infty$.
3. *Oscillation.* Consider $f_n(x) = \sin(nx)$. As n increases the function oscillates widely. This is sort of like the worst.⁹

We would like uniform convergence, so we want conditions to avoid lack of uniform convergence. Keep in mind to counterexamples. To avoid translation, work with compact space, or if not compact, have the functions decay uniformly. To avoid oscillation, we can bound the derivative, which is a restriction on each function $|f'_n(x)| \leq M$. The Cauchy criterion is too much. To avoid going to infinity, just bound f : $|f_n(x)| \leq M$ for all $n \in \mathbb{N}, x \in X$.

The bounding of derivatives can be a bit strong. We aren't always working with differentiable functions, so we introduce a similar concept.

Definition 7.4 (Equicontinuous Family)

A family of functions \mathcal{F} on E is said to be **equicontinuous** if $\forall \epsilon > 0$ there exists a $\delta > 0$ s.t.

$$|x - y| < \delta \implies |f(x) - f(y)| < \epsilon \quad (871)$$

for all $x, y \in E, f \in \mathcal{F}$.

So this doesn't even depend on f . You can think of this as uniformly continuous for a class of functions that doesn't depend on f . The first class of equicontinuous functions you should know are those with bounded derivatives.

Lemma 7.3 (Functions with Bounded Derivatives Are Equicontinuous)

Fix $M \geq 0$. Then

$$\mathcal{F}_n := \{f : [0, 1] \rightarrow \mathbb{R} \mid |f'(x)| \leq M\} \quad (872)$$

⁹It turns out that this is the same as (2) under the Fourier transform.

is an equicontinuous family.

Proof. For any $f \in \mathcal{F}$, the MVT $|f(x) - f(y)| = |f'(c)(x - y)|$ for some $c \in (x, y)$. But since $f'(c)$ is bounded by M , take $\delta = \epsilon/M$.

Example 7.12 ()

$\mathcal{F} = \{\sin(nx)\}_{n \in \mathbb{N}}$ is not equicontinuous on $[0, 1]$ since

$$\left| \sin\left(n \frac{\pi}{2n}\right) - \sin\left(n \frac{\pi}{n}\right) \right| = 1 \quad (873)$$

for all n . So setting $x_n = \frac{\pi}{2n}$, $y_n = \frac{\pi}{n}$, we have $d(x_n, y_n) \rightarrow 0$ while $d(f(x_n), f(y_n)) \geq 1$. So this is not equicontinuous.

Definition 7.5 (Pointwise Bounded)

Given a sequence of functions (f_n) over E , we say the sequence is **pointwise bounded** if it satisfies each of the equivalent conditions.

1. There exists some function $\phi(x)$ s.t. $|f_n(x)| < \phi(x)$ for all $x \in E, n \in \mathbb{N}$.
2. For every $x \in E$, the sequence $(|f_n(x)|)_n$ is bounded.

Definition 7.6 (Uniformly Bounded)

Given a sequence of functions (f_n) over E , we say the sequence is **uniformly bounded** if there exists some M s.t. $|f_n(x)| \leq M$ for all $x \in E, n \in \mathbb{N}$.

Therefore, uniform boundedness is stronger, since this bound doesn't even depend on x .

Lemma 7.4 (Uniform Boundedness Implies Pointwise Boundedness)

Uniform boundedness implies pointwise boundedness.

Proof. Take $\phi(x) = M$.

We may wonder what the conditions are for the converse. The following theorem gives us those conditions.

Theorem 7.7 (Conditions for Uniform Boundedness)

If K is compact, with

1. f_n is continuous on K for each n
2. f_n pointwise bounded.
3. f_n equicontinuous on K

Then $\{f_n\}$ is uniformly bounded.

Proof.

If (f_n) is pointwise bounded on E and E_1 is a countable subset of E , it is always possible to find a subsequence (f_{n_k}) that converges for every $x \in E_1$. As an intuitive example, suppose

1. $(f_{2n}(q_1))$ converges
2. $(f_{3n}(q_2))$ converges
3. $(f_{5n}(q_3))$ converges
4. $(f_{7n}(q_4))$ converges

So combining the first two, we have that $(f_{6n}(q_i))$ converges for $i = 1, 2$. Continuing on, $(f_{30n}(q_i))$ converges for $i = 1, 2, 3$. But you can't do this infinitely. So if you want a single subsequence s.t. all the sequences converges, we can do

$$f_2, f_6, f_{30}, f_{210}, f_{2310}, \dots \quad (874)$$

Since

1. if you take out f_2 , it is a subsequence of (f_{3n}) which converges for q_2 , and
2. if you also take out f_6 , it is a subsequence of (f_{6n}) which converges for q_1, q_2 , and
3. if you also take out f_{30} , it is a subsequence of (f_{30n}) which converges for q_1, q_2, q_3

so $f_{n_k}(q_i)$ converges for all i . Now let's formalize this argument.

Lemma 7.5 ()

Let (f_n) be a sequence of functions on $[0, 1]$ that's uniformly bounded. Let $\{q_m\}_{m=1}^\infty$ be a countable set of numbers in $[0, 1]$. Then \exists a subsequence (f_{n_k}) for which $f_{n_k}(q_m)$ is convergent for all $m \in \mathbb{N}$.

Proof. Intuitively, if we find a sequence of functions, we want to look at each point—say 1—and look at $(f_n(1))_n$. $(f_n(1))$ is bounded and so contains a convergent subsequence $(f_{n_k}(1))_k$. Now with this subsequence, we look at $(f_{n_k}(0))_k$ which is bounded and therefore $(f_{n_{k_j}}(0))_j$ converges, and $(f_{n_{k_j}}(1))_j$ must converge as a subsequence of convergent $(f_{n_k}(1))_k$. Now do this for all q 's, and we get a single subsequence that converges for all of them.

For ease of notation, let f_{ij} denote the j th term of the i th subsequence. Then there exists $(f_{n,1})_n$ s.t. $(f_{n,1}(q_1))_n$ converges. Take a subsequence $f_{n,2}$ of $f_{n,1}$ s.t. $(f_{n,2}(q_2))_n$ converges. Given $(f_{n,k})_n$, find a subsequence of it, called $(f_{n,k+1})_n$ for which $(f_{n,k+1}(q_{k+1}))_n$ converges. Now $(f_{n,n})_n$ is a subsequence of the original one (n th term of n th subsequence) for which $(f_{q,n})_n$ is eventually a subsequence of $(f_{n,j})_n$ for any fixed j .

Now the Ascoli's theorem gives us conditions to get rid of translation, oscillations, and infinity. To prove the second statement, we will need a lemma, so we state it now, along with providing a neat trick for constructing sequences.

Theorem 7.8 (Arzela-Ascoli's Theorem)

We claim the following.

1. If a sequence of continuous functions $f_n : [0, 1] \rightarrow \mathbb{R}$ (or more generally, over a compact set) converges uniformly, then they form an equicontinuous family.
2. If a sequence of functions $f_n : [0, 1] \rightarrow \mathbb{R}$ is equicontinuous and so uniformly bounded, then it has a uniformly convergent subsequence.

Proof. Assume (f_n) is uniformly convergent. Then it is uniformly Cauchy. To prove equicontinuity, given a $\epsilon > 0$ we need to find a $\delta > 0$ for all the functions. Since f_n is uniformly Cauchy, $\exists N$ s.t. if $n \geq N$, then

$$\sup_{x \in [0, 1]} |f_n(x) - f_N(x)| < \epsilon/3 \quad (875)$$

Consider the first N functions f_1, \dots, f_N . They are all continuous on a compact set and so uniformly continuous. So for each f_i , there exists a δ_i s.t. $|x - y| < \delta_i \implies |f_i(x) - f_i(y)| < \epsilon$. So take $\delta = \frac{1}{3} \min_i \delta_i > 0$. So for all $1 \leq i \leq N$,

$$|x - y| < \delta \implies |f_i(x) - f_i(y)| < \epsilon/3 \quad (876)$$

and for $n \geq N$,

$$|f_n(x) - f_n(y)| \leq \underbrace{|f_n(x) - f_N(x)|}_{< \epsilon/3} + \underbrace{|f_N(x) - f_N(y)|}_{< \epsilon/3} + \underbrace{|f_N(y) - f_n(y)|}_{< \epsilon/3} < \epsilon \quad (877)$$

For the second part, let $E = \mathbb{Q} \cap [0, 1]$. It is a good thing that E is dense in $[0, 1]$. Let (f_n) be an equicontinuous on $[0, 1]$ and uniformly bounded. Due to the lemma, there exists a $(f_{n_k})_k$ so that the f_{n_k} converges pointwise on E (since E is countable). We will now use equicontinuity of $(f_{n_k})_k$ to prove it's uniformly Cauchy on $[0, 1]$, which will imply that it's convergent. To make notation easier we will call $f_{n_k} = g_k$. Let $\epsilon > 0$. Since g_k is equicontinuous, $\exists \delta > 0$ s.t.

$$|x - y| < \delta \implies |g_k(x) - g_k(y)| < \epsilon \quad (878)$$

Since $E = \{q_1, q_2, \dots\}$ is dense in $[0, 1]$, $\{B_\delta(q_i)\}_{i=1}^\infty$ is an open cover of $[0, 1]$. Since $[0, 1]$ is compact, there exists a finite subcover

$$[0, 1] \subset \bigcup_{j=1}^N B_\delta(q_{i_j}) \quad (879)$$

Since $(g_k(q_{i_j}))$ converges for each $1 \leq j \leq N$, there exists M_j s.t.

$$n, m \geq M_j \implies |g_n(q_{i_j}) - g_m(q_{i_j})| < \epsilon \quad (880)$$

Take $M = \max_j M_j$. Now if $m, n \geq M$, given $x \in [0, 1]$ $\exists q_i$ with $1 \leq i \leq N$ so that $x \in B_\delta(q_i)$, and so

$$|g_n(x) - g_m(x)| \leq \underbrace{|g_n(x) - g_n(q_i)|}_{< \epsilon} + \underbrace{|g_n(q_i) - g_m(q_i)|}_{< \epsilon} + \underbrace{|g_m(q_i) - g_m(x)|}_{< \epsilon} < 3\epsilon \quad (881)$$

where the first and third inequalities come from equicontinuity, and the middle come from convergence on E . So by setting $\delta/3$ we are done.

Example 7.13 ()

An important application is in the existence of minimizers/maximizers for optimization problems involving functions. To minimize

$$J(f) = \int_0^1 \sqrt{1 + f'(t)} dt \quad (882)$$

is the length of the curve of $f : [0, 1] \rightarrow \mathbb{R}$. To minimize the length of the curve, we must search over a set of functions. So to use EVT, you must know what the compact subsets of functions.

Ascoli's theorem exactly characterizes these compact subsets. These compact subsets of function spaces is the closure of equicontinuous functions.

Corollary 7.3 ()

A set of functions $K \subset C([0, 1])$ is compact iff it is, under the supremum metric $\sup_{x \in [0, 1]}$,
1. closed

2. bounded
3. equicontinuous

The first two are needed for finite dimensions. The third condition is for function spaces.

Theorem 7.9 (Contraction Mapping Theorem)

Let (X, d) be a metric space with $J : X \rightarrow X$ and let there exist $c < 1$ s.t.

$$d(J(x), J(y)) \leq cd(x, y) \text{ for all } x, y \in X \quad (883)$$

Then there exists a unique $x^* \in X$ s.t. $J(x^*) = x^*$.

7.4 The Stone-Weierstrass Theorem

The Stone-Weierstrass theorem is a bit more general, while the Weierstrass approximation theorem is for polynomials.

Theorem 7.10 (Weierstrass Approximation Theorem)

If $f \in C([a, b])$, there exists a sequence of polynomials (p_n) that converges uniformly to f .

Lemma 7.6 ()

If $f : [0, 1] \rightarrow \mathbb{R}$ is continuously differentiable on $[0, 1]$, then

$$\lim_{n \rightarrow \infty} \int_0^1 f(x) \sin(nx) dx = 0 \quad (884)$$

This means that as $n \rightarrow \infty$, the integral becomes small.

Proof.

Now we prove a stronger version.

Theorem 7.11 ()

For every $f \in C([0, 1])$,

$$\lim_{n \rightarrow \infty} \int_0^1 f(x) \sin(nx) dx = 0 \quad (885)$$

Proof. Let $\epsilon > 0$. Then by the Weierstrass approximation theorem^a, there exists a polynomial $p : [0, 1] \rightarrow \mathbb{R}$ for which

$$\sup_{x \in [0, 1]} |p(x) - f(x)| < \epsilon \quad (886)$$

Now consider

$$\left| \int_0^1 f(x) \sin(nx) dx \right| \leq \left| \int_0^1 p(x) \sin(nx) dx \right| + \left| \int_0^1 (f(x) - p(x)) \sin(nx) dx \right| \quad (887)$$

$$\leq \left| \int_0^1 p(x) \sin(nx) dx \right| + \epsilon \quad (888)$$

$$\leq 2\epsilon \quad (889)$$

where the first inequality is the triangle inequality, the second is due to the Weierstrass approximation theorem, and the third is due to $p(x)$ being infinitely differentiable, and so by the lemma above it is $\leq \epsilon$.

^aaka, the set of polynomials is dense in the set of continuous functions with the supremum metric. Remember polynomial interpolation, which is for a finite number of points. This is a little different.

Theorem 7.12 ()

Proof. Suppose for the sake of contradiction that $\sin(n_k x) \rightarrow g(x)$ uniformly for subsequence $(n_k)_k$. Then $g(x)$ must be continuous on $[0, 1]$. Then

$$\int_a^b g(x)^2 dx = \lim_{n \rightarrow \infty} \int_0^1 g(x) \sin(n_k x) dx = 0 \quad (890)$$

due to the theorem, which implies that $g = 0$. But since $g(nx) = \pm 1$ for some x for all n , we have a contradiction.

7.5 Approximation of the Identity

Definition 7.7 (Approximation of the Identity)

A family of functions $\{\varphi_\epsilon\}$ parameterized by ϵ is called an **approximation of the identity** if

1. $\int_{-\infty}^{\infty} \varphi_\epsilon(y) dy = 1$
2. $\lim_{\epsilon \rightarrow 0} \int_{|y| > \delta} \varphi_\epsilon(y) dy = 0$ for all $\delta > 0$
3. $\varphi_\epsilon \geq 0$.^a

^aThis condition is flexible, but it makes things a bit easier for now.

Example 7.14 ()

Consider the functions f_ϵ satisfying $f(-\epsilon) = f(\epsilon) = 0$, $f(0) = 1/\epsilon$, and everything in between linearly interpolated. Then this is an approximation of the identity.

Example 7.15 ()

Take any $\varphi \geq 0$ s.t. $\int_{-\infty}^{\infty} \varphi = 1$ and define

$$\varphi_\epsilon = \frac{1}{\epsilon} \varphi\left(\frac{x}{\epsilon}\right) \quad (891)$$

which we can think of as squeezing the function horizontally to 0 and making the amplitude very large.

Then we see that

$$\int_{-\infty}^{\infty} \varphi_{\epsilon}(y) dy = \int_{-\infty}^{\infty} \frac{1}{\epsilon} \varphi\left(\frac{y}{\epsilon}\right) = \int_{-\infty}^{\infty} \varphi(x) dx = 1 \quad (892)$$

Fix $\delta > 0$. Then

$$\int_{\delta}^{\infty} \varphi_{\epsilon}(x) dx = \int_{\delta}^{\infty} \frac{1}{\epsilon} \varphi\left(\frac{x}{\epsilon}\right) dx = \int_{\delta/\epsilon}^{\infty} \varphi(y) dy \rightarrow 0 \text{ as } \epsilon \rightarrow 0 \quad (893)$$

Since δ is fixed, we have $\delta/\epsilon \rightarrow +\infty$ as $\epsilon \rightarrow 0$, and so this integral of the tail above converges to 0.

The approximation of the identity (AoI) has an amazing property.

Theorem 7.13 ()

Let $\{\varphi_{\epsilon}\}$ be an AoI. Assume $f : \mathbb{R} \rightarrow \mathbb{R}$ is bounded and continuous. Then

$$\lim_{\epsilon \rightarrow 0} \int_{-\infty}^{\infty} \varphi_{\epsilon}(y) f(y) dy = f(0) \quad (894)$$

Figure 32: The triangle has area 1. Now if you integrate the product of $\varphi_{\epsilon}(y)$ and $f(y)$, it's like taking the product and multiplying by f . But as $\epsilon \rightarrow 0$, the triangle's area is 1, and at the end you just multiply by $f(0)$.

Proof. We have

$$\left| \int_{-\infty}^{\infty} \varphi_{\epsilon}(y) f(y) dy - f(0) \right| = \left| \int_{-\infty}^{\infty} \varphi_{\epsilon}(y) (f(y) - f(0)) dy \right| \quad (895)$$

$$\leq \left| \int_{-\delta}^{\delta} \varphi_{\epsilon}(y) (f(y) - f(0)) dy \right| + \left| \int_{|y| > \delta} \varphi_{\epsilon}(y) (f(y) - f(0)) dy \right| \quad (896)$$

$$\leq \sup_{y \in [-\delta, \delta]} |f(y) - f(0)| + 2M \int_{|y| > \delta} \varphi_{\epsilon}(y) dy \quad (897)$$

where the final step follows from the left integral is less than $\sup_{[-\delta, \delta]} |f(y) - f(0)|$, and for the right integral, we have $f(y) - f(0) \leq 2 \sup |f| \leq 2M$. Since you don't know the limit exists, you take the limsup,

$$\limsup_{\epsilon \rightarrow 0} \left| \int_{-\infty}^{\infty} \varphi_{\epsilon}(y) f(y) dy - f(0) \right| \leq \sup_{y \in [-\delta, \delta]} |f(y) - f(0)| \rightarrow 0 \text{ as } \delta \rightarrow 0 \quad (898)$$

Corollary 7.4 (Convolution)

If $\{f_{\epsilon}\}$ is an AoI with f bounded and continuous and $x \in \mathbb{R}$, we have

$$\lim_{\epsilon \rightarrow 0} \int_{-\infty}^{\infty} \varphi_{\epsilon}(y) f(x - y) dy = f(x) \quad (899)$$

This is called the **convolution** of f with φ_{ϵ} .

Definition 7.8 (Dirac Delta Function)

$\varphi_\epsilon \rightarrow \delta_0$ as $\epsilon \rightarrow 0$, the Dirac delta function. This is a limit.

Theorem 7.14 ()

Consider the functions

$$\varphi_n(x) = \begin{cases} c_n(1-x^2)^n & \text{if } x \in [-1, 1] \\ 0 & \text{else} \end{cases}, \quad c_n = \left(\int_{-1}^1 (1-x^2)^n dx \right)^{-1} \quad (900)$$

Then,

$$\int_{-1}^1 \varphi_n(x) dx = \int_{-\infty}^{\infty} \varphi_n(x) dx = 1 \quad (901)$$

so $\{\varphi_n\}$ is an approximation of the identity.

Proof. Note that c_n is chosen such that $\int_{-\infty}^{\infty} \varphi_n(x) dx = 1$ and $\varphi_n \geq 0$. We want to show

$$\int_{|x|>\delta} \varphi_n(x) dx \rightarrow 0 \text{ as } n \rightarrow \infty \text{ for all } \delta > 0 \quad (902)$$

We claim that $c_n \leq 10\sqrt{n}$. since we wish to upper bound the multiplicative inverse of an integral, it suffices to lower bound the inverse—i.e. the integral itself.

$$\int_{-1}^1 (1-x^2)^n dx = 2 \int_0^1 (1-x^2)^n dx \geq 2 \int_0^{1/\sqrt{n}} (1-x^2)^n dx \quad (903)$$

$$\geq 2 \int_0^{1/\sqrt{n}} (1-nx^2) dx \quad (904)$$

where the last inequality follows from the binomial inequality $(1-s)^n \geq 1-sn$. Therefore, the final integral is now computable, so it equals

$$= 2 \left(x - \frac{nx^3}{3} \right) \Big|_0^{1/\sqrt{n}} = \frac{4}{3} \frac{1}{\sqrt{n}} \implies c_n \leq \left(\frac{4}{3} \frac{1}{\sqrt{n}} \right)^{-1} < \sqrt{n} \quad (905)$$

Now we have

$$\int_{|x|>\delta} \varphi_n(x) dx = 2 \int_{\delta}^{+\infty} \varphi_n(x) dx \quad (906)$$

$$= 2 \int_{\delta}^1 c_n(1-x^2)^n dx \quad (907)$$

$$\leq 20\sqrt{n} \int_{\delta}^1 (1-x^2)^n dx \quad (908)$$

$$\leq 20\sqrt{n}(1-\delta^2)^n \rightarrow 0 \text{ as } n \rightarrow \infty \quad (909)$$

Note that we could have claim that the bound be $c_n \leq (10\sqrt{n})^{1000}$ and this would still be true.

Theorem 7.15 (Stone-Weierstrass Theorem)

Let $f \in C([a, b])$. Then $\forall \epsilon > 0, \exists$ a polynomial p s.t.

$$d(f, p) := \sup_{x \in [a, b]} d(f(x), p(x)) < \epsilon \quad (910)$$

This is equivalent to saying that $\mathbb{R}[x]$ is dense in $C([a, b])$, or that for any $f \in C([a, b])$, \exists sequence (p_n) of polynomials s.t. $p_n \rightarrow p$ uniformly on $[a, b]$.

Proof. By translation and dilation, it suffices to take $[a, b] = [-1, 1]$. This is because translation/dilations are automorphisms (?). It also suffices to consider only f for which $f(-1) > 0$ and $f(1) = k$ for some number k . This is because we can always replace f with \tilde{f} defined by

$$\tilde{f}(x) := f(x) - \left(\frac{(1+x)f(1) + (1-x)f(-1)}{2} \right) \quad (911)$$

Let's extend f by 0 outside of $[-1, 1]$. f is now a bounded continuous function \mathbb{R} implying that f is uniformly continuous. Then we can take the integral.

$$\varphi_n(x) = \begin{cases} \frac{(1-x^2)^n}{\int_{-1}^1 (1-x^2)^n dx} & \text{if } x \in [-1, 1] \\ 0 & \text{else} \end{cases} \quad (912)$$

Since f is bounded, uniformly continuous on \mathbb{R} , and since $\{\varphi\}$ is an approximation of the identity,

$$\int_{-\infty}^{\infty} \varphi_n(t) f(x-t) dt \rightarrow f(x) \quad (913)$$

and so p_n is defined on $[-\frac{1}{2}, \frac{1}{2}]$ with $p_n \rightarrow f$ uniformly.

Now we can use the exact same strategy to prove convergence of Fourier series.

Definition 7.9 (L^2 Inner Product)

The L^2 inner product is defined on $C([a, b])$ as

$$\langle f, g \rangle := \frac{1}{b-a} \int_a^b f(x) \overline{g(x)} dx \quad (914)$$

where f, g are complex valued.

We know that orthonormal bases behave nicely. We present one particularly important one.

Lemma 7.7 ()

The functions $\{e^{inx}\}_{n \in \mathbb{Z}}$ are orthonormal in $C([0, 2\pi])$.

Proof. We have for $n, m \in \mathbb{Z}$

$$\langle e^{inx}, e^{imx} \rangle = \frac{1}{2\pi} \int_0^{2\pi} e^{inx} \overline{e^{imx}} dx = \frac{1}{2\pi} \int_0^{2\pi} e^{inx} e^{-imx} dx = \frac{1}{2\pi} \int_0^{2\pi} e^{i(n-m)x} dx \quad (915)$$

So if $n = m$, then $\langle f, g \rangle = 1$. If not, then

$$\langle f, g \rangle = \frac{1}{2\pi i(n-m)} e^{i(n-m)x} \Big|_0^{2\pi} = 0 \quad (916)$$

and we are done.

Lemma 7.8 ()

Define

$$\varphi_N(x) = \sum_{k=-N}^N e^{ikx} \quad (917)$$

Then, the family $\{\varphi_N\}_{N=0}^{\infty}$ forms an AoI. This is called a *generalized AoI*.

With this, we can prove that any sufficiently smooth (i.e. C^1) functions can be approximated with Fourier series.