

A Book of Abstract Algebra: Solutions to Chapter 5

Tushar Tyagi

May 26, 2016

Notes

A subgroup S is called a subgroup of a group G , if:

1. It is closed on the given operation, i.e. the operation (\cdot) of two elements produces an element $\in S$.
2. It is closed under inverse, i.e. the inverse of each element of S is in S .

Also, each subgroup is a group as well, and therefore follows the three group laws:

1. Associativity
2. Identity
3. Inverse

The *identity*, e of the group is shared by the subgroup.

Trivial & Proper Subgroups

1. The one-element subset $\{e\}$ and the entire group G are the smallest and the largest subgroups of G and are called *trivial subgroups*.
2. All the other subgroups of G are called *proper subgroups*.

Cyclic Groups and Subgroups

If a group (or a subgroup) is generated by a single element, we call that group *Cyclic* and it is written as $\langle a \rangle$, where a is called the *generator* and is the single element which, along with the identity and a^{-1} , can define the entire group.

Defining Equations

A set of equations, involving only the generators and their inverses, is called a set of *defining equations*. These equations can completely define the operation table of the group.

Solutions

Set A

1. $G = \langle R, + \rangle, H = \{ \log a : a \in \mathbb{Q}, a > 0 \}$

- Addition:

Let $a, b \in \mathbb{Q}$

$$\log a + \log b = \log ab$$

$\because a, b \in \mathbb{Q},$

$\therefore ab \in \mathbb{Q}, ab > 0,$

$$\Rightarrow \log ab \in H$$

- Identity:

The identity element would not change the value of $\log a$ under addition. $\log 1$ or 0 is the identity element, since:

If $\log a + \log b = \log a$, then $\log b = 0$, and $b = 1$.

- Inverse:

$$\begin{aligned} \log a + \log a^{-1} &= e \\ \Rightarrow \log a &= -\log a^{-1} \\ \Rightarrow \log a &= \log\left(\frac{1}{a^{-1}}\right) \\ \Rightarrow a &= \frac{1}{a^{-1}} \end{aligned}$$

Since $a \in \mathbb{Q}, \frac{1}{a^{-1}} \in \mathbb{Q}, \therefore \log a^{-1} \in H$

2. $G = \langle R, + \rangle, H = \{\log a : a \in \mathbb{Z}, a > 0\}$

- Addition:

Same reasoning as previous question.

- Inverse:

As calculated in the previous question, $a^{-1} = \frac{1}{a}$ Since $a \in \mathbb{Z}, a^{-1} \notin \mathbb{Z}$

3. $G = \langle R, + \rangle, H = \{x \in \mathbb{R} : \tan x \in \mathbb{Q}\}$

- Addition:

Let $x, y \in \mathbb{R}$.

$$\therefore \tan(x + y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$

If $x = y = 45^\circ$, then $\tan x = \tan y = 1$, which makes the denominator undefined, and therefore addition is not defined for H .

4. $G = \langle R, \cdot \rangle, H = \{2^n 3^m, m, n \in \mathbb{Z}\}$

- Multiplication:

Let $n, m, n', m' \in \mathbb{Z}$.

$$2^n 3^m \cdot 2^{n'} 3^{m'} = 2^{n+n'} 3^{m+m'}$$

$$\therefore n + n', m + m' \in \mathbb{Z}$$

$$\therefore 2^{n+n'} 3^{m+m'} \in H$$

- Inverse:

Since $2^n 3^m \in \mathbb{Z}$,

The inverse is $\frac{1}{2^n 3^m} = 2^{-n} 3^{-m} \therefore -n, -m \in \mathbb{Z}, \therefore 2^{-n} 3^{-m} \in \mathbb{Z}$

5. $G = \langle R \times R, + \rangle, H = \{(x, y) : y = 2x\}$

- Addition:

$$(x, y) + (x', y') = (x + x', y + y')$$

$$\therefore x, x', y, y' \in \mathbb{R},$$

$$\therefore x + x', y + y' \in \mathbb{R}$$

$$\therefore (y + y') = 2(x + x')$$

- Inverse:
 $e = (0, 0)$ Inverse: $(-x, -y)$
 $\because y = 2x$
 $\Rightarrow -y = -2x$
 $\Rightarrow y' = 2x'$

6. $G = \langle R \times R, + \rangle, H = \{(x, y) : x^2 + y^2 > 0\}$

- Addition
 $(x, y) + (x', y') = (x + x', y + y')$
 $\Rightarrow (x + x')^2 + (y + y')^2 > 0,$
 \therefore Addition operation is defined for H .
- Inverse:
 $\because e = (0, 0)$
Inverse is $(-x, -y)$
Let $x, y \in H$,
 $\therefore x^2 + y^2 > 0,$
 $\Rightarrow (-x)^2 + (-y)^2 > 0 \in H$
So inverse is defined.

7. Let C and D be sets, with $C \subseteq D$. Prove that P_C is a subgroup of P_D
In a way, $G = \langle P_D, + \rangle, H = \langle P_C, + \rangle$

- Identity:
(common to both P_C and P_D): $\{\phi\}$
- Inverse:
 $A^{-1} = A$
We have already proved this in Chapter 3, Exercise C.
- Addition:
Let $A, B \subseteq H$,
 $A + B = (A - B) \cup (B - A)$ Since $A \subseteq P_C, (A - B) \subseteq P_C$. Similarly,
 $(B - A) \subseteq P_C$. So the operation of symmetric difference is closed
on subgroup H .