

Skin-related Sensory Processes

Electrosensation and mechanosensation

Topics we will be covering:

- General sensory system stuff
 - What is the general structure of a sensory cell
 - How do they fit into the bigger picture of the nervous system
- Skin sensory systems
 - Electrosensation
 - Passive and active
 - Mechanosensation
- How does the central nervous system process sensory inputs?

General features of sensory receptors

General features of sensory receptors

- How are sensory cells structured?
- How do sensory cells fit into the bigger picture of the nervous system?


How are sensory cells structured?

How are sensory cells structured?

- There are two ways cells that receive sensory input can be structured:



How are sensory cells structured?

- There are two ways cells that receive sensory input can be structured:
 - The sensory cell itself can be a specialized **neuron** that sends an action potential to the central nervous system




How are sensory cells structured?

- There are two ways cells that receive sensory input can be structured:
 - The sensory cell itself can be a specialized **neuron** that sends an action potential to the central nervous system
 - Or it can be a **non-neuronal sensory cell** that connects to an afferent neuron that sends signals to the central nervous system




How do sensory cells fit into the nervous system?

- Either way, the sensory cell must send signals:




How do sensory cells fit into the nervous system?

- Either way, the sensory cell must send signals:
 - Via the afferent neurons of the peripheral nervous system



How do sensory cells fit into the nervous system?

- Either way, the sensory cell must send signals:
 - Via the afferent neurons of the peripheral nervous system
 - Into the central nervous system (spine and brain) for **interpretation and processing**



Interpretation and Processing of Sensory Inputs

Interpretation and Processing of Sensory Inputs

- How does an organism distinguish between different types of stimulus?
- How do sensory cells respond to prolonged stimuli?
- How do sensory cells respond to stimuli of different intensities?
- How does the central nervous system interpret sensory input?

Interpretation and Processing of Sensory Inputs

- How does an organism distinguish between different types of stimulus?
- How do sensory cells respond to prolonged stimuli?
- How do sensory cells respond to stimuli of different intensities?
- How does the central nervous system interpret sensory input?

Again, I'm going in a slightly different order than the lecture did, which I think will make much more sense.

Good resource:

http://droualb.faculty.mjc.edu/Course%20Materials/Physiology%20101/Chapter%20Notes/Fall%202007/chapter_10%20Fall%202007.htm

How do we distinguish different types of stimulus?

How do we distinguish different types of stimulus?

- We are able to feel many different kinds of things with our skin:

How do we distinguish different types of stimulus?

- We are able to feel many different kinds of things with our skin:
 - A firm poke

How do we distinguish different types of stimulus?

- We are able to feel many different kinds of things with our skin:
 - A firm poke
 - A finger brushing along the skin

How do we distinguish different types of stimulus?

- We are able to feel many different kinds of things with our skin:
 - A firm poke
 - A finger brushing along the skin
 - A breeze blowing across our skin


How do we distinguish different types of stimulus?

- We are able to feel many different kinds of things with our skin:
 - A firm poke
 - A finger brushing along the skin
 - A breeze blowing across our skin
- Why can we feel the difference between these stimuli?




How do we distinguish different types of stimulus?

- Why can we feel the difference between these stimuli?
 - Each stimulus activates different combinations of mechanoreceptors.



How do we distinguish different types of stimulus?

- Why can we feel the difference between these stimuli?
 - Each stimulus activates different combinations of mechanoreceptors.
 - Poke: activates Merkel cells strongly, activates zigzag hair follicles moderately




How do we distinguish different types of stimulus?

- Why can we feel the difference between these stimuli?
 - Each stimulus activates different combinations of mechanoreceptors.
 - Poke: activates Merkel cells strongly, activates zigzag hair follicles moderately
 - Stroke: activates Merkel cells mildly, activates zigzag hair follicles strongly




How do we distinguish different types of stimulus?

- Why can we feel the difference between these stimuli?
 - Each stimulus activates different combinations of mechanoreceptors.
 - Poke: activates Merkel cells strongly, activates zigzag hair follicles moderately
 - Stroke: activates Merkel cells mildly, activates zigzag hair follicles strongly
 - Breeze: Merkel cells not active, activates zigzag hair follicles strongly



How do we distinguish different types of stimulus?

- Why can we feel the difference between these stimuli?
 - Each stimulus activates different combinations of mechanoreceptors.
 - Poke: activates Merkel cells strongly, activates zigzag hair follicles moderately
 - Stroke: activates Merkel cells mildly, activates zigzag hair follicles strongly
 - Breeze: Merkel cells not active, activates zigzag hair follicles strongly
 - **Different combinations create different sensations.**



How do sensory cells respond to prolonged stimuli?

How do sensory cells respond to prolonged stimuli?

- Some types of stimuli go on for a long time at a fairly constant rate

How do sensory cells respond to prolonged stimuli?

- Some types of stimuli go on for a long time at a fairly constant rate
 - For example, the sensation of your pants touching your legs.

How do sensory cells respond to prolonged stimuli?

- Some types of stimuli go on for a long time at a fairly constant rate
 - For example, the sensation of your pants touching your legs.
- Different sensory cells respond to prolonged, constant stimuli in different ways.

How do sensory cells respond to prolonged stimuli?


- Some types of stimuli go on for a long time at a fairly constant rate
 - For example, the sensation of your pants touching your legs.
- Different sensory cells respond to prolonged, constant stimuli in different ways.
- Two types of response:

How do sensory cells respond to prolonged stimuli?

- Some types of stimuli go on for a long time at a fairly constant rate
 - For example, the sensation of your pants touching your legs.
- Different sensory cells respond to prolonged, constant stimuli in different ways.
- Two types of response:
 1. Tonic:


How do sensory cells respond to prolonged stimuli?

- Some types of stimuli go on for a long time at a fairly constant rate
 - For example, the sensation of your pants touching your legs.
- Different sensory cells respond to prolonged, constant stimuli in different ways.
- Two types of response:
 1. Tonic: slowly-adapting sense receptors that continue to fire action potentials for as long as the stimulus is occurring.




How do sensory cells respond to prolonged stimuli?

- Some types of stimuli go on for a long time at a fairly constant rate
 - For example, the sensation of your pants touching your legs.
- Different sensory cells respond to prolonged, constant stimuli in different ways.
- Two types of response:
 1. Tonic: slowly-adapting sense receptors that continue to fire action potentials for as long as the stimulus is occurring.
 - a. Ex: pain receptors.




How do sensory cells respond to prolonged stimuli?

- Some types of stimuli go on for a long time at a fairly constant rate
 - For example, the sensation of your pants touching your legs.
- Different sensory cells respond to prolonged, constant stimuli in different ways.
- Two types of response:
 1. Tonic: slowly-adapting sense receptors.
 2. Phasic: rapidly-adapting receptors that only fire when a stimulus begins or changes.





How do sensory cells respond to prolonged stimuli?

- Some types of stimuli go on for a long time at a fairly constant rate
 - For example, the sensation of your pants touching your legs.
- Different sensory cells respond to prolonged, constant stimuli in different ways.
- Two types of response:
 1. Tonic: slowly-adapting sense receptors.
 2. Phasic: rapidly-adapting receptors.
 - a. Ex: Pacinian corpuscles (sense vibrations)



How do sensory cells respond to prolonged stimuli?

- Two types of response:
 1. Tonic: slowly-adapting sense receptors.
 2. Phasic: rapidly-adapting receptors.
- You should understand these two types of response
- You should know, for each mechanoreceptor we will cover, whether it is tonic or phasic.



How do sensory cells respond to stimuli of different intensities?

How do sensory cells respond to stimuli of different intensities?

- Your arm can be poked gently, or very hard.

How do sensory cells respond to stimuli of different intensities?


- Your arm can be poked gently, or very hard.
 - You can distinguish the different intensity of stimulus. How?

How do sensory cells respond to stimuli of different intensities?


- Your arm can be poked gently, or very hard.
 - You can distinguish the different intensity of stimulus. How?
- The traditional model of intensity discrimination is called **frequency coding**.

How do sensory cells respond to stimuli of different intensities?


- Your arm can be poked gently, or very hard.
 - You can distinguish the different intensity of stimulus. How?
- The traditional model of intensity discrimination is called **frequency coding**.
- As stimulus intensity increases, the rate of action potential firing increases.



mild
stimulus




moderate
stimulus



strong
stimulus

How do sensory cells respond to stimuli of different intensities?


- Your arm can be poked gently, or very hard.
 - You can distinguish the different intensity of stimulus. How?
- The traditional model of intensity discrimination is called **frequency coding**.
- As stimulus intensity increases, the rate of action potential firing increases.



mild
stimulus

moderate
Stimulus

strong
stimulus




How do sensory cells respond to stimuli of different intensities?

- You
-
- Th
- As

Take-home message:

The action potential **firing rate** is what encodes information about **stimulus intensity**.




In other words, **stimulus intensity** is encoded in the **firing rate**.

mild stimulus more intense stimulus very intense stimulus




How do sensory cells respond to stimuli of different intensities?


- Your arm can be poked gently, or very hard.
 - You can distinguish the different intensity of stimulus. How?
- The traditional model of intensity discrimination is called **frequency coding**.
- As stimulus intensity increases, the rate of action potential firing increases.




mild
stimulus



moderate
Stimulus




strong
stimulus




How do sensory cells respond to stimuli of different intensities?

- Your arm can be poked gently, or very hard.
 - You can distinguish the different intensity of stimulus. How?
- The traditional model of intensity discrimination is called **frequency coding**.
- As stimulus intensity increases, the rate of action potential firing increases.
- **Note some features of this relationship:**




How do sensory cells respond to stimuli of different intensities?

- Your arm can be poked gently, or very hard.
 - You can distinguish the different intensity of stimulus. How?
- The traditional model of intensity discrimination is called **frequency coding**.
- As stimulus intensity increases, the rate of action potential firing increases.
- Note some features of this relationship:
 - The **threshold**: minimum stimulus intensity required to trigger an action potential.




How do sensory cells respond to stimuli of different intensities?

- Your arm can be poked gently, or very hard.
 - You can distinguish the different intensity of stimulus. How?
- The traditional model of intensity discrimination is called **frequency coding**.
- As stimulus intensity increases, the rate of action potential firing increases.
- Note some features of this relationship:
 - The **threshold**: minimum stimulus intensity required to trigger an action potential.
 - The **saturation point**: maximum stimulus intensity where action potential firing no longer increases




How do sensory cells respond to stimuli of different intensities?

- Your arm can be poked gently, or very hard.
 - You can distinguish the different intensity of stimulus. How?
- The traditional model of intensity discrimination is called **frequency coding**.
- As stimulus intensity increases, the rate of action potential firing increases.
- Note some features of this relationship:
 - The **threshold**: minimum stimulus intensity required to trigger an action potential.
 - The **saturation point**: maximum stimulus intensity where action potential firing no longer increases
 - **Dynamic range**: the range of stimuli that produce a range of nerve responses.




How do sensory cells respond to stimuli of different intensities?

- As stimulus intensity increases, the rate of action potential firing increases.
- Note some features of this relationship:
 - The **threshold**: minimum stimulus intensity required to trigger an action potential.
 - The **saturation point**: maximum stimulus intensity where action potential firing no longer increases
 - **Dynamic range**: the range of stimuli that produce a range of nerve responses.
 - Different types of receptors will have different **thresholds** and different **dynamic ranges** (different levels of sensitivity).




How do sensory cells respond to stimuli of different intensities?

- Different types of receptors will have different **thresholds** and different **dynamic ranges** (different levels of sensitivity).
- Which line represents the receptor with the **highest threshold**?




How do sensory cells respond to stimuli of different intensities?

- Different types of receptors will have different **thresholds** and different **dynamic ranges** (different levels of sensitivity).
- Which line represents the receptor with the **highest threshold**?




How do sensory cells respond to stimuli of different intensities?

- Different types of receptors will have different **thresholds** and different **dynamic ranges** (different levels of sensitivity).
- Which line represents the receptor with the **lowest threshold**?




How do sensory cells respond to stimuli of different intensities?

- Different types of receptors will have different **thresholds** and different **dynamic ranges** (different levels of sensitivity).
- Which line represents the receptor with the **lowest threshold**?




How do sensory cells respond to stimuli of different intensities?

- Different types of receptors will have different **thresholds** and different **dynamic ranges** (different levels of sensitivity).
- Which line represents the receptor with the **smallest dynamic range**?




How do sensory cells respond to stimuli of different intensities?

- Different types of receptors will have different **thresholds** and different **dynamic ranges** (different levels of sensitivity).
- Which line represents the receptor with the **smallest dynamic range**?




How do sensory cells respond to stimuli of different intensities?

- Different types of receptors will have different **thresholds** and different **dynamic ranges** (different levels of sensitivity).
- Which line represents the receptor with the **largest dynamic range**?





How do sensory cells respond to stimuli of different intensities?

- Different types of receptors will have different **thresholds** and different **dynamic ranges** (different levels of sensitivity).
- Which line represents the receptor with the **largest dynamic range**?



Intensity sensing: take-aways

- You should understand what **frequency coding** means
- You should understand the **three features** of a frequency coding graph
- You should know that **different receptors have different threshold and dynamic ranges**, and what that means about their **sensitivity**.



How does the CNS interpret sensory inputs?


- Sensory processing involves more than just the sensory receptor neurons.

How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:


How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:
 - First: sensory neurons




How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:
 - First: sensory neurons
 - Second: interneurons




How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:
 - First: sensory neurons
 - Second: interneurons




How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:
 - First: sensory neurons
 - Second: interneurons
 - Third: interneurons




How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:
 - First: sensory neurons
 - Second: interneurons
 - Third: interneurons
 - Fourth: can still be interneurons, or the somatosensory cortex in the brain.




How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:
 - First: sensory neurons
 - Second: interneurons
 - Third: interneurons
 - Fourth: can still be interneurons, or the somatosensory cortex in the brain.
 - The fourth order in the cortex allows for **integration** and **processing** (through feedback loops - excitatory and/or inhibitory)



How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:
 - First: sensory neurons
 - Second: interneurons
 - Third: interneurons
 - Fourth: can still be interneurons, or the somatosensory cortex in the brain.
 - The fourth order in the cortex allows for **integration** and **processing** (through feedback loops - excitatory and/or inhibitory)
 - The signals are organized in the thalamus before heading to the cortex.



How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:
 - First: sensory neurons
 - Second: interneurons
 - Third: interneurons
 - Fourth: can still be interneurons, or the somatosensory cortex in the brain.
 - The fourth order in the cortex allows for **integration** and **processing** (through feedback loops - excitatory and/or inhibitory)
 - The signals are organized in the thalamus before heading to the cortex.

Signals from each sensory neuron travel **in parallel**

How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:
 - First: sensory neurons
 - Second: interneurons
 - Third: interneurons
 - Fourth: can still be interneurons, or the somatosensory cortex in the brain.
 - The fourth order in the cortex allows for **integration** and **processing** (through feedback loops - excitatory and/or inhibitory)
 - The signals are organized in the thalamus before heading to the cortex.

Signals from each sensory neuron travel **in parallel**

This allows the somatosensory cortex to use **receptor type, location, phasic/tonic properties, receptive field overlaps, conduction velocity** and other events happening nearby over time to make **specific interpretations** about the type of stimulus.

How does the CNS interpret sensory inputs?

- Sensory processing involves more than just the sensory receptor neurons.
- It can be thought of as a four-order process:
 - First: sensory neurons
 - Second: interneurons
 - Third: interneurons
 - Fourth: can still be interneurons, or the somatosensory cortex in the brain.
- Lots of signals are **ignored** by the somatosensory cortex so that only critical stimuli get attention.

What is lateral inhibition?

What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.

What is lateral inhibition?



- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Why is this important?

What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Why is this important?
 - This creates a stronger contrast across space, allowing better discrimination of sensory stimuli.



What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Why is this important?
 - This creates a stronger contrast across space, allowing better discrimination of sensory stimuli.





What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Imagine a stimulus stretching the skin.





What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Imagine a stimulus stretching the skin.
- Many stretch receptors will be under the area being stretched.





What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Imagine a stimulus stretching the skin.
- Many stretch receptors will be under the area being stretched.





What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Imagine a stimulus stretching the skin.
- Many stretch receptors will be under the area being stretched.
- This could lead to widespread, nonspecific action potential firing.





What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Many stretch receptors will be under the area being stretched.
- This could lead to widespread, nonspecific action potential firing.
- As a result, the brain couldn't pinpoint stimulus location.





What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Many stretch receptors will be under the area being stretched.
- Lateral inhibition means that the central, most-excited neuron inhibits its neighbors.





What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Many stretch receptors will be under the area being stretched.
- Lateral inhibition means that the central, most-excited neuron inhibits its neighbors.





What is lateral inhibition?


- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Many stretch receptors will be under the area being stretched.
- Lateral inhibition means that the central, most-excited neuron inhibits its neighbors.
- This creates finer spatial discrimination.




What is lateral inhibition?

- Lateral inhibition is the ability of an excited neuron to reduce the activity of its neighbors.
- Many stretch receptors will be under the area being stretched.
- Lateral inhibition means that the central, most-excited neuron inhibits its neighbors.
- This creates finer spatial discrimination.







without lateral inhibition



VS.



with lateral inhibition



What is lateral inhibition?

- It is important to note that lateral inhibition does not occur **only at the level of the sensory neurons.**


What is lateral inhibition?

- It is important to note that lateral inhibition does not occur **only at the level of the sensory neurons**.
- It also happens at the **second order** of sensory processing, via **inhibitory interneurons**.

What is lateral inhibition?


- It is important to note that lateral inhibition does not occur **only at the level of the sensory neurons.**
- It also happens at the **second order** of sensory processing, via **inhibitory interneurons.**

firing
rate




What is lateral inhibition?

- It is important to note that lateral inhibition does not occur **only at the level of the sensory neurons**.
- It also happens at the **second order** of sensory processing, via **inhibitory interneurons**.
- The central afferent neuron fires more rapidly than its neighbors.




What is lateral inhibition?

- It is important to note that lateral inhibition does not occur **only at the level of the sensory neurons**.
- It also happens at the **second order** of sensory processing, via **inhibitory interneurons**.
- The central afferent neuron fires more rapidly than its neighbors.
- This triggers inhibitory interneurons.




What is lateral inhibition?

- It is important to note that lateral inhibition does not occur **only at the level of the sensory neurons**.
- It also happens at the **second order** of sensory processing, via **inhibitory interneurons**.
- This triggers inhibitory interneurons.
- They inhibit the neighbors more.




What is lateral inhibition?

- It is important to note that lateral inhibition does not occur **only at the level of the sensory neurons**.
- It also happens at the **second order** of sensory processing, via **inhibitory interneurons**.
- This triggers inhibitory interneurons.
- They inhibit the neighbors more.
- And strengthen central firing.




What is lateral inhibition?

- It is important to note that lateral inhibition does not occur **only** at the level of the sensory neurons.
- It also happens at the **second order** of sensory processing, via **inhibitory interneurons**.
- Afferent neurons synapse onto second order neurons.



What is lateral inhibition?

- It is important to note that lateral inhibition does not occur **only** at the level of the sensory neurons.
- It also happens at the **second order** of sensory processing, via **inhibitory interneurons**.
- Afferent neurons synapse onto second order neurons.
- These neurons also show the pattern of lateral inhibition.



Lateral inhibition: takeaways

- How does position within a receptive field affect action potential frequency/
spike rate?
More central = more frequent
- What is lateral inhibition?
- Why is it important?
Allows fine spatial discrimination
- Physiologically, how does it happen?
Excited neuron inhibits neighbors
- Does it only involve first order processing?
No, second order interneurons too.

What is population coding?

What is population coding?

- It is the integration of information from **populations of neurons** to interpret sensory stimuli.


What is population coding?

- It is the integration of information from **populations** of neurons to interpret sensory stimuli.
- Signals from multiple neurons, travelling to the brain **in parallel**, allow the brain to **pinpoint the location** of a stimulus.



What is population coding?

- It is the integration of information from **populations** of neurons to interpret sensory stimuli.
- Signals from multiple neurons, travelling to the brain **in parallel**, allow the brain to **pinpoint the location** of a stimulus.



From Dr. Selvaraj:

“Frequency coding, population coding and lateral inhibition are core concepts to be understood for solving problems.


For the CNS circuits, they should know that there are 3 orders of transmission and that it is channeled through the thalamus to the cortex, the 4th order being processing within the cortex.”

Electrosensation

Electrosensation


- How can organisms sense electricity?
- What is the difference between active and passive electrosensation?

Why would organisms need to sense electricity?




Why would organisms need to sense electricity?

- Animals immersed in water passively produce electrical fields in the water around them due to muscle and nerve activity, and ion pumps in gills.




Why would organisms need to sense electricity?

- Animals immersed in water passively produce electrical fields in the water around them due to muscle and nerve activity, and ion pumps in gills.
- Therefore, electroreception would be advantageous for predators, especially in environments where vision is limited.



Why would organisms need to sense electricity?

- Animals immersed in water passively produce electrical fields in the water around them due to muscle and nerve activity, and ion pumps in gills.
- Therefore, electroreception would be advantageous for predators, especially in environments where vision is limited.
- Some organisms also communicate via electrical signals.




How do organisms sense electricity?

How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems**
 - **Active systems**


How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.




How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - Do not produce any intentional electrical signals of their own



How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - Do not produce any intentional electrical signals of their own
 - Analogous to our light vision: we simply see light reflected from other things, we don't produce any light of our own.



How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - Do not produce any intentional electrical signals of their own
 - Analogous to our light vision: we simply see light reflected from other things, we don't produce any light of our own.
 - **How is the electricity sensed?**

How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - Do not produce any intentional electrical signals of their own
 - Analogous to our light vision: we simply see light reflected from other things, we don't produce any light of our own.
 - **How is the electricity sensed?**
 - **Via ampullary receptors (Ampullae of Lorenzini)**




How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - Do not produce any intentional electrical signals of their own
 - Analogous to our light vision: we simply see light reflected from other things, we don't produce any light of our own.
 - **How is the electricity sensed?**
 - **Via ampullary receptors (Ampullae of Lorenzini)**
 - Clusters of jelly-filled pores; jelly is high in **keratan sulfate**



How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - Do not produce any intentional electrical signals of their own
 - Analogous to our light vision: we simply see light reflected from other things, we don't produce any light of our own.
 - **How is the electricity sensed?**
 - **Via ampullary receptors (Ampullae of Lorenzini)**
 - Clusters of jelly-filled pores; jelly is high in **keratan sulfate**
 - Electrical currents in the water move through the jelly and trigger very sensitive calcium channels, causing an action potential



The lightning bolt is an artist's interpretation- this is NOT how a physicist would depict this!

How do organisms sense electricity?

- There are two types of electrosensory systems:

- Passive systems: passively detect electricity produced by other organisms

Sharks rely on this sensory system so much that they will preferentially attack electric diodes over dead, chopped up fish!




(slide 8 of lecture 20)

... currents in the water move through the jelly and trigger very sensitive calcium channels, causing an action potential

The lightning bolt is an artist's interpretation- this is NOT how a physicist would depict this!

How do organisms sense electricity?


- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - **Active systems:**



Ex: elephant nose fish


How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - **Active systems:** actively produces pulses or waves of electricity,




How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - **Active systems:** actively produces pulses or waves of electricity, and sense how objects in the environment distort those signals



How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - **Active systems:** actively produces pulses or waves of electricity, and sense how objects in the environment distort those signals
 - Analogous to how a bat uses echolocation to sense objects around it





How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - **Active systems:** actively produces pulses or waves of electricity, and sense how objects in the environment distort those signals
 - Analogous to how a bat uses echolocation to sense objects around it
 - **How is distortion of electrical signals sensed?**



How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - **Active systems:** actively produces pulses or waves of electricity, and sense how objects in the environment distort those signals
 - Analogous to how a bat uses echolocation to sense objects around it
 - **How is distortion of electrical signals sensed?**
 - **Tuberous receptors** detect changes in a) the **timing** of the returning electrical impulse



How do organisms sense electricity?

- There are two types of electrosensory systems:
 - **Passive systems:** passively detect electricity produced by other living organisms.
 - **Active systems:** actively produces pulses or waves of electricity, and sense how objects in the environment distort those signals
 - Analogous to how a bat uses echolocation to sense objects around it
 - **How is distortion of electrical signals sensed?**
 - **Tuberous receptors** detect changes in a) the **timing** of the returning electrical impulse or b) the **amplitude**



How do organisms sense electricity?





- How is distortion of electrical signals sensed?
 - Tuberous receptors detect changes in a) the **timing** of the returning electrical impulse or b) the **amplitude**
- You should know these two examples from class:

- Moray Eels:

- Create electrical pulses at irregular intervals
- Pay attention to the timing **and** amplitude of what comes back to figure out what's in the environment

- Elephant fish:

- Create regularly-timed electrical pulses
- Can vary the amplitude of the pulse
- Sense the difference between the amplitude of what it sends out and the amplitude that comes back, and use that to figure out what objects are in the environment




Ribbon synapses

Ribbon synapses

- Some sharks, skates and rays have a special kind of synapse between their electrosensory cells and afferent neurons: **the ribbon synapse**.

Ribbon synapses


- Some sharks, skates and rays have a special kind of synapse between their electrosensory cells and afferent neurons: **the ribbon synapse**.



- These synapses are characterized by a **ribbon**, a cellular structure with many **neurotransmitter vesicles** attached to it.

Ribbon synapses


- Some sharks, skates and rays have a special kind of synapse between their electrosensory cells and afferent neurons: **the ribbon synapse**.



- These synapses are characterized by a **ribbon**, a cellular structure with many **neurotransmitter vesicles** attached to it.
- This facilitates extremely rapid, precise, sustained neuronal communication
 - Lots of neurotransmitters can be released very rapidly

Ribbon synapses

- Some sharks, skates and rays have a special kind of synapse between their electrosensory cells and afferent neurons: **the ribbon synapse**.



- These synapses are characterized by a **ribbon**, a cellular structure with many **neurotransmitter vesicles** attached to it.
- This facilitates extremely rapid, precise, sustained neuronal communication
 - Lots of neurotransmitters can be released very rapidly
- For more info: Bellono, N.W., Leitch, D.B. & Julius, D. Molecular tuning of electroreception in sharks and skates. *Nature* 558, 122–126 (2018)
doi:10.1038/s41586-018-0160-9

From Dr. Selvaraj:

“Students should know about the electrosensory system (active and passive), the two types of active electrolocation.”

Mechanosensation


Mechanosensation

- How do cells turn a mechanical stimulus into a neural signal?
- How can organisms sense many different mechanical stimuli?

Mechanoreceptors have unique ion channels


Mechanoreceptors have unique ion channels

- Mechanoreceptors must translate mechanical stimuli into a neural signal




Mechanoreceptors have unique ion channels

- Mechanoreceptors must translate mechanical stimuli into a neural signal
- They do this with **special mechanosensitive ion channels**




Mechanoreceptors have unique ion channels

- Mechanoreceptors must translate mechanical stimuli into a neural signal
- They do this with **special mechanosensitive ion channels**
 - There are two groups of these channels whose names you should know:
 - Transient receptor potential channels
 - Piezo genes




Mechanoreceptors have unique ion channels

- Mechanoreceptors must translate mechanical stimuli into a neural signal
- They do this with **special mechanosensitive ion channels**
 - There are two groups of these channels whose names you should know:
 - **Transient receptor potential channels**
 - Mechanical force causes second messenger release which causes a TRPV1 channel to open
 - **Piezo genes**



Mechanoreceptors have unique ion channels

- Mechanoreceptors have unique ion channels
- They do this via TRPV1 channels. Dr. Selvaraj specifically mentioned this in his study guide.
 - There are TRPV1 channels
 - Transient receptor potential channels
 - Mechanical force causes second messenger release which causes a TRPV1 channel to open
 - Piezo genes



Mechanoreceptors have unique ion channels

-
-
- For more info if you're curious:


now:

<https://www.britannica.com/science/transient-receptor-potential-channel>

[https://www.cell.com/current-biology/pdf/S0960-9822\(17\)30083-0.pdf](https://www.cell.com/current-biology/pdf/S0960-9822(17)30083-0.pdf)

This one is an especially good reference for understanding mechanoreception.

stimulus



signal

Mechanoreceptors are found in the skin.

General anatomy of skin

This is extra background information not covered directly in the in-class lecture.

General anatomy of skin

This is extra background information not covered directly in the in-class lecture.

- Two major layers:
 - Dermis: most external

epidermis

dermis

General anatomy of skin

This is extra background information not covered directly in the in-class lecture.

- Two major layers:
 - Dermis: most external
 - Epidermis

epidermis

dermis

General anatomy of skin

This is extra background information not covered directly in the in-class lecture.

- Two major layers:
 - Dermis: most external
 - Epidermis
 - Composed of three sublayers, which you don't need to know

epidermis

dermis


Types of skin

Types of skin

- Mammals have two types of skin:


Types of skin

- Mammals have two types of skin:
 - **Glabrous** (nonhairy): lips, palms, feet, parts of genitals




Types of skin

- Mammals have two types of skin:
 - **Glabrous** (nonhairy): lips, palms, feet, parts of genitals
 - **Hairy**: basically everywhere else



Types of skin


- Mammals have two types of skin:
 - **Glabrous** (nonhairy): lips, palms, feet, parts of genitals
 - **Hairy**: basically everywhere else
- These types of skin have distinct types of mechanoreceptor cells, as you will see.



Mechanoreceptors of glabrous skin


Mechanoreceptors of glabrous skin

- Glabrous skin has four major mechanoreceptors:




Mechanoreceptors of glabrous skin

- Glabrous skin has four major mechanoreceptors:
 - **Pacinian corpuscles:** located in the dermis.
 - These are **phasic** receptors that sense **vibrations**, as deformation of the dermis causes their membranes to leak sodium ions and send action potentials.




Mechanoreceptors of glabrous skin

- Glabrous skin has four major mechanoreceptors:
 - Pacinian corpuscles: phasic receptors that sense vibrations.
 - Ruffini endings: located in the deep dermis (also in hairy skin).
 - These are tonic receptors that respond to skin stretching/deformation.




Mechanoreceptors of glabrous skin

- Glabrous skin has four major mechanoreceptors:
 - Pacinian corpuscles: **phasic** receptors that sense **vibrations**.
 - Ruffini endings: **tonic** receptors that respond to **skin stretch/deformation**.
 - **Meissner corpuscles**: found high in the dermis (also in hairy skin).
 - **Phasic** receptors that detect **movement across the skin** (described as “slip” or “flutter” movements)




Mechanoreceptors of glabrous skin

- Glabrous skin has four major mechanoreceptors:
 - Pacinian corpuscles: **phasic** receptors that sense **vibrations**.
 - Ruffini endings: **tonic** receptors that respond to **skin stretch/deformation**.
 - Meissner corpuscles: **phasic** receptors that detect **movement across skin**
 - **Merkel's discs**: in the basal layer of the epidermis (also in hairy skin). **Tonic** receptors that sense **indentation at a very fine scale**.



Mechanoreceptors of glabrous skin


- Glabrous skin has four major mechanoreceptors:
 - **Pacinian corpuscles:** **phasic** receptors that sense **vibrations**.
 - **Ruffini endings:** **tonic** receptors that respond to **skin stretch/deformation**.
 - **Meissner corpuscles:** **phasic** receptors that detect **movement across skin**
 - **Merkel's discs:** **tonic** receptors that sense **indentation at a very fine scale**.



Mechanoreceptors of hairy skin

Mechanoreceptors of hairy skin

- There is a specialized group of mechanoreceptors associated with hair follicles that you need to know.



Mechanoreceptors of hairy skin


- There is a specialized group of mechanoreceptors associated with hair follicles that you need to know.
- **Longitudinal lanceolate endings:** tonic receptors that surround the follicle and detect **when the hair bends**. Different combinations give rise to the different sensory properties of different hairs.



Free nerve endings


Free nerve endings

- These are found in both types of skin.




Free nerve endings

- These are found in both types of skin.
- They detect “**noxious touch**” which mostly means **pain**.



Free nerve endings

- These are found in both types of skin.
- They detect “**noxious touch**” which mostly means **pain**.
- They are **tonic**.



Mechanoreceptors you need to memorize:

End organ	Skin type	Stimulus	Type of response
Pacinian corpuscle	Glabrous (nonhairy)	High-frequency vibration	Phasic
Ruffini ending	Both	Skin stretching	Tonic
Meissner corpuscle	Both	Movement across the skin	Phasic
Merkel's disc	Both	Indentation	Tonic
Longitudinal lanceolate endings	Hairy	Hair follicle deflection	Tonic
Free nerve endings	Both	"Noxious mechanical" aka pain	Tonic

Mechanoreceptors you need to memorize:

End organ	Skin type	Stimulus	Type of response
Pacinian corpuscle	Glabrous (nonhairy)		
R			
M			
M			
Long lanc			
Free		"Noxious mechanical" aka pain	Tonic

According to Dr. Selvaraj, you 100% need to memorize this information, so make flashcards and start practicing with them.

You **DO NOT** need to know the physiological subtypes or the associated fibers listed in the lecture slides.

Practice questions:

You are holding a vibrating sander for a while as you refinish your floor. After a while, you realize that you are no longer sensing the vibrations in your hand. Why is this?

- a. The vibrations damaged mechanoreceptors in your hand.
- b. The vibrations are too small for the receptive fields of your Pacinian corpuscles.
- c. Pacinian corpuscles are phasic receptors.
- d. The stimulus is above the saturation point for your Pacinian corpuscles.

Practice questions:

You are holding a vibrating sander for a while as you refinish your floor. After a while, you realize that you are no longer sensing the vibrations in your hand. Why is this?

- a. The vibrations damaged mechanoreceptors in your hand.
- b. The vibrations are too small for the receptive fields of your Pacinian corpuscles.
- c. **Pacinian corpuscles are phasic receptors.**
- d. The stimulus is above the saturation point for your Pacinian corpuscles.

Practice questions:

You are holding a vibrating sander for a while as you refinish your floor. After a while, you realize that you are no longer sensing the vibrations in your hand. Why is this?

- a. The vibrations damaged mechanoreceptors in your hand.
 - i. This is highly unlikely, and would require very very strong vibrations for a really long time. Plus, you would feel pain if this happened.
- b. The vibrations are too small for the receptive fields of your Pacinian corpuscles.
 - i. There is no such thing as a stimulus that is spatially too small for a receptive field; you would simply sense it as a single stimulus.
- c. **Pacinian corpuscles are phasic receptors.**
- d. The stimulus is above the saturation point for your Pacinian corpuscles.
 - i. You would still feel the vibrations if this was the case; you simply would not be able to detect if they were getting more intense.

Practice questions:

Cats are able to take in lots of information about the world around them via their whiskers. Why would this be the case?

- a. Whiskers are hair follicles with lots of longitudinal lanceolate endings.
- b. Whiskers are in glabrous skin that has many free nerve endings.
- c. The mechanoreceptors surrounding cat whiskers are very phasic.
- d. Whiskers are hair follicles with lots of Pacinian corpuscles.

Practice questions:

Cats are able to take in lots of information about the world around them via their whiskers. Why would this be the case?

- a. Whiskers are hair follicles with lots of longitudinal lanceolate endings.
- b. Whiskers are in glabrous skin that has many free nerve endings.
- c. The mechanoreceptors surrounding cat whiskers are very phasic.
- d. Whiskers are hair follicles with lots of Pacinian corpuscles.

Practice questions:

Cats are able to take in lots of information about the world around them via their whiskers. Why would this be the case?

- a. Whiskers are hair follicles with lots of longitudinal lanceolate endings.
- b. Whiskers are in glabrous skin that has many free nerve endings.
 - i. Free nerve endings detect pain, which is generally not a stimulus used to construct a map of the surrounding world.
- c. The mechanoreceptors surrounding cat whiskers are very phasic.
 - i. Phasic receptors adapt rapidly and stop transmitting signals, which would not be useful for building a map of the world
- d. Whiskers are hair follicles with lots of Pacinian corpuscles.
 - i. Pacinian corpuscles are not found in hairy skin.