

treecm: an introduction

Marco Bascietto

April 5, 2012

1 Examples

1.1 Plot centre of mass

We will make use of the data set bundled in the package to plot a basic view of masses of branches and logs of a stone pine sampled by the author:

```
> library(treecm)
> data(treeData)
> print(treeData)
```

```
$fieldData
      azimuth dBase dTip length tipD height tilt toBePruned biomass
L1         275    73   41   10.2  2.50   0.00   80      FALSE 1740.88372
L2         275    41   16    3.9  2.75  10.20   80      FALSE  183.92713
B1         190    15    0    NA  7.95  10.10    0      FALSE  119.69839
B2         200    22    0    NA  7.95  10.40    0      FALSE  246.69214
B3         230    15    0    NA  7.95  10.40    0      FALSE  119.69839
B4         200    18    0    NA  7.95  11.15    0      FALSE  168.88783
B5         180     7    0    NA  7.95  11.30    0      FALSE   28.38618
B6         150     6    0    NA  7.95  11.30    0      FALSE   21.21769
B7         340    16    0    NA  3.95  11.30    0      FALSE  135.21104
B8         220    13    0    NA  7.95  11.80    0      FALSE   91.35675
B9         165    19    0    NA  7.95  11.80    0      FALSE  187.04037
B10        280     8    0    NA  3.95  11.90    0      FALSE   36.52644
B11        170     9    0    NA  7.95  11.90    0      FALSE   45.62402
B12        265     8    0    NA  7.95  12.20    0      FALSE   36.52644
B13         75     6    0    NA  3.95  12.20    0      FALSE   21.21769
B14        180     6    0    NA  7.95  12.20    0      FALSE   21.21769
B15        170     6    0    NA  7.95  12.60    0      FALSE   21.21769
B16        120     5    0    NA  7.95  12.60    0      FALSE   15.03793
B17         10    14    0    NA  3.95  13.00    0      FALSE  105.07799
B18        180    13    0    NA  7.95  13.00    0      FALSE   91.35675
B19        260    13    0    NA  7.95  13.20    0      FALSE   91.35675
B20         75     6    0    NA  3.95  13.20    0      FALSE   21.21769
```

B21	75	10	0	NA	3.95	13.75	0	FALSE	55.66636
B22	215	7	0	NA	7.95	13.75	0	FALSE	28.38618
B23	140	7	0	NA	7.95	13.75	0	FALSE	28.38618
C	275	16	0	3.0	3.00	14.10	80	FALSE	135.21104

```
$density
```

```
[1] 620
```

```
$allometryFUN
```

```
function (x, diameter)
```

```
{
```

```
  a <- 0.7201
```

```
  b <- 1.8882
```

```
  powerEquation(a, b, as.real(x[diameter]))
```

```
}
```

```
$branchesCM
```

```
[1] 1
```

This data set has been collected for a 17.1 metres tall stone pine whose stem was tilted approx. 20° from the vertical plane (or 80° from the horizontal plane). The stem has been sectioned in two logs (L1 and L2), and a final branch (C). The crown was made up of 23 branches (B1-B23), all of them horizontal (ie tilted 0°). The package recognizes rows as branches because their diameter at tip is 0.

Please notice that some rules have to be followed in order to record sound data in the field:

- the diameter of the tip of L1 is equal to the diameter of the base of L2. L2 tip diameter is, in turn, equal to C base diameter. Height figures must match as well as diameter measures
- the distance of the tip of the branch (`tipD`) is not the length of the branch but the distance between tree base (the origin of the cartesian plot) and branch tip
- note that only the `length` of C branch has been recorded as it is the only branch not being horizontal. Non horizontal branches affect tree CM z-coordinate. When non-horizontal branches are present, and if one is interested in the z-coordinate of CM, than one should record branch length and its angle from the horizontal plane (`tilt`). Otherwise branch `length` is not needed.

Let's get going and compute the centre of mass of this pine:

```
> vectors <- treeVectors(treeData)
> CM      <- centreOfMass(vectors)
> summary(CM)
```

Coordinates of the centre of mass:


Cartesian (x/m, y/m, z/m): -2.09 , -1.98 , 7.70

Polar (angle/degrees, distance/m, height/m): 226 , 2.88 , 7.70

The core of the package is the `summary` method for CM object. The centre of mass for this stone pine lies 2.88 metres South-West of tree base (226° from magnetic North), 7.70 metres above ground. Cartesian coordinates are provided as well, though not so usefull as polar ones.

A simple visualization of tree centre of mass and its logs and branches is achieved simply by:

```
> plot(vectors, main = "A stone pine centre of mass")
> plot(CM)
```




In a cartesian coordinate system whose origin lies at tree base, the masses of logs branches and branches are plotted as vectors pointing inwards, towards the ground. Each circle represent a branch or log mass. Circle radius is proportional to branch or log mass. Likewise, the centre of mass is plotted as a vector pointing inwards, in red colour. Its height component is written alongside its label as z coordinate. A red arrow exemplifies the direction the tree will follow should it break from its base.

If we measured the polar coordinates of the building(s) around the tree we would have plot them alongside the tree using the `plotPolarSegment` function:

```

> plot(vectors,
+   main = "A stone pine centre of mass",
+   xlim = c(-8, 10),
+   ylim = c(-12, 4)
+ )
> plot(CM)
> plotPolarSegment(210, 10.6, 140, 14.4)

```



1.2 Snow load

Snow may increase crown load substantially, sometimes breaking entire branches. As a side effect, snow-loaded crowns may alter tree centre of mass by moving it upwards and, in asymmetric crowns, towards the part of crown under heavier load.

Let's model a snow load that doubles the biomass of branches higher than 12 m:

```

> rows <- substr(row.names(treeData$fieldData), 1, 1)
> q1 <- subset(treeData$fieldData, subset=(height < 12 | rows == "L"))
> q2 <- subset(treeData$fieldData, subset=(height > 12 & rows != "L"))
> q2$biomass <- q2$biomass * 2
> treeData$fieldData <- rbind(q1, q2)
> rm(list = c("q1", "q2", "rows"))


```

Let's recalculate the vectors under snow load and plot the results:

```
> vectors <- treeVectors(treeData)
> CM      <- centreOfMass(vectors)
> summary(CM)

Coordinates of the centre of mass:
Cartesian (x/m, y/m, z/m): -1.96 , -1.92 , 8.62
Polar (angle/degrees, distance/m, height/m): 225 , 2.74 , 8.62

> plot(vectors,
+   main = "A stone pine centre of mass under 2x snow load",
+   xlim = c(-8,10),
+   ylim = c(-12,4)
+ )
> plot(CM)
> plotPolarSegment(210, 10.6, 140, 14.4)
```



Tree centre of mass has clearly shifted upwards and towards the house...

1.3 Wind load

Winds may increase load on some sectors of the crown and decrease it in other sectors. We would like to model the effect of a prevailing Southbound

wind that halves branches mass in the northern sector and doubles it in the southern sector.

```
> data(treeData)
> rows <- substr(row.names(treeData$fieldData), 1, 1)
> treeData$fieldData <- within(
+   treeData$fieldData, {
+     biomass[((azimuth >= 270 | azimuth < 90) & rows != "L")] <- biomass[((azimuth >= 270 | a
+     biomass[((azimuth >= 90 | azimuth < 270) & rows != "L")] <- biomass[((azimuth >= 90 | az
+   })
> rm(rows)
> vectors <- treeVectors(treeData)
> CM <- centreOfMass(vectors)
> summary(CM)
```

Coordinates of the centre of mass:

Cartesian (x/m, y/m, z/m): -2.08 , -3.20 , 8.69

Polar (angle/degrees, distance/m, height/m): 213 , 3.82 , 8.69

Under a heavy Southbound wind the CM of the tree will move considerably towards South and 1.2 metres farther away from tree base. Although too simplistic a model the results lead to the conclusion that dynamic forces in prevailing wind conditions should be taken into account when assessing tree stability.

1.4 Effect of pruning


As far as static forces are concerned, in an effort to move centre of mass toward tree base, we could prune a few heavy branches. Let's have a look how CM would move if we cut B2 and B4. We will make use of function `switchBranchPruningStatus`:

```
> library(treecm)
> data(treeData)
> vectors <- treeVectors(treeData)
> CM <- centreOfMass(vectors)
> op <- par(mfrow = c(2, 1), mai = c(0.5,0.5,0.5,0.2))
> plot(vectors,
+   main = "Centre of mass of a stone pine"
+ )
> plot(CM)
> treeData <- switchBranchPruningStatus(treeData, c(4, 6))
> vectors <- treeVectors(treeData)
> CM <- centreOfMass(vectors)
> plot(vectors,
+   main = "Centre of mass of a stone pine, B2 and B4 removed"
+ )
```

```

> plot(CM)
> par(op)
> rm(op)

```



CM has actually moved 0.5 metres towards tree base, and farther away from the house. As a matter of facts, branch pruning has not been a reasonable action towards a safer tree.

1.5 Slenderness ratio

The slenderness ratio of a tree is defined as $SR = \frac{h}{d}$ where h is the height of the tree trunk, and d is the diameter of the tree Mattheck et al. (1995). The SR is a measure of tree stability and is extensively used in tree stability measures carried out by Visual Tree Assessment (VTA). SR in the range $30 \leq SR \leq 70$ are considered optimal, whereas $SR > 70$ lead to consider the tree at risk of breaking due to its excessive slenderness. The authors have applied the same concept to tree branches as well. While SR in vertical trees has a physical meaning Mattheck et al. (1995), branches are not usually vertical. As the branch starts to deviate from the verticality (as most of the branches do) the arm of the moment gets longer, reaching a maximum limit in horizontal branches. The longer the arm, the higher the stress on the branch. In order to estimate the added stress imposed by branch angle we improved Mattheck's formula by adding a component proportional to branch tilt angle:


$$SR_c = \frac{l}{d} \cdot (1 + \cos\alpha)$$

where α is branch tilt angle (i.e. 90° for a vertical branch, 0° for an horizontal branch). In vertical branches $SR_c = SR$, in horizontal branches $SR_c = 2SR$. As far as we know this is the first attempt to apply the slenderness ratio to branches. Optimal (safe) branches could be in the range $30 \leq SR_c \leq 70$.

When **treeData** object is filled with branches length (not the length of the projection on the ground, from the tree base to their tip) than SR_c can be computed and plotted:

```
> data(treeData)
> # assign length to branches
> treeData$fieldData$length <- c(10.2, 3.9, 7, 7, 7, 7, 7, 7, 3.95, 7, 7, 3.95, 7, 7, 3.95,
> vectors <- treeVectors(treeData)
> SR <- treeSR(treeData,vectors)
> plot(SR, main = "Branches slenderness ratio", xaxt='n', yaxt = 'n', xlab = "", ylab = "")
```

Branches slenderness ratio



The 2D plot charts branches azimuth as arrows whose length is SR_c . The longer the arrows the more slender the branch. Arrows pointing inside the red circle are considered to be stable, whereas longer arrows are considered as risky ($SR_c \geq 70$). The plot may be a visual clue on the process of branch pruning selection.

2 The problem

Estimating the coordinates of the centre of mass (barycentre) of a tree is crucial to judge its static stability. The centre of mass of a perfectly static tree lies inside its circumference at its base, that is the x and y coordinates of the barycentre lie inside the $\pi \cdot r^2$ surface where r is the radius of tree base.

The more distant the centre of mass from tree base the higher the constraints the tree poses on the soil through its roots. When concerns about tree stability are raised and the tree needs to be consolidated a proper cabling system has to be put in place. Knowing in advance the direction the tree would fall in case of breakage at its base is necessary to properly engineer the cabling system.

Nevertheless, the estimate of barycentre position is useful to assess the effects of different pruning SRhemes as far as tree balance is concerned.

3 Data collection

Data collection to estimation of the centre of mass is carried out in three steps:

1. Field measurements
2. Visual check for correctness of assumptions
3. Collection of correct allometric equation in order to estimate branch and foliage biomass

3.1 Field measurements

A few field measurements are needed to estimate centre of mass position at the stem level and at the branch level. Field data are easily taken climbing the tree using tree climbing techniques or by hydraulic platforms. A few instruments are needed including:

- A forestry caliper to measure diameter of logs and branches
- A clinometer, or ipsometer or any other instrument to measure height of branches or logs
- A measuring tape to measure length of branch or log projections on the ground

The stem is ideally sectioned in logs in order to compute their volume and biomass. The measurements to be taken on each log include:

- Diameter at the base of the log, in cm
- Diameter at the top of the log, in cm
- Length of the log, in m

- Azimuth of the log, in case it is not vertical, in degrees from North (0° North, 180° South)
- Length of log projection on the ground, from the tree base to the log tip (0 for a vertical stem), in m
- Height above ground of the base of the log, in m
- Log tilt from the horizontal plane (eg a vertical log is tilted by 90° , an horizontal log is tilted by 0°), in degrees (optional)

Each branch contributes to the position of the centre of mass by means of their wooden component and their foliage component. Every part of a tree carrying foliage is considered to as a branch. This definition applies to tree tip as well, although some trees may have lost their tip or have it removed during topping operations. The measurements to be taken on each branch include:

- Diameter at the base of the branch, in cm
- Azimuth of the branch, in degrees, usually measured with a compass (0° North, 180° South)
- Length of branch projection on the ground, from the tree base to the branch tip, in m
- Height above ground of the branch insertion into the stem
- Branch tilt from the horizontal plane (eg a vertical branch is tilted by 90° , an horizontal branch is tilted by 0°), in degrees (optional)

3.2 Visual check for correctness of assumptions

3.2.1 Relative position of centre of masses of branches and logs

The position of the centre of mass of a tree is computed taking into account the centre of mass of each branch and log. Pinpointing the centre of mass along a branch, taking into account branch form factor and the pattern of distribution of leaves biomass along it, would require many more field measures raising the time spent on it and the costs of the sampling.

Since the package aims to help engineering a consolidation system, the centre of mass is by default located at branches or logs tip. This leads to an estimate of the coordinated of tree centre of mass that is further away from the base than the real one. This difference can be regarded as an inherent safety factor.

The package behaviour can be modified in order to let the position branches and logs centre of mass to get nearer to their base. The relative position of the centre of mass of branches and logs can be set as a real number ranging from 0.01 (base) to 1 (tip, the default behaviour). Setting can be done during import of field data using function `importField` and its parameter `bCM` or by using the setter function `setBranchesSRM`.

3.2.2 The density

Log mass is estimated by converting its volume (as measured in the field) to fresh weight. The conversion factor is usually referred to as density. Wood density is usually quite conservative among individual of the same tree species. Density values are commonly found in published literature. The following table (Niklas (K. J. et al.)) can be a useful resource (density in $\frac{kg}{m^3}$, measured at 50% moisture content):

```
> library(treecm)
> data(Dst)
> print(Dst)
```

	species	group	density
1	Abies alba	conifer	545
2	Abies alba	conifer	577
3	Abies balsama	conifer	529
4	Abies grandis	conifer	449
5	Abies procera	conifer	465
6	Agathis vitiensis	conifer	673
7	Araucaria angustifolia	conifer	689
8	Chaemaecyparis lawsoniana	conifer	497
9	Larix decidua	conifer	673
10	Larix eurolepis	conifer	577
11	Larix kaempferi	conifer	609
12	Picea abies	conifer	497
13	Picea alba	conifer	529
14	Picea omorika	conifer	497
15	Picea sitchensis	conifer	481
16	Picea sitchensis	conifer	529
17	Pinus caribaea	conifer	977
18	Pinus contorta	conifer	593
19	Pinus holfordiana	conifer	513
20	Pinus nigra	conifer	609
21	Pinus nigra	conifer	705
22	Pinus pinaster	conifer	609
23	Pinus ponderosa	conifer	561
24	Pinus radiata	conifer	577
25	Pinus radiata	conifer	641
26	Pinus strobus	conifer	433
27	Pinus sylvestris	conifer	625
28	Podocarpus sp.	conifer	641
29	Podocarpus guatemalensis	conifer	657
30	Pseudotsuga menziesii	conifer	625
31	Pseudotsuga menziesii	conifer	673
32	Thuja heterophylla	conifer	545
33	Thuja heterophylla	conifer	593

34	Thuja heterophylla	conifer	609
35	Thuja plicata	conifer	465
36	Aesculus hippocastanum	dicot	657
37	Acacia mollissima	dicot	897
38	Acer psedudoplatanus	dicot	721
39	Afzelia quanzensis	dicot	1137
40	Alnus glutinosa	dicot	675
41	Alstonia boonei	dicot	497
42	Anthocephalus chinensis	dicot	567
43	Aspidosperma sp.	dicot	993
44	Autranella congolensis	dicot	1144
45	Berlinia confusa	dicot	849
46	Betula sp.	dicot	801
47	Brachstegia nigerica	dicot	865
48	Brachylaena hutchinsii	dicot	1153
49	Byrsonima coriacea	dicot	865
50	Calophyllum brasiliense	dicot	817
51	Canarium schweinfurthii	dicot	593
52	Carpinus betulus	dicot	865
53	Cassispourea malasana	dicot	897
54	Castanea sativa	dicot	657
55	Cedrela odorata	dicot	433
56	Celtis sp.	dicot	961
57	Ceratopetalum apetalum	dicot	733
58	Chlorophora excelsa	dicot	817
59	Cordia millenii	dicot	545
60	Cullenia ceylanica	dicot	769
61	Cyanomeria alexandri	dicot	1121
62	Cylicodiscus gabunensis	dicot	1185
63	Dipterocarpus sp.	dicot	929
64	Dipterocarpus acutangulus	dicot	913
65	Dipterocarpus caudiferus	dicot	753
66	Dipterocarpus zeylanicus	dicot	977
67	Drychalanops beccarii	dicot	865
68	Drychalanops keithii	dicot	902
69	Drychalanops lanceolata	dicot	865
70	Entandrophragma angolense	dicot	705
71	Entandrophragma cylindricum	dicot	833
72	Entandrophragma utile	dicot	833
73	Eperua sp.	dicot	1169
74	Erythrophteum sp.	dicot	1362
75	Eucalyptus pilularis	dicot	897
76	Eucalyptus marginata	dicot	1009
77	Eucalyptus microcorys	dicot	1234
78	Eucalyptus paniculata	dicot	1346
79	Eucalyptus versicolor	dicot	1041

80	<i>Eusideroxylon zwageri</i>	dicot	1282
81	<i>Fagus sylvatica</i>	dicot	833
82	<i>Fraxinus excelsior</i>	dicot	801
83	<i>Gmelina arborea</i>	dicot	625
84	<i>Gonystylus macrophyllum</i>	dicot	785
85	<i>Gossweilerodendron balsamiferum</i>	dicot	641
86	<i>Guarca excelsa</i>	dicot	689
87	<i>Guarca thompsonii</i>	dicot	817
88	<i>Heritiera simplicifolia</i>	dicot	753
89	<i>Heritiera simplicifolia</i>	dicot	801
90	<i>Hevea brasiliensis</i>	dicot	865
91	<i>Hopea sengal</i>	dicot	817
92	<i>Khaya anthotheca</i>	dicot	657
93	<i>Khaya grandiflora</i>	dicot	817
94	<i>Khaya ivorensis</i>	dicot	641
95	<i>Khaya nyascia</i>	dicot	705
96	<i>Koordersiodendron pinnatum</i>	dicot	1089
97	<i>Loniciocarpus castillo</i>	dicot	1169
98	<i>Lophira alata</i>	dicot	1292
99	<i>Lovoa trichilioides</i>	dicot	673
100	<i>Maesopsis veminii</i>	dicot	609
101	<i>Mansonia altissima</i>	dicot	801
102	<i>Mora excelsa</i>	dicot	1137
103	<i>Muragne sp.</i>	dicot	689
104	<i>Nauclea diderrichii</i>	dicot	945
105	<i>Nectrandra sp.</i>	dicot	689
106	<i>Newtonia buchaneni</i>	dicot	705
107	<i>Nothofagus sprocera</i>	dicot	561
108	<i>Ocotea rodiaei</i>	dicot	1250
109	<i>Ocotea usambarensis</i>	dicot	769
110	<i>Octomeles sumatrana</i>	dicot	481
111	<i>Olea hochstetteri</i>	dicot	1121
112	<i>Oxystigma oxyphyllum</i>	dicot	801
113	<i>Parashorea sp.</i>	dicot	705
114	<i>Parashorea malaanonan</i>	dicot	641
115	<i>Parashorea tomentelia</i>	dicot	577
116	<i>Peltogyne sp</i>	dicot	1105
117	<i>Pericopsis elata</i>	dicot	977
118	<i>Pipradeniostrum africanum</i>	dicot	849
119	<i>Platanus hybrida</i>	dicot	785
120	<i>Populus canadensis</i>	dicot	529
121	<i>Populus x canescens</i>	dicot	577
122	<i>Populus x canescens</i>	dicot	481
123	<i>Protium decendrum</i>	dicot	801
124	<i>Prunus avium</i>	dicot	753
125	<i>Pseudosindora palustris</i>	dicot	833

126	<i>Pterocarpus angolensis</i>	dicot	881
127	<i>Pterygota bequaertii</i>	dicot	849
128	<i>Pterygota macrocarpa</i>	dicot	705
129	<i>Qualea</i> sp.	dicot	897
130	<i>Quercus</i> sp.	dicot	833
131	<i>Quercus cerris</i>	dicot	929
132	<i>Quercus rubra</i>	dicot	865
133	<i>Ricinodendron rautanenii</i>	dicot	224
134	<i>Salix</i> x <i>alba</i>	dicot	529
135	<i>Salix alba</i> var. <i>coerulea</i>	dicot	513
136	<i>Salix fragilis</i>	dicot	529
137	<i>Sclerocarpa</i> sp.	dicot	657
138	<i>Scottellia coriacea</i>	dicot	849
139	<i>Shorea acuminatissima</i>	dicot	609
140	<i>Shorea faguetiana</i>	dicot	673
141	<i>Shorea gibbosa</i>	dicot	625
142	<i>Shorea guiso</i>	dicot	993
143	<i>Shorea hakeifolia</i>	dicot	689
144	<i>Shorea leptoclados</i>	dicot	545
145	<i>Shorea macrophylla</i>	dicot	449
146	<i>Shorea parviflora</i>	dicot	513
147	<i>Shorea pauciflora</i>	dicot	689
148	<i>Shorea smithiana</i>	dicot	513
149	<i>Shorea superba</i>	dicot	945
150	<i>Shorea superba</i>	dicot	1057
151	<i>Shorea waltonii</i>	dicot	529
152	<i>Staudtia stipitata</i>	dicot	1139
153	<i>Sterculia oblonga</i>	dicot	913
154	<i>Sterculia rhinopetala</i>	dicot	961
155	<i>Swartzia leiocalycine</i>	dicot	1298
156	<i>Symphonia globulifera</i>	dicot	881
157	<i>Syncarpia glomulifera</i>	dicot	1025
158	<i>Tarrietia utilis</i>	dicot	817
159	<i>Tectona grandis</i>	dicot	801
160	<i>Tectona grandis</i>	dicot	817
161	<i>Terminalia amazonica</i>	dicot	961
162	<i>Tieghemelia heckerii</i>	dicot	801
163	<i>Tilia vulgaris</i>	dicot	657
164	<i>Triplochiton scleroxylon</i>	dicot	465
165	<i>Ulmus glabra</i>	dicot	753
166	<i>Ulmus hollandica</i>	dicot	641
167	<i>Ulmus procera</i>	dicot	641
168	<i>Virola koschnyi</i>	dicot	657
169	<i>Vochvsia</i> sp.	dicot	657
170	<i>Vochysia hondurensis</i>	dicot	577

3.3 Finding a correct allometric equation in order to estimate branch and foliage biomass

It is not feasible to weight the branches of a living tree. As a result branch and foliage biomass is estimated using branch diameter at base. Models relating size or biomass to diameter of trees or branches are known as allometric equations. They usually take the form of $Y = a \cdot X^b$ where Y is branch biomass, X is branch diameter, a and b are parameters estimated on a subsample of branches.

When subsampling is not possible one should rely on published allometric equations and feed them to `treecm`. Currently `treecm` ships with three allometric equations:

- `branchBiomassPine`, tested on stone pine trees (not on branches), 40+ cm diameter, returns biomass (dry weight), Cutini et al. (2009)
- `branchBiomassPinePorte`, tested on maritime pine branches, 10- cm diameter, returns biomass (dry weight), Porté et al. (2002)
- `branchBiomassPineASRa`, tested on stone pine branches, 8-16 cm diameter, predominantly from the lower layers of the crown, returns fresh weight

The proper allometric equation to be used must be fed to `treecm` when importing field data using function `importFieldData`, parameter `branchesAllometryFUN`.

We welcome contributions to increase the available list of allometric equations.

Note that one should pick an allometric equation that yields fresh mass of branches in order to get results as closer as possible to the real tree centre of mass. Function `branchBiomassPineASRa` returns fresh weight values for stone pine branches.

4 Correct layout of CSV file

A sample CSV data file is provided in the `data` directory. Function `importFieldData` loads and stores CSV files and along with needed data. CSV files are made up of 9 columns. The first row has to hold column headers. Headers are case sensitive. Each row holds individual log or branch data. Headers include:

1. **code** a simple code assigned to each log or branch (usually Lx for logs, Bx for branches)
2. **azimuth** orientation, ie: compass bearing in degrees
3. **dBase** diameter of log or branch basal section, in cm
4. **dTip** diameter of log or branch tip (always 0 for branches), in cm
5. **length** log length (leave it empty in case of branches), in m
6. **tipD** distance of the tip of the log or branch to tree base (different from branch length when tree stem is not vertical)

7. **height** height of log basal section of height of branch insertion on stem
8. **tilt** log or branch tilt from the horizontal plane (eg a vertical branch is tilted by 90°, an horizontal branch is tilted by 0°), in degrees (optional, only useful to estimate z coordinate of centre of mass)
9. **toBePruned** boolean to simulate branch pruning

There are a few simple rules to be followed in order to layout a correct CSV file, please refer to page 2

5 Contribute!

treecm is an ongoing project hosted on GitHub (<https://github.com/mbask/treecm>). Many areas need to be expanded including:

- branch biomass estimation; allometric equations are used to estimate fresh branch and foliage biomass. So far only branch biomass for stone pine and maritime pine have been developed or integrated into the software from published data. We need to expand further the number of species represented, particularly for those species common in urban area such as cedars, magnolias, oaks
- The package does not estimate the position of the centre of mass of tree branches. This position may vary according to foliage mass and its distribution along the branches, branch tapering, quantity of water in leaves (ie: shaded or lit leaves) *etc.* The position must be fed to the package during data loading, as the variable **branchesSRM**. Although going for the safe road, setting a branch centre of mass position on its tip may not be sufficiently precise should one assess wood quality as a function of load balance. Work is under way in order to model branch load balance
- As far as position of centre of mass in logs, the package does not tell branches and logs apart. The position of CM in logs follows branches CM position settings, though not realistic.

References

- Cutini, A. and Hajny, M. and Gugliotta, O. and Manetti, M. and Amorini, E. 2009, Effetti della struttura del popolamento sui modelli di stima del volume e della biomassa epigea (Pineta di Castelfusano - Roma) *Forest@*, **6**, 75–84
- Niklas, K. J. and Spatz, H.-C., 2010, Worldwide correlations of mechanical properties and green wood density *American Journal of Botany*, **97**, 1587–1594

- Porté, A. and Trichet, P. and Bert, D. and Loustau, D. 2002, Allometric relationships for branch and tree woody biomass of Maritime pine (*Pinus pinaster* Ait.) *Forest Ecology and Management*, **158**, 71–83
- Mattheck, C. and Breloer, H., 1995, *The Body Language of Trees: A Handbook for Failure Analysis (Research for Amenity Trees)*. HMSO (London).