

EXERCISES AND EXACT SOLUTIONS

Resistance Signals Require Stewardship, Not Panic

Independent practice, facilitated discussion, and assessment reference

Production Unit	PU-B07-C27
Notebook	V7_B_N07
Version	0.1
Assessment boundary	Learning evidence—not operational authorization

Instructions

Complete Part A before consulting the solutions. State assumptions explicitly. Where evidence is insufficient, write an abstention or escalation response rather than inventing precision.

Exercise 1

Deduplicate isolates.

- Required output: a concise decision-ready response with owner, evidence, uncertainty, and boundary where applicable.

Exercise 2

Handle breakpoint changes.

- Required output: a concise decision-ready response with owner, evidence, uncertainty, and boundary where applicable.

Exercise 3

Explain why resistance rates are not prescribing rules.

- Required output: a concise decision-ready response with owner, evidence, uncertainty, and boundary where applicable.

Exact solutions and scoring guide

Solution 1

Use the agreed first-isolate-per-patient/episode/specimen rule.

Solution 2

Reinterpret stored measurements where valid or stratify by breakpoint version; do not splice incomparable rates.

Solution 3

Treatment also depends on syndrome, specimen, patient factors, severity, local guidance, toxicity, availability, and clinician judgment.

Scoring rubric

Criterion	Strong evidence	Revision required
Decision framing	Owner, action, horizon, authority named	Model output substituted for decision
Evidence	Concepts, grain, provenance, limits explicit	Data treated as self-explanatory
Uncertainty	Abstention/escalation used appropriately	False precision or hidden limitations
Governance	Human authority and safeguards retained	Authority silently delegated