

Project planning

New real-time GNSS algorithms for detection and measurement of potential
geoeffective stellar flares

Author

David Moreno Borràs

Supervisor

Manuel Hernández-Pajares

Specialization

Computing

March 27, 2019

Abstract

Solar flares are sudden electromagnetic emissions on the Sun's surface that release large amounts of magnetic energy. These flares emit radiation that has an effect on Earth's ionosphere electron content and are detected by telescopes such as Swift or Fermi by performing gamma-ray observations from low Earth orbit. (Swift y fermi no son para solar flares son para GRB en general!!, quizas aqui deberia ir alguno que se centre en el sol especificamente)

Another approach for detecting these events is possible: the ionosphere electron content variation can be studied using data from the Global Navigation Satellite System (GNSS), specifically the Global Positioning System (GPS).....???

Algorithms for detecting solar flares without knowing the location of the source, that is, the position of the Sun relative to Earth, as well as a study on the feasibility of the detection of such events for the challenging scenario of far-away stars, are presented, aiming to find an alternative detection method to that of the telescopes and using free, open-source data.

Keywords: Solar flares, stellar flares, algorithms, GNSS, GPS, IGS, gamma ray bursts?

Contents

1	Background	2
1.0.1	Global Navigation Satellite Systems	2
1.0.2	Ionosphere	2
1.0.3	Stellar flares	3
1.0.4	Gamma-Ray Bursts	3
1.1	Study on the feasibility of stellar flare detection	4
1.1.1	Sources of data and possible candidates	4
1.1.2	The Neil Gehrels Swift Observatory and its data	5
1.1.3	Objective function	6
1.1.4	Obtaining the data	7
1.1.5	Results	7
2	Solar flare detection	8
2.1	Pseudocode	8

Chapter 1

Background

As the project has a large background in physics and astronomy, some of the relevant topics that are going to be studied are introduced: Global Navigation Satellite Systems, the Ionosphere, Stellar Flares and Gamma-Ray Bursts.

1.0.1 Global Navigation Satellite Systems

During the project both GNSS and GPS will be mentioned. GNSS stands for Global Navigation Satellite Systems and GPS for Global Positioning System

In 1998 the International GNSS Service (IGS) was created as a collaboration of several members of the scientific community: Center for Orbit Determination in Europe (CODE), (European Space Agency) ESA, Jet Propulsion Laboratory (JPL) and Polytechnic University of Catalonia (UPC). This voluntary federation has made available open access GNSS data since its creation.

Its data is provided by more than 300 GPS receivers around the globe and is processed by the previous institutions which compute the global distribution of the TEC.

de donde sacamos los datos

Referencias en este apartado: The IGS VTEC maps: a reliable source of ionospheric information since 1998

explicado en pagina 5 del libro este Reference: GNSS – Global Navigation Satellite Systems: GPS, GLONASS, Galileo, and more

1.0.2 Ionosphere

The ionosphere is a layer of the Earth's atmosphere that lies 75-1000km above its surface. (<http://solar-center.stanford.edu/SID/activities/ionosphere.html>)

Due to the free electrons and ionized molecules it has, it is capable of affecting radio wave propagation, thus having an effect on Global Navigation Satellite Systems (GNSS) technology, this phenomena allows these satellites to be used as a global scanner for the ionosphere.

The main physical quantity used for describing the electron content of the ionosphere is the Total Electron Content (TEC), the TEC is the total number of electrons between two

points (r_1, r_2) along a cylinder of base $1m^2$. Slant TEC (STEC), in particular can be defined as the TEC in which r_1 and r_2 are a satellite and a receiver's positions.

(aquí citar a GPS as a solar observational instrument: Real-time estimation of EUV photons flux rate during strong, medium, and weak solar flares)

The unique properties the ionosphere has enable us to use the data provided by the GNSS technology to study a powerful phenomena that occurs in many stars across the universe: stellar flares.

Referencias en este apartado: 1International reference ionosphere 1990, 2The ionosphere: effects, GPS modeling and the benefits for space geodetic techniques

1.0.3 Stellar flares

Flares from stars, in particular those that have the Sun as a source, more noticeable due to its proximity, are sudden flashes of brightness in the surface of stars which release large amounts of energy across the whole electromagnetic spectrum (X-rays and Extreme Ultraviolet (EUV)) rays. Flares that have the Sun as a source can increase the electron content of the ionosphere and have an effect on waves passing through it, affecting satellite communications and causing a delay. This phenomena is the key element of the project, as it enables us to study these events.

Satellites can also be harmed by this effects: the flares heat up the outer atmosphere, which in turn increases the drag on these satellites reducing their lifetime in orbit. GPS as a solar observational instrument: Real-time estimation of EUV photons flux rate during strong, medium, and weak solar flares se explica el efecto del sol sobre ionosfera

The previous phenomena applies to flares originating in the Sun, whether a flare that has a star from outside the Solar System as a source has an effect on the Earth's atmosphere or not is one of the topic that is going to be studied in this project.

1.0.4 Gamma-Ray Bursts

Throughout the project, in particular when studying the feasibility of stellar flares detection, another type of event will be mentioned and studied as well: Gamma-Ray Bursts (GRBs):

GRBs are highly energetic explosions that occur in distant galaxies, releasing large amounts of radiation, in particular Gamma rays, hence the name of the event. These bursts, despite being millions of light years away from Earth are so powerful they might still have an impact on the ionosphere, like the aforementioned stellar flares. The main difference with flares originating from stars is that GRBs are thought to be originated from the death of massive stars, that is, supernovas.

This event, despite not being a stellar flare, the phenomena we aim to detect, is going to be studied in the following sections to test the currently working algorithms mainly for two reasons: it has been studied and cataloged by telescopes such as the Fermi Observatory and there's is available information that we can use for our study, and the large amounts of energy they emit it make GRBs a more feasible target to detect.

Fermi Observations of High-Energy Gamma-Ray Emission from GRB 080916C

1.1 Study on the feasibility of stellar flare detection

Flares from far-away stars and Gamma-ray Bursts, albeit more powerful than flares that have the Sun as their source, may not be possible to detect due to the large distances that separate them from our measurement tool: the ionosphere.

Before starting to adapt the algorithm for detecting solar flares to this scenario, a study was conducted parallel to its development, to see if the energy from flares originating in far-away stars could be detected using the already existing method, namely the GNSS Solar Flare Activity Indicator (GSFLAI) algorithms [3].

To study if this was feasible, the algorithms were run on certain candidates of flares and GRBs to see if they could be detected.

The project supervisor, Manuel Hernández-Pajares, who as mentioned before has previously performed several studies on the subject, provided the GSFLAI algorithm to test the candidates. The GSFLAI algorithm takes into consideration the location of the source (the Sun) to see if there's a relation between an increase in the VTEC of the ionosphere and the solar-zenith angle to determine if this increase is caused by a solar flare. [GNSS measurement of EUV photons flux rate during strong and mid solar flares]

However, its execution may take up to 2 hours, because of this the aim was to:

- Find a database for possible candidates, several online archives with information about previously recorded Gamma Ray Bursts were considered.
- Select an appropriate source of this pool of candidates by writing a quick script that yielded an ordered list of the best candidates based on certain factors, instead of selecting a random source.

1.1.1 Sources of data and possible candidates

The three main databases we considered for the study were:

- The GRB collection website of Jochen Greiner, scientist at the Max-Planck-Institute for extraterrestrial Physics (MPE) [5], which offers a collection of detected GRBs by different telescopes and observatories.
- The Magnetar Outburst Online Catalog (MOOC), developed by the Institute of Space Sciences (CSIC-IEEC, Barcelona) [6]. We also had the pleasure to meet one of the leaders of this project, Nanda Rea, and discuss
- The Neil Gehrels Swift Observatory website and archive by the National Aeronautics and Space Administration (NASA), Goddard Space Flight Center [7] which contains an archive of detected GRBs by the Swift observatory and is constantly updated.

Because of the layout of the website and how the data could be accessed, the option with which we started was the Swift Database, as the data could be visualized in an HTML table and was easily accessible.

1.1.2 The Neil Gehrels Swift Observatory and its data

The Swift Observatory is a NASA mission with international participation, designed to observe GRBs and their afterglows to study topics such as the origins of GRBs or what they can reveal about the early stages of the universe [9]. The observatory is equipped with three main instruments that work with each other to study GRBs [2] [7]:

- The **Burst Alert Telescope (BAT)**, tasked with detecting the GRBs and computing their positions. This triggers the spacecraft to point the other telescopes to the burst so it can be studied in more detail.
- The **X-ray Telescope (XRT)**, used for studying the X-ray radiation and taking images of the bursts which in turn help increase the accuracy of the location estimation.
- The **UV/Optical Telescope (UVOT)**, which serves a similar purpose to the XRT, but studies the ultraviolet band of the spectrum.

For each detected GRB, the data obtained by the different telescopes is given. The parameters that are relevant to our study and determine the fitness of each of the candidates are:

- The **name of the burst**, given by the date it was detected. For example, the GRB named 190220A was detected the 20th of February of 2019.
- The **Universal Time (UT)** of the detection, that is, hh:mm:ss of the day given by the name.
- The fluence detected by the BAT component, in units of keV. that is.
- The **UVOT magnitude**, measured by the UV Telescope.
- The location that triggered the detection, given as **Right Ascension (Ra)** and **Declination (Dec)**.

Right Ascension and Declination are two concepts similar to longitude and latitude, respectively, used to describe the location of objects in the sky, in particular in a sphere of infinite radius that with the Earth as its center called the **celestial sphere**.

Taking this into account, Right Ascension is the equivalent of longitude, expressed in degrees (or more commonly in hours, minutes and seconds) and Declination, the equivalent of latitude, is expressed in degrees between the two poles: $+90^\circ$ and -90° . [10]

This reference system is used to describe the position of objects in the sky, and it is the one used by the Swift telescope to specify the location of the sources. Afterwards these concepts will play an important role when computing the angle formed between the source and the Sun.

1.1.3 Objective function

Our main goal in this section was to obtain a list of GRB candidates ordered from more to less probable to be detected by the algorithm, that is, their fitness. To obtain this score we had to define an objective function, taking into consideration two factors:

- The **strength** of the burst, given by the UVOT magnitude. If this value was not available (as it was the case with many of the candidates) the BAT fluence was considered as its strength. This values were already given by the archive and no additional computations were required.
- The **angle between the burst and the Sun**, this was an important factor because bursts having an effect on the night hemisphere should be more noticeable than those hitting the day one, where the Sun has a bigger influence.

Computing the angle

As mentioned before, the Swift archive gives us the Right Ascension (Ra) and Declination (Dec) where the source is thought to be located.

The location of the Sun, on the other hand, is unknown. But we do know the time when the burst was detected.

The supervisor, Manuel Hernández-Pajares, provided me an algorithm which takes date (year, month, day and UT) and a planet of the Solar System (or the Sun, our case) as the input and returns its location in the celestial sphere, that is, its Ra and Dec.

This algorithm belongs to the **Starlink Project** (Rutherford Appleton Laboratory), which provided open-source software like the one at hand to astronomical institutions. Although it was shut down in 2005, the code still remains and we could use it for our study [1].

Knowing both the location of the burst and the Sun, the angle between both can be computed and used as a parameter for the objective function.

This computation is done by performing the dot product of the two unit vectors that can be obtained from the Ra and Dec of the objects. These are the unit vectors of

The unit vectors are given by:

```
coordSun = [math.cos(sunDec)*math.cos(sunRa), math.cos(sunDec)*math.sin(sunRa),
math.sin(sunDec)] coordGRB = [math.cos(dec)*math.cos(ra), math.cos(dec)*math.sin(ra),
math.sin(dec)]
```

```
angle = math.acos(coordSun[0]*coordGRB[0] + coordSun[1]*coordGRB[1] + coordSun[2]*coordGRB[2])
angle = angle*180/math.pi
```

$$\sum_{i=0}^{\infty} a_i x^i \quad (1.1)$$

The dot product of the two unit vectors obtained from gave us the angle between the burst and the Sun.

1.1.4 Obtaining the data

Regarding the scrapping of the website to parse the data and obtain this ordered list, **Python** was chosen because the problem required a quick implementation, and Python's libraries offered a great tool to develop a simple solution as quick as possible.

In the script, the website with the table of bursts (see figure x) is scrapped using Python's **BeautifulSoup** library, which has an HTML and XML parser that allows us to easily select and obtain data from a given website.

Insertion sort was used so we could insert every considered GRB into a list of sorted candidates as we were traversing the table.

Regarding the distribution of weight between both factors, strength and angle, we

The best 10 candidates of the resulting sorted table (pie de pagina: bursts registered up to the 26th of February of 2019), is shown here:

We proceeded to study these bursts

1.1.5 Results

Chapter 2

Solar flare detection

Before starting to write the main solar flare detection algorithm a first program was developed in which the location of the sun was known and we knew when the solar flare had taken place.

To test this, the data used was that of the called "Halloween Storm, in whi

2.1 Pseudocode

Algorithm 1 My algorithm

```
1: procedure MYPROCEDURE
2:    $stringlen \leftarrow \text{length of } string$ 
3:    $i \leftarrow patlen$ 
4: top:
5:   if  $i > stringlen$  then return false
6:    $j \leftarrow patlen$ 
7: loop:
8:   if  $string(i) = path(j)$  then
9:      $j \leftarrow j - 1.$ 
10:     $i \leftarrow i - 1.$ 
11:    goto loop.
12:   close;
13:    $i \leftarrow i + \max(delta_1(string(i)), delta_2(j)).$ 
14:   goto top.
```

testing [4] [8] [7]

Bibliography

- [1] (Archived website) The Starlink Project. <https://web.archive.org/web/20120123062045/http://starlink.jach.hawaii.edu/starlink/WelcomePage>. [Online; accessed 10-March-2019].
- [2] N. Gehrels, G. Chincarini, P. Giommi, K. Mason, J. Nousek, A. Wells, N. White, S. Barthelmy, D. Burrows, L. Cominsky, et al. The swift gamma-ray burst mission. *The Astrophysical Journal*, 611(2):1005, 2004.
- [3] M. Hernández-Pajares, A. García-Rigo, J. M. Juan, J. Sanz, E. Monte, and A. Aragón-Ángel. GNSS measurement of EUV photons flux rate during strong and mid solar flares. *Space Weather*, 10(12):1–16, 2012.
- [4] M. Hernández-Pajares, J. Juan, J. Sanz, R. Orus, A. Garcia-Rigo, J. Feltens, A. Komjathy, S. Schaer, and A. Krankowski. The igs vtec maps: a reliable source of ionospheric information since 1998. *Journal of Geodesy*, 83(3-4):263–275, 2009.
- [5] Institute of Space Sciences (CSIC-IEEC. Magnetar Outburst Online Catalog. <http://magnetars.ice.csic.es/#/parameters>. [Online; accessed 3-March-2019].
- [6] Institute of Space Sciences (CSIC-IEEC. Magnetar Outburst Online Catalog. <http://magnetars.ice.csic.es/#/parameters>. [Online; accessed 3-March-2019].
- [7] Jochen Greiner, Max-Planck-Institute for extraterrestrial Physics. GRB collection. <http://www.mpe.mpg.de/~jcgr/bgrgen.html>. [Online; accessed 3-March-2019].
- [8] D. Martínez Cid. First study on the feasibility of stellar flares detection with gps. B.S. thesis, Universitat Politècnica de Catalunya, 2016.
- [9] P. W. Roming, T. E. Kennedy, K. O. Mason, J. A. Nousek, L. Ahr, R. E. Bingham, P. S. Broos, M. J. Carter, B. K. Hancock, H. E. Huckle, et al. The swift ultra-violet/optical telescope. *Space Science Reviews*, 120(3-4):95–142, 2005.
- [10] Solar System Exploration, NASA Science. Reference Systems. <https://solarsystem.nasa.gov/basics/chapter2-2/>. [Online; accessed 5-March-2019].