

# Práctica 0

Álvaro Francisco Ruiz Cornejo

14/9/2021

## Ejercicio 1

```
(dado_honesto = sample(1:6, size = 100, replace = TRUE))
```

```
## [1] 6 6 6 3 1 4 5 6 4 4 2 2 6 6 4 1 1 6 4 4 4 2 3 4 3 3 2 2 5 6 6 5 6 5 3 2 6
## [38] 4 2 1 6 2 5 6 6 3 2 1 6 6 3 5 4 5 6 5 5 3 1 4 2 4 2 2 3 5 5 1 5 4 3 3 5 2
## [75] 1 3 4 4 1 5 6 6 5 3 6 2 6 1 3 1 2 4 5 2 5 4 3 5 4 6
```

```
table(dado_honesto) #frecuencia absoluta R básico
```

```
## dado_honesto
## 1 2 3 4 5 6
## 11 16 15 18 18 22
```

```
signif(prop.table(table(dado_honesto)),2) # frecuencia relativa R básico
```

```
## dado_honesto
## 1 2 3 4 5 6
## 0.11 0.16 0.15 0.18 0.18 0.22
```

```
t = tibble(col_tirada = dado_honesto)
```

```
t %>%
  count(col_tirada) #frecuencia absoluta con dplyr
```

```
## # A tibble: 6 x 2
##   col_tirada     n
##   <int> <int>
## 1         1    11
## 2         2    16
## 3         3    15
## 4         4    18
## 5         5    18
## 6         6    22
```

```
t %>%
  count(col_tirada) %>%
  mutate(col_tirada, relFreq = prop.table(n), n = NULL) #frecuencia relativa con dplyr
```

```
## # A tibble: 6 x 2
##   col_tirada relFreq
##       <int>   <dbl>
## 1         1    0.11
## 2         2    0.16
## 3         3    0.15
## 4         4    0.18
## 5         5    0.18
## 6         6    0.22
```

## Ejercicio 2

Trucamos un dado para duplicar la posibilidad de sacar un seis

```
(dado_cargado = sample(1:6, size = 100, replace = TRUE,
  prob = c((1/7),(1/7), (1/7), (1/7), (1/7),(2/7))))
```

```
## [1] 6 4 5 5 1 6 1 6 5 6 6 6 3 3 3 6 2 6 6 1 5 6 2 6 6 6 6 5 5 6 5 6 1 5 2 3 3
## [38] 6 1 2 1 6 1 2 1 6 6 6 5 4 3 3 6 5 3 2 6 3 5 6 4 4 3 6 6 5 2 4 3 6 5 5 3 5
## [75] 2 4 6 5 3 6 5 4 5 5 4 1 4 2 6 6 2 1 1 6 5 6 6 6 6 1
```

```
table(dado_cargado) #frecuencia absoluta
```

```
## dado_cargado
##  1  2  3  4  5  6
## 12 10 13  9 20 36
```

```
signif(prop.table(table(dado_cargado)),2) # frecuencia relativa
```

```
## dado_cargado
##  1  2  3  4  5  6
## 0.12 0.10 0.13 0.09 0.20 0.36
```

## Ejercicio 3

```
(v1 = rep(seq(from = 1, to = 4, by = 1), each = 4))
```

```
## [1] 1 1 1 1 2 2 2 2 3 3 3 3 4 4 4 4
```

```
(v2 = rep(seq(from = 1, to = 5, by = 1), times = c(1,2,3,4,5)))
```

```
## [1] 1 2 2 3 3 3 4 4 4 4 5 5 5 5
```

```
(v3 = rep(seq(from = 1, to = 4, by = 1), times = 4))
```

```
## [1] 1 2 3 4 1 2 3 4 1 2 3 4 1 2 3 4
```

## Ejercicio 4

```
(mpg2 <- mpg %>%
  filter(class == 'pickup') %>%
  select(starts_with("c")))
```

```
## # A tibble: 33 x 3
##   cyl   cty class
##   <int> <int> <chr>
## 1     6    15 pickup
## 2     6    14 pickup
## 3     6    13 pickup
## 4     6    14 pickup
## 5     8    14 pickup
## 6     8    14 pickup
## 7     8     9 pickup
## 8     8    11 pickup
## 9     8    11 pickup
## 10    8    12 pickup
## # ... with 23 more rows
```

## Ejercicio 5

### 5.1


```
(pob_region = census %>%
  group_by(region) %>%
  summarise(poblacion_total = sum(pop), n = n()))
```

```
## # A tibble: 4 x 3
##   region poblacion_total   n
##   <dbl+lbl>         <dbl> <int>
## 1 1 [NE]           49135283     9
## 2 2 [N Cntrl]     58865670    12
## 3 3 [South]       74734029    16
## 4 4 [West]        43172490    13
```

## 5.2

```
ggplot(pob_region) +  
  geom_col(aes(region, poblacion_total), fill = 'orange',  
            color = 'black')
```

```
## Don't know how to automatically pick scale for object of type haven_labelled/vctrs_vctr/double. Defa
```



## 5.3

```
(orden <- census %>%  
  arrange(desc(pop)))
```

```
## # A tibble: 50 x 12  
##   state      region  pop poplt5 pop5_17 pop18p pop65p popurban medage  death  
##   <chr>    <dbl+lbl> <dbl> <dbl>    <dbl> <dbl>  <dbl>  <dbl>  <dbl>  <dbl>  
## 1 Califor~ 4 [West]  2.37e7 1.71e6 4680558 1.73e7 2.41e6 21607606 29.9 186428  
## 2 New York 1 [NE]    1.76e7 1.14e6 3551938 1.29e7 2.16e6 14858068 31.9 171769  
## 3 Texas    3 [South] 1.42e7 1.17e6 3137045 9.92e6 1.37e6 11333017 28.2 108019  
## 4 Pennsylv~ 1 [NE]    1.19e7 7.47e5 2375838 8.74e6 1.53e6 8220851 32.1 123261  
## 5 Illinois 2 [N Cnt~ 1.14e7 8.42e5 2400796 8.18e6 1.26e6 9518039 29.9 102230
```

```
## 6 Ohio      2 [N Cnt~ 1.08e7 7.87e5 2307170 7.70e6 1.17e6 7918259 29.9 98268
## 7 Florida   3 [South] 9.75e6 5.70e5 1789412 7.39e6 1.69e6 8212385 34.7 104190
## 8 Michigan  2 [N Cnt~ 9.26e6 6.85e5 2066873 6.51e6 9.12e5 6551551 28.8 75102
## 9 New Jer~  1 [NE]    7.36e6 4.63e5 1527572 5.37e6 8.60e5 6557377 32.2 68762
## 10 N. Caro~ 3 [South] 5.88e6 4.04e5 1253659 4.22e6 6.03e5 2822852 29.6 48426
## # ... with 40 more rows, and 2 more variables: marriage <dbl>, divorce <dbl>
```

## 5.4

```
(tasa_matdiv <- census %>%
  select(state, marriage, divorce) %>%
  mutate(tasa = divorce/marriage) %>%
  arrange(desc(tasa))
```

```
## # A tibble: 50 x 4
##   state      marriage divorce  tasa
##   <chr>      <dbl>    <dbl> <dbl>
## 1 Oregon      23004    17762 0.772
## 2 Indiana     57853    40006 0.692
## 3 Florida    108344    71579 0.661
## 4 Arizona     30223    19908 0.659
## 5 Alaska       5361     3517 0.656
## 6 California  210864   133541 0.633
## 7 New Mexico   16641    10426 0.627
## 8 N. Carolina  46718    28050 0.600
## 9 Washington   47728    28642 0.600
## 10 Arkansas    26513    15882 0.599
## # ... with 40 more rows
```

## 5.5

```
census %>%
  summarise(state, medage,
             mediana = median(medage), desvMediana = (medage-mediana),
             propMayorEdad = pop65p/pop) %>%
  arrange(desc(propMayorEdad)) %>%
  head(10)
```

```
## # A tibble: 10 x 5
##   state      medage mediana desvMediana propMayorEdad
##   <chr>      <dbl>    <dbl>    <dbl>    <dbl>
## 1 Florida      34.7     29.8      4.95      0.173
## 2 Arkansas      30.6     29.8      0.850     0.137
## 3 Rhode Island  31.8     29.8      2.05      0.134
## 4 Iowa          30      29.8      0.25      0.133
## 5 Missouri      30.9     29.8      1.15      0.132
## 6 S. Dakota     28.9     29.8     -0.850     0.132
## 7 Nebraska      29.7     29.8     -0.0500    0.131
## 8 Kansas        30.1     29.8      0.350     0.130
## 9 Pennsylvania  32.1     29.8      2.35      0.129
## 10 Massachusetts 31.2     29.8      1.45      0.127
```

## 5.6

```
cortes = seq(min(census$medage), max(census$medage), length.out = 11)
ggplot(data = census, aes(x=medage)) +
  geom_histogram(mapping = aes(y=stat(density)), breaks = cortes,
                fill = "orange", color = "black") +
  geom_density(color = "red", size = 1.5)
```

