The Theoretical Minimum Quantum Mechanics - Solutions

L03E02

M. Bivert

January 4, 2023

Exercise 1. Prove that Eq. 3.16 is the unique solution to Eqs. 3.14 and 3.15.

Let's recall all the equations, 3.14, 3.15 and 3.16

$$\begin{pmatrix} (\sigma_z)_{11} & (\sigma_z)_{12} \\ (\sigma_z)_{21} & (\sigma_z)_{22} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \tag{1}$$

$$\begin{pmatrix} (\sigma_z)_{11} & (\sigma_z)_{12} \\ (\sigma_z)_{21} & (\sigma_z)_{22} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = -\begin{pmatrix} 0 \\ 1 \end{pmatrix}$$
 (2)

$$\begin{pmatrix} (\sigma_z)_{11} & (\sigma_z)_{12} \\ (\sigma_z)_{21} & (\sigma_z)_{22} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\tag{3}$$

By developing the matrix product and identifying the vectors components, the first two equations make a system of four equations involving four unknowns $(\sigma_z)_{11}$, $(\sigma_z)_{12}$, $(\sigma_z)_{21}$ and $(\sigma_z)_{22}$:

$$\begin{cases}
1(\sigma_z)_{11} + 0(\sigma_z)_{12} &= 1 \\
1(\sigma_z)_{21} + 0(\sigma_z)_{22} &= 0 \\
0(\sigma_z)_{11} + 1(\sigma_z)_{12} &= 0 \\
0(\sigma_z)_{21} + 1(\sigma_z)_{22} &= -1
\end{cases}
\Leftrightarrow
\begin{cases}
(\sigma_z)_{11} &= 1 \\
(\sigma_z)_{21} &= 0 \\
(\sigma_z)_{12} &= 0 \\
(\sigma_z)_{22} &= -1
\end{cases}
\Leftrightarrow
\begin{bmatrix}
\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}
\end{bmatrix}$$