

Assignment for subcellular protein localization

1. Name of chosen protein: HWP1
Organism: *Candida albicans*
Locus: CAU64206
NCBI Nucleotide accession #: U64206
NCBI protein accession #: AAC96368
2. Predicted function: HWP1 possibly functions as an adhesin in *Candida albicans*. It has also been demonstrated to play a role in the morphological change (from yeast form to hyphae form) of *Candida albicans*.
3. The adhesin function of HWP1 implies that it localizes to the cell wall. I did not find any data directly proving this though. In addition, I think it would be interesting to check the change of HWP1 localization during the morphological change of *Candida albicans*.
4. Results from different protein localization:

I. TargetP 1.1 Server - prediction results (Technical University of Denmark)

TargetP 1.1 predicts the subcellular location of eukaryotic proteins. The location assignment is based on the predicted presence of any of the N-terminal presequences: chloroplast transit peptide (cTP), mitochondrial targeting peptide (mTP) or secretory pathway signal peptide (SP).

targetp v1.1 prediction results

Number of query sequences: 1

Cleavage site predictions not included.

Using NON-PLANT networks.

Name	Len	mTP	SP	other	Loc	RC

Sequence	634	0.031	0.859	0.098	S	2

cutoff		0.000	0.000	0.000		

DESCRIPTION

Name Sequence name truncated to 20 characters

Len Sequence length

cTP, mTP, SP, Final NN scores on which the final prediction is based (Loc, see below). Note that the scores are not really probabilities, and they do not necessarily add to one. However, the location with the highest score is the most likely according to TargetP, and the relationship between the scores (the reliability class, see below) may be an indication of how certain the prediction is.

Loc	<p>Prediction of localization, based on the scores above; the possible values are:</p> <p>C Chloroplast, i.e. the sequence contains cTP, a chloroplast transit peptide;</p> <p>M Mitochondrion, i.e. the sequence contains mTP, a mitochondrial targeting peptide;</p> <p>S Secretory pathway, i.e. the sequence contains SP, a signal peptide;</p> <p>_ Any other location;</p> <p>* "don't know"; indicates that cutoff restrictions were set (see instructions) and the winning network output score was below the requested cutoff for that category.</p>
RC	<p>Reliability class, from 1 to 5, where 1 indicates the strongest prediction. RC is a measure of the size of the difference ('diff') between the highest (winning) and the second highest output scores. There are 5 reliability classes, defined as follows:</p> <p>1 : $\text{diff} > 0.800$</p> <p>2 : $0.800 > \text{diff} > 0.600$</p> <p>3 : $0.600 > \text{diff} > 0.400$</p> <p>4 : $0.400 > \text{diff} > 0.200$</p> <p>5 : $0.200 > \text{diff}$</p> <p>Thus, the lower the value of RC the safer the prediction.</p>

In a short word, TargetP 1.1 predicts HWP1 will go through the secretory pathway, and it contains a secretory pathway signal peptide (SP).

II. Results of PSORTII

PSG: a new signal peptide prediction method

N-region: length 2; pos.chg 1; neg.chg 0

H-region: length 23; peak value 8.53

PSG score: 4.13

GvH: von Heijne's method for signal seq. recognition

GvH score (threshold: -2.1): -4.50

possible cleavage site: between 27 and 28

>>> Seems to have no N-terminal signal peptide

ALOM: Klein et al's method for TM region allocation

Init position for calculation: 1

Tentative number of TMS(s) for the threshold 0.5: 2

Number of TMS(s) for threshold 0.5: 1

INTEGRAL Likelihood = -3.93 Transmembrane 618 - 634

PERIPHERAL Likelihood = 5.46 (at 398)

ALOM score: -3.93 (number of TMSs: 1)

MITOP: Prediction of membrane topology (Hartmann et al.)

Center position for calculation: 625

Charge difference: 0.0 C(0.0) - N(0.0)

N >= C: **N-terminal side will be inside**>>> **Single TMS is located near the C-terminus**>>> **membrane topology: type Nt (cytoplasmic tail 1 to 617)**

MITDISC: discrimination of mitochondrial targeting seq

R content: 1 Hyd Moment(75): 7.98

Hyd Moment(95): 7.26 G content: 1

D/E content: 1 S/T content: 4

Score: -2.96

Gavel: prediction of cleavage sites for mitochondrial preseq

R-2 motif at 12 MRL|ST

NUCDISC: discrimination of nuclear localization signals

pat4: none

pat7: none

bipartite: none

content of basic residues: 2.1%

NLS Score: -0.47

KDEL: ER retention motif in the C-terminus: none

ER Membrane Retention Signals:

XXRR-like motif in the N-terminus: RLST

none

SKL: peroxisomal targeting signal in the C-terminus: none

SKL2: 2nd peroxisomal targeting signal: none

VAC: possible vacuolar targeting motif: none

RNA-binding motif: none

Actinin-type actin-binding motif:

type 1: none

type 2: none

NMYR: N-myristoylation pattern : none

Prenylation motif: none

memYQRL: transport motif from cell surface to Golgi: none

Tyrosines in the tail: too long tail

Dileucine motif in the tail: none

checking 63 PROSITE DNA binding motifs: none

checking 71 PROSITE ribosomal protein motifs: none

checking 33 PROSITE prokaryotic DNA binding motifs: none

NNCN: Reinhardt's method for Cytoplasmic/Nuclear discrimination

Prediction: nuclear

Reliability: 94.1

COIL: Lupas's algorithm to detect coiled-coil regions

total: 0 residues

Results of the k-NN Prediction

k = 9/23

21.7 %: nuclear

13.0 %: vesicles of secretory system

13.0 %: cytoplasmic

13.0 %: Golgi

13.0 %: mitochondrial

8.7 %: endoplasmic reticulum

8.7 %: plasma membrane

4.3 %: extracellular, including cell wall

4.3 %: cytoskeletal

>> **prediction for QUERY is nuc (k=23)**

III. Results of WOLFPSORT

queryProtein (634 aa)

PSG: a new signal peptide prediction method

N-region: length 2; pos.chg 1; neg.chg 0

H-region: length 23; peak value 3.51

PSG score: -0.89

GvH: von Heijne's method for signal seq. recognition

GvH score (threshold: -2.1): -4.50

possible cleavage site: between 27 and 28

>>> **Seems to have no N-terminal signal peptide**

ALOM: Klein et al's method for TM region allocation

Init position for calculation: 1

Tentative number of TMS(s) for the threshold 0.5: 2

Number of TMS(s) for threshold 0.5: 1

INTEGRAL Likelihood = -3.93 Transmembrane 618 - 634

PERIPHERAL Likelihood = 5.46 (at 398)

ALOM score: -3.93 (number of TMSs: 1)

MTOP: Prediction of membrane topology (Hartmann et al.)

Center position for calculation: 625

Charge difference: 0.0 C(0.0) - N(0.0)

N >= C: N-terminal side will be inside

>>> **Single TMS is located near the C-terminus**

>>> **membrane topology: type Nt (cytoplasmic tail 1 to 617)**

MITDISC: discrimination of mitochondrial targeting seq

R content: 1 Hyd Moment(75): 7.98

Hyd Moment(95): 7.26 G content: 1

D/E content: 1 S/T content: 4

Score: -2.96

Gavel: prediction of cleavage sites for mitochondrial preseq


R-2 motif at 12 MRL|ST

NUCDISC: discrimination of nuclear localization signals

pat4: none
 pat7: none
 bipartite: none
 content of basic residues: 2.1%
 NLS Score: -0.47
 KDEL: ER retention motif in the C-terminus: none
 ER Membrane Retention Signals:
 XXRR-like motif in the N-terminus: RLST none
 SKL: peroxisomal targeting signal in the C-terminus: none
 PTS2: 2nd peroxisomal targeting signal: none
 VAC: possible vacuolar targeting motif: none
 RNA-binding motif: none
 Actinin-type actin-binding motif:
 type 1: none
 type 2: none
 NMYR: N-myristoylation pattern : none
 Farnesylation/Geranylgeranylation motif: none
 memYQRL: transport motif from cell surface to Golgi: none
 Tyrosines in the tail: too long tail
 Dileucine motif in the tail: none
 checking 63 PROSITE DNA binding motifs: none
 checking 71 PROSITE ribosomal protein motifs: none
 checking 33 PROSITE prokaryotic dna binding motifs: none
Final Results:
 48.0 %: extracellular, including cell wall 12.0 %: cytoplasmic 12.0 %: nuclear 12.0 %:
 endoplasmic reticulum 8.0 %: vesicles of secretory system 4.0 %: plasma membrane 4.0 %:
 mitochondrial >> **prediction for queryProtein is exc**

WOLFPSORT is an updated version of PSORTII. Comparing the results from two programs, we can see that they gave same/similar results in most aspects. Both programs said that HWP1 doesn't seem to have an N-terminal signal peptide. HWP1 looks more like a peripheral protein rather than an integral one, with single trans-membrane sequence near its C-terminus and its N-terminus facing inside the cell. Nonetheless, those two programs gave controversial results of HWP1 localization. PSORTII suggests that HWP1 localizes to nucleus, while WOLFPSORT predicts HWP1 localizes to extracellular space or cell wall. Given adhesion function of HWP1, the result of WOLFPSORT makes more sense. And it is consistent with the result of TargetP (Because WOLFPSORT has been updated?) However, we can not rule out the possibility that HWP1 may localize to nucleus.

5. HWP1 were overall predicted to be a peripheral membrane protein by all the programs. I further used iPSORT to do an AAindex Analysis, and got



Based on this Hydropathy Index of HWP1, I would predict a trans-membrane sequence at its N-terminus. (If I have understood the hydropathy index correctly) I don't know why both PSORTII and WOLFPSORT predict a trans-membrane sequence at the C-terminus of HWP1.


6. Although either PSORTII or WOLFPSORT did not find any N-terminal signal peptide, iPSORT does find a signal peptide in the N-terminus of HWP1. This is consistent with and further consolidates the result of TargetP, which says HWP1 has a signal sequence and is likely a secretory protein.

iPSORT Prediction

Predicted as: having a signal peptide

Values used for reasoning

Node	Answer	View	Substring	Value(s)	Plot
1. Signal peptide?	Yes	Average Hydropathy (KYTJ820101)	[6,20]	1.52667 (>= 0.953? Yes)	show



7. Base on the results of prediction programs, as well as previous data in literature. I would propose that HWP1 has a low expression level in the yeast-form cells of *Candida*. The expression of HWP1 will be induced during *Candida* cells change from yeast-form to hyphae-form, and HWP1 will localize to the cell wall.

8. Considering the expression level may affect a protein's subcellular localization, I would take the advantage of homologous recombination in *Candida*, and replace the endogenous gene of HWP1 with a fusion gene in which HWP1 being tagged with a fluorophore. In this case, the fusion gene will be expressed under endogenous promoter of HWP1, and hopefully has a similar expression level as HWP1. Some better-known adhesion (e.g. ALS3) will be used as cell wall marker. I expected to see dim or no fluorescence from fusion protein in yeast-form cells. After shifting conditions and cells becoming hyphae, I expect to see strong signal and the fluorescence localizes to the cells' surface. If I also tagged ALS3 with another fluorophore, I expect to see co-localization of HWP1 and ALS3. A possible problem is the fluorophore may affect the localization of HWP1. An alternative way to get around is to use HWP1 antibody (which we already have) to do immuno-staining. The down-side of this approach is non-specific binding and staining.

Both approaches may require certain amount of HWP1 to exist inside cells. If HWP1 keeps being expressed at a low level, we can make *Candida* cultures, do cell fractionation. Use HWP1 antibody to detect the amount of HWP1 in each fraction.

Appendix

Candida albicans hyphal wall protein 1 (HWP1) gene, complete cds

FEATURES	Location/Qualifiers
source	1..2682 /organism="Candida albicans" /mol_type="genomic DNA" /strain="SC5314" /db_xref="taxon:5476"
<u>gene</u>	445..2682 /gene="HWP1"
<u>mRNA</u>	445..2682 /gene="HWP1"
<u>CDS</u>	503..2407 /gene="HWP1" /note="hyphal surface protein" /codon_start=1 /transl_table=12 /product="hyphal wall protein 1" /protein_id="AAC96368.1" /db_xref="GI:1915979"

/translation="MRLSTAQLIAIAYYMLSIGATVPQVDGQGETEEALIQKRSYDYY
QEPCDDYPQQQQQQEPCDYPQQQQQEPCDYPQQQPQEPCDYPQQQPQEPCDYPQQPQE
PCDYPQQPQEPCDNPPQPDVPCDNPPQPDVPCDNPPQPDIPCDNPPQPDIPCDNPPQP
DQPDNDNPPIPNIPTDWIPNIPTDWIPDIPEKPTTPATTPNIPATTTTTSESSSSSSSSSS
SSTTPKTSASTTPESSVPATTPNTSVPTTSSESTTPATSPPESSVPVTSGSSILATTSE
SSSAPATTPNTSVPTTTTTETKSSSTPLTTTTTEHDTTVVTVTSCSNSVCTESEVTTGVI
VITSKDIYTTYCPLTETTPVSTAPATETPTGTVSTSTEQSTTVITVTSCSESSCTES
EVTGTVVVVTSEETVYTTFCPLTENTPGTDSTPEASIPPMETIPAGSESSMPAGETSP
AVPKSDVPATESAPVPEMTPAGSQPSIPAGETSPAVPKSDVPATESAPPEMTPAGTE
TKPAAPKSSAPATEPSPVAPGTESAPAGPGASSPKSSVLASETSPIAPGAETAPAGS
SGAITIPSSAVVSTTEGAIPTTLESVPLMQPSANYSSVAPISTFEGAGNNMRLTFGA
AIIGIAAFLI"

ORIGIN

1 ggatccaaaa acaaggaatt cggaaattct gacgataaat gtcgactcac aattcattgt
61 aaaaaggag agtttggtta ggctcataat cgcttataat gtacctctaa agtaatctaa
121 aacaaacaca acctttctaa aacctataat aataacccta atggctcaca accgggataa
181 gttagtagc ccagctgttt ttttttgcc ttattttat gactacattt tgttcactt
241 ttgttgcca cttaataacc gttttgcaa cttctcttg taccacctgt atcegcctt
301 ttaacatag caactctgt aaagtcctt tctttccca ctatttatc attcttgaaa
361 tatgtaatca gaatagttt tcaaaaacta taaataacgg tcaaaataac cggtatttt
421 caatttccat tcaactgtt ttctcaaca tatcaaacac aacaggaatc tctatagtc

481 actcgcttt agttcgta atagagatt atcaactgct caacttattg ctatcgctta
 541 ttacatgtta tcaattgggg ccaactgtccc acaggtagac ggtcaagggtg aaacagagga
 601 agctcttatt caaaagagat cttatgatta ctatcaagaa ccatgtgatg attacccaca
 661 acaacaacaa caacaagagc cttgtgatta ccacaacaa caacagcagg aagaaccttg
 721 tgattacca caacaacaac cacaagagcc atgtgactat ccacaacagc cacaagaacc
 781 ttgtgactac ccacaacaac cacaagaacc ttgtgactac ccacaacaac cacaagaacc
 841 ttgcgacaat ccacctcaac ctgatgttcc ttgtgacaat cctctcaac ctgatgttcc
 901 ttgtgacaat cctctcaac ctgatattcc ttgtgacaat cctctcaac ctgatattcc
 961 ttgtgacaat cctctcaac ctgatcagcc tgatgacaat cctctattc caaacattcc
 1021 aaccgattgg attccaata ttccaactga ttggatccca gatattccag aaaagccaac
 1081 aactccagct actactccaa acattctgc tacaactact acttctgaat calcatcttc
 1141 ttcttcttct tcatcatcat ctactactcc aaaaacttct gcttcaacta cacctgaate
 1201 ttctgttcca gctaccactc caaacacttc tgttccaaca acttcttcag aatcaactac
 1261 tccagctact agcccagaaa gttctgttcc agttacttct ggatcatcta ttttagctac
 1321 cacttcagaa tcatcatctg ctccagctac tactccaaat acatctgttc caaccactac
 1381 tactgaaacc aatcatcaa gtactccatt aactactact actgaacatg atacaactgt
 1441 tgtcactgtt acttcatgtt ctaacagtgt ttgtaccgaa agtgaagtta ctactggtgt
 1501 tattgtcatc acatctaaag atactattta caccacttac tgtccattga ctgaaactac
 1561 tccagtttct actgctccag ccaactgaac accaactggg acagtatcca ctctactga
 1621 acaatcaact actgttatta ctgttacttc atgttctgaa agctcttgta ccgaatctga
 1681 agttactact ggtgtgttg ttgttacttc tgaggaaact gtctacacta cattctgtcc
 1741 attgactgaa aacactccag gtactgattc aactccagaa gcttccattc cacctatgga
 1801 aacaattcct gctggttcag aatcatccat gcctgccggg gaaacctctc cagctgttcc
 1861 aaaatcagat gttccagcta ctgaatcagc tccagttcct gaaatgactc cagctggttc
 1921 acaaccatct attctgccg gtgaaacctc tccagctgtt ccaaatcag atgttccagc
 1981 tactgaatct gctctctc ctgaaatgac tccagctggg actgaaacta aaccagctgc
 2041 tccaaaatca tcagctctg ccaactgaacc ttccccagtt gctccaggta ctgaatccgc
 2101 accagctggg ccaggtgctt cttcttctcc aaaatcttct gtttggcta gtgaaacctc
 2161 accaattgct ccaggtgctg aaaccgctcc agctggctca agtgggtgcta ttactattcc
 2221 ggaatctagt gctgtcgtct ctacgactga aggtgctatt ccaactacat tagaatcagt
 2281 tccactcatg caaccatctg ccaattactc aagtgtcgt cctatttcta catttgaagg
 2341 tgctggtaac aacatgagat tgactttcgg tgctgctatt attggtattg ctgcattctt
 2401 gatctaattc taataactga tactaagttt tgtcttttt ttgggatttc tttttttct
 2461 aattttgatt gtttttcaat ttgggtttt caatattatt gacaagagtc attttattga
 2521 atatttgatt tgtttactac attaaagggt ataggtactt ttagtttta aaaattgatt
 2581 tgttcaaatt gtttatcttt ttcttctct tctacttgc ttgtttctg ttttcgggtc
 2641 atagttgata gcttttaata aataccctt ttttttaca at