

Tables

Table 1. Number of adults with a diagnosis of opioid use disorder who were prescribed buprenorphine.

Period	Mean change in number of individuals prescribed buprenorphine (95% CI)	P-value	Mean change in period prevalence (95% CI)	P-value
Segment 1: Monthly change between 1 to 29 months	2600 (2539, 2660)	<0.001	2.56 (2.49, 2.63)	<0.001
Segment 2: Monthly change between 29 to 50 months	2846 (2753, 2939)	<0.001	0.96 (0.91, 1.01)	<0.001
Segment 3: Monthly change between 50 to 85 months	3361 (3320, 3402)	<0.001	0.69 (0.64, 0.74)	<0.001
Difference in slopes between Segment 2 and Segment 1	246 (135, 357)	<0.001	-1.60 (-1.68, -1.51)	<0.001
Difference in slopes between Segment 3 and Segment 2	515 (413, 617)	<0.001	-0.27 (-0.34, -0.20)	<0.001
Difference at 29 months	1612 (133, 3091)	0.033	-2.88 (-4.34, -1.42)	<0.001
Difference at 50 months	1454 (13, 2894)	0.048	1.21 (0.27, 2.15)	0.012

CI, confidence interval

Segment 1: time < 29

Segment 2: time ≥ 29 and time < 50

Segment 3: time ≥ 50

Table 2. Number of adults with a diagnosis of opioid use disorder who were prescribed buprenorphine grouped by SVI quantiles.

Outcome: Number of buprenorphine	SVI = 1		SVI = 2		SVI = 3		SVI = 4	
Period	Mean change in buprenorphine dispensed (95% CI)	P-value	Mean change in buprenorphine dispensed (95% CI)	P-value	Mean change in buprenorphine dispensed (95% CI)	P-value	Mean change in buprenorphine dispensed (95% CI)	P-value
Segment 1: Monthly change between 1 to 29 months	307 (295, 319)	<0.001	424 (412, 436)	<0.001	478 (466, 490)	<0.001	486 (474, 498)	<0.001
Segment 2: Monthly change between 29 to 50 months	361 (342, 380)	<0.001	483 (464, 501)	<0.001	543 (525, 562)	<0.001	526 (507, 544)	<0.001
Segment 3: Monthly change between 50 to 85 months	389 (380, 397)	<0.001	545 (536, 553)	<0.001	669 (660, 677)	<0.001	743 (734, 751)	<0.001
Difference in slopes between Segment 2 and Segment 1	54 (32, 75)	<0.001	59 (37, 81)	<0.001	65 (43, 87)	<0.001	39 (17, 62)	<0.001
Difference in slopes between Segment 3 and Segment 2	28 (7, 48)	0.007	62 (41, 82)	<0.001	125 (105, 146)	<0.001	217 (197, 237)	<0.001
Difference at 29 months	202 (-92, 497)	0.179	335 (41, 630)	0.026	446 (151, 740)	0.003	363 (69, 658)	0.016
Difference at 50 months	-58 (-345, 228)	0.690	42 (-245, 329)	0.774	38 (-249, 325)	0.795	37 (-250, 324)	0.801
Outcome: Period prevalence of buprenorphine	SVI = 1		SVI = 2		SVI = 3		SVI = 4	
Period	Mean change in period prevalence (95% CI)	P-value	Mean change in period prevalence (95% CI)	P-value	Mean change in period prevalence (95% CI)	P-value	Mean change in period prevalence (95% CI)	P-value
Segment 1: Monthly change between 1 to 29 months	3.62 (3.53, 3.70)	<0.001	4.33 (4.23, 4.42)	<0.001	4.57 (4.50, 4.69)	<0.001	4.01 (3.90, 4.13)	<0.001
Segment 2: Monthly change between 29 to 50 months	1.63 (1.48, 1.78)	<0.001	1.79 (1.69, 1.88)	<0.001	1.89 (1.76, 2.02)	<0.001	1.47 (1.34, 1.59)	<0.001
Segment 3: Monthly change between 50 to 85 months	0.95 (0.85, 1.04)	<0.001	1.12 (1.04, 1.49)	<0.001	1.40 (1.31, 1.49)	<0.001	1.74 (1.63, 1.84)	<0.001
Difference in slopes between Segment 2 and Segment 1	-1.99 (-2.15, -1.82)	<0.001	-2.54 (-2.67, 1.82)	<0.001	-2.68 (-2.86, -2.50)	<0.001	-2.54 (-2.71, -2.37)	<0.001
Difference in slopes between Segment 3 and Segment 2	-0.68 (-0.85, -0.51)	<0.001	-0.67 (-0.79, -0.55)	<0.001	-0.49 (-0.65, -0.33)	<0.001	0.27 (0.10, 0.43)	<0.001
Difference at 29 months	-1.93 (-3.43, 0.43)	0.012	-1.71 (-3.19, -0.24)	0.023	-1.91 (-4.01, 0.18)	0.074	-3.10 (-5.24, -0.96)	0.005
Difference at 50 months	-0.51 (-2.51, 1.50)	0.620	-0.23 (-1.65, 1.19)	0.620	0.16 (-1.48, 1.81)	0.848	2.62 (1.14, 4.10)	0.001

SVI, social vulnerability index

CI, confidence interval

Segment 1: time < 29

Segment 2: time >= 29 and time < 50

Segment 3: time >= 50

Figures

Figure 1A. Trend of the number of individuals with opioid use disorder with a buprenorphine prescription.




Figure 1B. Trend of the period prevalence of individuals with opioid use disorder with a buprenorphine prescription.




Figure 2A. Trends of the number of individuals with opioid use disorder with a buprenorphine prescription by Social Vulnerability Index (SVI) Quantiles.




Figure 2B. Trends of the period prevalence of individuals with opioid use disorder with a buprenorphine prescription by Social Vulnerability Index (SVI) Quantiles.

