

MSc Computational Science
joint programme UvA/VU



A DIFFUSION-BASED MODEL OF SPATIAL INTERACTIONS IN *ASPERGILLUS* SPP. GERMINATION



FIRST EXPERIMENTS

Presented by Boyan Mihaylov

December 13, 2024

Supervisor: Prof. dr. Han Wösten, Utrecht University

Examiner: Dr. Jaap Kaandorp, University of Amsterdam

OUTLINE



1 Introduction

2 General Assumptions

Simulation space

Inhibitor

3 Experiment 1: Immediate release

Assumptions

Analytical verification

Results

4 Experiment 2: Slow release

Assumptions

Analytical verification

Results


5 Discussion

INTRODUCTION



Hypothesis

- » Upon inoculation in an aqueous medium, a germination inhibitor diffuses away from the conidium.
 - Experiments have highlighted **1-octen-3-ol** as a likely candidate [2, 3, 4]
- » Once its concentration at the spore falls below a certain threshold, the conidium breaks dormancy and enters a swelling phase.
 - It has been observed that in a **externally non-inhibited** spore swelling begins around **4 hours** from inoculation
- » An increased density of spores drives the local inhibitor concentration high, reducing overall germination.
 - Densities above 10^5 spores/mL exhibit germination inhibition [4, 6]



OUTLINE



1 Introduction

2 General Assumptions

Simulation space

Inhibitor

3 Experiment 1: Immediate release

Assumptions

Analytical verification

Results

4 Experiment 2: Slow release

Assumptions

Analytical verification

Results

5 Discussion

GENERAL ASSUMPTIONS



Simulation space

» 2D lattice \equiv thin layer of medium clamped between 2 impermeable sheets


- $L = 3000 \mu\text{m}$
- Discrete blocks of size $\delta x = \delta y = \delta z = 5 \mu\text{m}$
- $N = 600$ subdivisions
- Lattice volume $V_L = 4.5 \times 10^7 \mu\text{m}^3 = 4.5 \times 10^{-5} \text{ mL}$
- Assuming water as extracellular medium

» Spore \equiv single block in the center of the lattice, inhibitor source

- Volume $V_s = 125 \mu\text{m}^3$
- Top and bottom side of spore are sealed

» Inhibitor - initial concentrations $c(x, y, t = 0)$:

- At spore: $c(x_s, y_s, t = 0) = c_0$
- Elsewhere: $c(x, y, t = 0) = 0$
- Assumed non-replenishable during process



GENERAL ASSUMPTIONS



Inhibitor

» Assumed to have known molecular properties of 1-octen-3-ol:

- Molecular weight: $m_M = 128.21 \text{ g/mol}$
- Density: $\rho = 0.837 \text{ g/mL}$



Figure: 1-octen-3-ol

» Assumed to have similar diffusion properties as glucose:

- Diffusion constant in water at temperature $T = 293.15\text{K}$:
 $D = 600 \mu\text{m}^2/\text{s}$ [5] (in agar only 5% lower[12])
(i.e. a molecule wanders on average $\langle x \rangle = \sqrt{6 \cdot D} = 60 \mu\text{m}$ per second)
- Permeation constant in an artificial lipid bilayer: $P_s = 1.9 \times 10^{-6} \mu\text{m/s}$
(average from [7] and [1], close to [11])
- Octanol-water partition coefficient: $K \approx 10^{-3}$



Figure: D-Glucose

» The documented threshold for 1-octen-3-ol inhibition is $10 \text{ mM} = 0.01 \text{ mol/L}$ [8].

» Once below this level, it is assumed swelling begins immediately

OUTLINE



1 Introduction

2 General Assumptions

Simulation space

Inhibitor

3 Experiment 1: Immediate release

Assumptions

Analytical verification

Results

4 Experiment 2: Slow release

Assumptions

Analytical verification

Results

5 Discussion

EXPERIMENT 1




Inhibitor assumptions

» At $t = 0$, all the inhibitor is contained in a cell wall at the interface between the spore and the medium, having the same diffusion coefficient as water.

- Cell wall thickness: $d \approx 130 \text{ nm}$ (*A. fumigatus*)[10]
- Total cell wall volume:
 $V_{cw} \approx 6 \cdot 25 \mu\text{m}^2 \cdot 0.13 \mu\text{m} = 19.5 \mu\text{m}^3$
- The maximum possible concentration is:

$$c_{cw} = \frac{\rho V_{cw}}{m_M V_s} = 6.5 \text{ mol/mL} \cdot 0.156 = 1.014 \text{ mol/mL} \quad (1)$$

- Equivalently, $N = N_A c_{cw} V_{cw} = 7.04 \times 10^{10}$ molecules.
- The experiments use an initial concentration $c_0 = c_{cw}$.



EXPERIMENT 1




Diffusion assumptions

- » Diffusion is simulated using the diffusion equation (Fick's Law):

$$\frac{\partial c}{\partial t} = D \nabla^2 c \quad (2)$$

- » Simulation run for $t_{\max} = 4 \text{ h} = 14\,400 \text{ s}$
- » The times for reaching threshold concentrations of $c_t = k c_0$ ($k = \{10^{-1}, 10^{-2}, \dots, 10^{-6}\}$) are recorded.
- » To prevent numerical error, the concentration in the simulations is normalised by a factor $\lambda = \frac{1}{c_0}$.



EXPERIMENT 1




Analytical verification

- » Assuming the initial condition is a delta pulse of concentration at coordinates (x_0, y_0)
- » The concentration at an arbitrary position (x, y) at time t can also be computed through the analytical formula[9]:

$$c(x, y, t) = \frac{N^{2/3}}{4\pi Dt} \exp\left(-\frac{(x - x_0)^2 + (y - y_0)^2}{4Dt}\right) \quad (3)$$

- » At the spore site, this equation becomes:

$$c(x_0, y_0, t) = \frac{(V_s c_0)^{2/3}}{4\pi Dt} \quad (4)$$





EXPERIMENT 1



Results

- » Results after 4 hours
- » Numerical vs. analytical
 - Alignment of numerical and analytical solution up to $t \approx 1000$ s, then finite size starts to take effect.
 - Under the assumed c_0 , the concentration in this period is low enough to make this deviation insignificant (few thousands of molecules).
- » Threshold times follow a power law
- » Under the given c_0 (maximum that the cell wall can fit), threshold of 10 mM reached within a few minutes
- » Diffusion is not a limiting factor



OUTLINE



1 Introduction

2 General Assumptions

Simulation space

Inhibitor

3 Experiment 1: Immediate release

Assumptions

Analytical verification

Results

4 Experiment 2: Slow release

Assumptions

Analytical verification

Results

5 Discussion

EXPERIMENT 2




Inhibitor assumptions

- » All the inhibitor is contained within the cell wall volume, separated from the medium by a membrane having the properties of an artificial lipid bilayer.
- » Maximum concentration limit
 - Under homogeneous distribution, standard temperature and pressure:

$$c_{\max} = \frac{\rho}{m_M} = \frac{0.837 \text{ g/mL}}{128.21 \text{ g/mol}} \approx 6.5 \text{ mol/mL} \quad (5)$$

- Equivalently, $N = N_A c_{\max} V_s = 4.89 \times 10^{14}$ molecules.



EXPERIMENT 2




Diffusion assumptions

- » Diffusion through the medium as before
- » Diffusion at the interface follows the flux equation

$$J' = -P_s \Delta c \quad (6)$$

where J' is the flux through the cell wall and Δc is the concentration difference across the membrane.

- » Simulation run for $t_{\max} = 4 \text{ h} = 14\,400 \text{ s}$
- » The times for reaching threshold concentrations of $c_t = kc_0$ ($k = \{10^{-1}, 10^{-2}, \dots, 10^{-6}\}$) are recorded.



EXPERIMENT 2



Analytical verification


- » Since the concentration outside (c_{out}) the spore with a volume V and a surface A diffuses much faster than inside (c_{in}), $c_{\text{out}} \approx \text{const.}$
- » Then, an analytical solution is:

$$c_{\text{in}}(t) = c_{\text{out}} - \Delta c(0)e^{-t/\tau}, \quad (7)$$

where τ is a decay constant:

$$\tau = \frac{V}{AP_s} \quad (8)$$

- » Only 2D communication in the numerical simulation $\rightarrow A$ is actually the **circumference** of the cross section $\times 1 \mu\text{m}$!





EXPERIMENT 2



Results

- » Results after 4 hours
- » Numerical vs. analytical
 - Good alignment of numerical and analytical solution (because medium does not saturate in the given time)
- » Almost linear decrease in concentration
- » Threshold of 10 mM never reached, neither is one tenth of the initial concentration



OUTLINE



1 Introduction

2 General Assumptions

Simulation space

Inhibitor

3 Experiment 1: Immediate release

Assumptions

Analytical verification

Results

4 Experiment 2: Slow release

Assumptions

Analytical verification

Results



5 Discussion

DISCUSSION



Validity of results

- » Known threshold for 1-octen-3-ol inhibition (10 mM)
 - **Experiment 1:** threshold reached within minutes
 - **Experiment 2:** threshold not reached within 4 h
- » The resulting concentrations can be rescaled to different c_0 :
 - **Experiment 1:** If the starting concentration is based on the entire spore volume V_s , then 1-octen-3-ol threshold reached within 30 minutes. More reasonable result, but unrealistic c_0 and permeability assumptions.
 - **Experiment 2:** If c_0 is only slightly above threshold, it can be reached within 4 h.
- » The truth lies in between:
 - Permeation of cell wall is between that of membrane and water
 - If permeating through the membrane, facilitated by proteins (e.g. glucose transporters) → increased P_s ?
 - Cell wall permeability modulated during transition from dormancy.
 - In three dimensions: more surface area for diffusion



DISCUSSION



Tuning the permeation coefficient

- » The analytical solution can be rewritten for P_s :

$$P_s = \frac{V_s}{tA} \ln \left(\frac{\Delta c(0)}{\Delta c(t)} \right) \quad (9)$$

- » Assuming the inhibitor stops being effective below $c_t = 10$ mM
- » Keeping the assumptions about c_0 , d , K , V_s and A_s
- » A hypothetical permeation constant P_s can be computed for a diffusion of 4 hours
- » This results in $P_s = 0.001\,174 \mu\text{m/s}$
- » Remains to be interpreted in terms of physical properties

DISCUSSION



Future outlook

- » Focus on 3D systems
- » Investigating the relationship between inhibitor concentration and spore density
- » Investigating effects on the outward flux when the medium is locally saturated



- [1] G J Bresseleers, H. L. Goderis, and Paul P. Tobback. "Measurement of the glucose permeation rate across phospholipid bilayers using small unilamellar vesicles. Effect of membrane composition and temperature.". In: *Biochimica et biophysica acta* 772 3 (1984), pp. 374–82. URL: <https://api.semanticscholar.org/CorpusID:9584953>.
- [2] Gilma Silva Chitarra et al. "1-Octen-3-ol inhibits conidia germination of *Penicillium paneum* despite of mild effects on membrane permeability, respiration, intracellular pH, and changes the protein composition.". In: *FEMS microbiology ecology* 54 1 (2005), pp. 67–75. URL: <https://api.semanticscholar.org/CorpusID:24273006>.
- [3] Gilma Silva Chitarra et al. "Germination of *Penicillium paneum* Conidia Is Regulated by 1-Octen-3-ol, a Volatile Self-Inhibitor". In: *Applied and Environmental Microbiology* 70 (2004), pp. 2823 –2829. URL: <https://api.semanticscholar.org/CorpusID:19828197>.

BIBLIOGRAPHY II



- [4] Erika Herrero-García et al. “8-Carbon oxylipins inhibit germination and growth, and stimulate aerial conidiation in *Aspergillus nidulans*.”. In: *Fungal biology* 115 4-5 (2011), pp. 393–400. URL: <https://api.semanticscholar.org/CorpusID:33687383>.
- [5] Rudolf Hober. “Physical chemistry of cells and tissues”. In: 1945. URL: <https://api.semanticscholar.org/CorpusID:11597902>.
- [6] Maryam Ijadpanahsaravi et al. “The impact of inter- and intra-species spore density on germination of the food spoilage fungus *Aspergillus niger*.”. In: *International journal of food microbiology* 410 (2023), p. 110495. URL: <https://api.semanticscholar.org/CorpusID:265268197>.
- [7] Henry V Jakubowski et al. “Fundamentals of Biochemistry, a free and new LibreText book for Undergraduate Courses”. In: *The FASEB Journal* 36 (2022). URL: <https://api.semanticscholar.org/CorpusID:248635799>.



- [8] Kana Miyamoto et al. "Formation of 1-octen-3-ol from *Aspergillus flavus* conidia is accelerated after disruption of cells independently of Ppo oxygenases, and is not a main cause of inhibition of germination". In: *PeerJ* 2 (2014). URL: <https://api.semanticscholar.org/CorpusID:8514039>.
- [9] Philip C. Nelson, Marko Radosavljevic, and Sarina Bromberg. "Biological Physics: Energy, Information, Life". In: 2003. URL: <https://api.semanticscholar.org/CorpusID:267787080>.
- [10] Malitha C. Dickwella Widanage et al. "Adaptative survival of *Aspergillus fumigatus* to echinocandins arises from cell wall remodeling beyond β -1,3-glucan synthesis inhibition". In: *Nature Communications* 15 (2024). URL: <https://api.semanticscholar.org/CorpusID:271598394>.
- [11] Reuben E. Wood, Fremont Philip Wirth, and Hywel Morgan. "Glucose permeability of lipid bilayer membranes.". In: *Biochimica et biophysica acta* 163 2 (1968), pp. 171–8. URL: <https://api.semanticscholar.org/CorpusID:29751436>.

BIBLIOGRAPHY IV



- [12] Tong Zhang and Hff Fang. “Effective Diffusion Coefficients of Glucose in Artificial Biofilms”. In: *Environmental Technology* 26 (2005), pp. 155 –160. URL: <https://api.semanticscholar.org/CorpusID:41242504>.