

MSc Computational Science
joint programme UvA/VU



A DIFFUSION-BASED MODEL OF SPATIAL INTERACTIONS IN ASPERGILLUS SPECIES FROM GERMINATION TO OUTGROWTH

THESIS TOPIC PROPOSAL

Presented by Boyan Mihaylov

October 23, 2024

Supervisor: Prof. dr. Han Wösten, Utrecht University

Examiner: Dr. Jaap Kaandorp, University of Amsterdam

OUTLINE



1 Introduction

Aspergilli as model species

Relevance of mycelial morphology

Linking germination to hyphal outgrowth

2 Scope of Proposal

The *Aspergillus* life cycle

Unifying principles

3 Model

Ideas and challenges

Data

Possible extensions

4 Outlook

Tentative Schedule

Next steps

ASPERGILLI AS MODEL SPECIES



Figure: *Aspergillus niger* growing on an onion (Wikipedia).




Figure: Number of published papers on PubMed about filamentous fungi[2].

RELEVANCE OF MYCELIAL MORPHOLOGY



» Dispersed mycelia

- higher enzyme productivity
- high viscosity and non-Newtonian behavior, more energy required for mixing

» Pellets

- easier to handle in industrial processes
- density-dependent diffusion limitations, productivity affected by size and compactness




Figure: Pellet (top) vs. free filament (bottom)[13].

LINKING GERMINATION TO HYPHAL OUTGROWTH



- » Outgrowth lag is correlated with germination time[6].
- » Spore density affects both germination[9] and hyphal morphology[16].
- » Medium provides **activation** in spore germination (signaling) and **nutrition** in mycelial growth (consumption).




Figure: Stages in coagulative pellet growth[19].




Figure: Relationship between germination time and outgrowth lag[6].

OUTLINE



1 Introduction

Aspergilli as model species

Relevance of mycelial morphology

Linking germination to hyphal outgrowth

2 Scope of Proposal

The *Aspergillus* life cycle

Unifying principles

3 Model

Ideas and challenges

Data

Possible extensions

4 Outlook

Tentative Schedule


Next steps

THE ASPERGILLUS LIFE CYCLE



QUESTION


- » Phases of interest - inoculation to hyphal outgrowth (excluding reproduction).
- » How to identify overarching principles among several different and complex phases that can be represented in a much simpler model?



SPORE AGGLOMERATION




- » Motion of conidia through medium guided by diffusion or currents
- » Electrostatic and salt bridging between cell wall surface polysaccharides
- » Aggregation of multiple conidia into future pellet cores



GERMINATION



- » **Carbon sensing leads to germination:** GanB(α)-SfaD(β)-GpgA(γ) is activated by carbon source and triggers a rapid and transient cAMP signal, which stimulates PKA activity[11]. RasA and RasB also play a role [5, 15, 1].
- » **Isotropic swelling:** breaking of dormancy, restructuring of membrane, switch from fermentative to respiratory metabolism, conidia become larger and more vulnerable.
- » **Polarisation and germ tube formation:** establishment of polarity axis, cortical markers recruit proteins to site of polarisation. A hyphal tip forms.




GERMINATION



QUESTION

- » **Carbon sensing leads to germination:** GanB(α)-SfaD(β)-GpgA(γ) is activated by carbon source and triggers a rapid and transient cAMP signal, which stimulates PKA activity[11]. RasA and RasB also play a role [5, 15, 1].
- » **Isotropic swelling:** breaking of dormancy, restructuring of membrane, switch from fermentative to respiratory metabolism, conidia become larger and more vulnerable.
- » **Polarisation and germ tube formation:** establishment of polarity axis, cortical markers recruit proteins to site of polarisation. A hyphal tip forms.




GERMINATION



QUESTION


- » **Carbon sensing leads to germination:** GanB(α)-SfaD(β)-GpgA(γ) is activated by carbon source and triggers a rapid and transient cAMP signal, which stimulates PKA activity[11]. RasA and RasB also play a role [5, 15, 1].
- » **Isotropic swelling:** breaking of dormancy, restructuring of membrane, switch from fermentative to respiratory metabolism, conidia become larger and more vulnerable.
- » **Polarisation and germ tube formation:** establishment of polarity axis, cortical markers recruit proteins to site of polarisation. A hyphal tip forms.



HYPHAL GROWTH




- » Traffic of **vesicles** (EEs) along microtubuli delivers proteins to hyphal tip.
- » Spitzenkörper (SPK) as central hub for distribution of vesicles to the cell membrane via actin filaments.
- » Compartmentalisation of hyphae via **septa**. Opening and closure of **septal pore** can control diffusion between different regions.
- » Secondary polarity axes cause formation of **lateral** branches.



HYPHAL GROWTH




- » Traffic of **vesicles** (EEs) along microtubuli delivers proteins to hyphal tip.
- » **Spitzenkörper (SPK)** as central hub for distribution of vesicles to the cell membrane via actin filaments.
- » Compartmentalisation of hyphae via **septa**. Opening and closure of **septal pore** can control diffusion between different regions.
- » Secondary polarity axes cause formation of **lateral branches**.



HYPHAL GROWTH




- » Traffic of **vesicles** (EEs) along microtubuli delivers proteins to hyphal tip.
- » **Spitzenkörper** (SPK) as central hub for distribution of vesicles to the cell membrane via actin filaments.
- » Compartmentalisation of hyphae via **septa**. Opening and closure of **septal pore** can control diffusion between different regions.
- » Secondary polarity axes cause formation of lateral branches.



HYPHAL GROWTH




- » Traffic of **vesicles** (EEs) along microtubuli delivers proteins to hyphal tip.
- » **Spitzenkörper** (SPK) as central hub for distribution of vesicles to the cell membrane via actin filaments.
- » Compartmentalisation of hyphae via **septa**. Opening and closure of **septal pore** can control diffusion between different regions.
- » Secondary polarity axes cause formation of **lateral branches**.



PELLETS, CLUMPS AND FREE FILAMENTS




- » Spore density, agglomeration and glucosamine production influence the hyphal **macromorphologies**[16].
- » External transport: different densities, e.g. in pellets, cause spatially differentiated nutrient supply, which reflects back on growth[13].
- » Internal transport: cytoplasmic streaming along hyphal network, glucose can be transported even if septal pores are closed. Hyphal fusion improves long-range transport of nutrients[18].



PELLETS, CLUMPS AND FREE FILAMENTS




- » Spore density, agglomeration and glucosamine production influence the hyphal **macromorphologies**[16].
- » **External transport:** different densities, e.g. in pellets, cause spatially differentiated nutrient supply, which reflects back on growth[13].
- » **Internal transport:** cytoplasmic streaming along hyphal network, glucose can be transported even if septal pores are closed. Hyphal fusion improves long-range transport of nutrients[18].



PELLETS, CLUMPS AND FREE FILAMENTS



- » Spore density, agglomeration and glucosamine production influence the hyphal **macromorphologies**[16].
- » **External transport:** different densities, e.g. in pellets, cause spatially differentiated nutrient supply, which reflects back on growth[13].
- » **Internal transport:** cytoplasmic streaming along hyphal network, glucose can be transported even if septal pores are closed. Hyphal fusion improves long-range transport of nutrients[18].



UNIFYING PRINCIPLES



» Diffuse and directed transport through space plays an important role in all phases:

- Motion of conidia leading to agglomeration;
- Transmission of carbon signals to conidial membrane proteins (possible self-inhibition of conidia due to crowding);
- Transport of vesicles in apical growth;
- Diffusion of nutrients and fungal products in media between filaments.

» Therefore, a PDE-based modelling approach seems appropriate, representing concentration changes through discretised space.

UNIFYING PRINCIPLES



- » Diffuse and directed transport through space plays an important role in all phases:
 - Motion of conidia leading to agglomeration;
 - Transmission of carbon signals to conidial membrane proteins (possible self-inhibition of conidia due to crowding);
 - Transport of vesicles in apical growth;
 - Diffusion of nutrients and fungal products in media between filaments.
- » Therefore, a PDE-based modelling approach seems appropriate, representing concentration changes through discretised space.

UNIFYING PRINCIPLES



- » Diffuse and directed transport through space plays an important role in all phases:
 - Motion of conidia leading to agglomeration;
 - Transmission of carbon signals to conidial membrane proteins (possible self-inhibition of conidia due to crowding);
 - Transport of vesicles in apical growth;
 - Diffusion of nutrients and fungal products in media between filaments.
- » Therefore, a PDE-based modelling approach seems appropriate, representing concentration changes through discretised space.

UNIFYING PRINCIPLES



- » Diffuse and directed transport through space plays an important role in all phases:
 - Motion of conidia leading to agglomeration;
 - Transmission of carbon signals to conidial membrane proteins (possible self-inhibition of conidia due to crowding);
 - Transport of vesicles in apical growth;
 - Diffusion of nutrients and fungal products in media between filaments.
- » Therefore, a PDE-based modelling approach seems appropriate, representing concentration changes through discretised space.

- » Diffuse and directed transport through space plays an important role in all phases:
 - Motion of conidia leading to agglomeration;
 - Transmission of carbon signals to conidial membrane proteins (possible self-inhibition of conidia due to crowding);
 - Transport of vesicles in apical growth;
 - Diffusion of nutrients and fungal products in media between filaments.
- » Therefore, a PDE-based modelling approach seems appropriate, representing concentration changes through discretised space.

UNIFYING PRINCIPLES



- » Diffuse and directed transport through space plays an important role in all phases:
 - Motion of conidia leading to agglomeration;
 - Transmission of carbon signals to conidial membrane proteins (possible self-inhibition of conidia due to crowding);
 - Transport of vesicles in apical growth;
 - Diffusion of nutrients and fungal products in media between filaments.
- » Therefore, a **PDE-based** modelling approach seems appropriate, representing **concentration changes through discretised space**.

OUTLINE



1 Introduction

Aspergilli as model species

Relevance of mycelial morphology

Linking germination to hyphal outgrowth

2 Scope of Proposal

The *Aspergillus* life cycle

Unifying principles

3 Model

Ideas and challenges

Data

Possible extensions

4 Outlook

Tentative Schedule

Next steps

MINIMAL MODEL



Minimum viable product of Thesis:

- » A **spatio-temporal model** of growing hyphal morphologies, starting from ungerminated spores.
- » Starting from a **minimal set of assumptions** and gradually incorporating more mechanisms where necessary.
- » The model should be able to **explain and represent** characteristic phenomena from phases like agglomeration, germination and mycelial outgrowth.

MINIMAL MODEL



Minimum viable product of Thesis:

- » A **spatio-temporal model** of growing hyphal morphologies, starting from ungerminated spores.
- » Starting from a **minimal set of assumptions** and gradually incorporating more mechanisms where necessary.
- » The model should be able to **explain and represent** characteristic phenomena from phases like agglomeration, germination and mycelial outgrowth.

MINIMAL MODEL



Minimum viable product of Thesis:

- » A **spatio-temporal model** of growing hyphal morphologies, starting from ungerminated spores.
- » Starting from a **minimal set of assumptions** and gradually incorporating more mechanisms where necessary.
- » The model should be able to **explain and represent** characteristic phenomena from phases like agglomeration, germination and mycelial outgrowth.

EXISTING MODELS

Diffusion-limited aggregation

- » Model by Witten and Sander (1981)[20]
- » Based on the aggregation of randomly moving particles, can also be expressed in terms of concentration-based probabilities
- » The interplay between aggregate and diffusion can be represented well




Figure: PDE-based DLA.




Figure: DLA-generated coral morphology section under laminar flow[12].

EXISTING MODELS



Vector-based growth

- » Many existing models generate the hypha through addition of discrete segments by vector operations.
- » Such formations can be superimposed over concentration gradients or be discretised on a lattice.



Figure: Simulated 3D fungal pellet[3].




Figure: Simulated *Physarum* transport network patterns[10].





Figure: Growth simulation of a fungal pellet, including DLA in spore agglomeration[17].

EXISTING MODELS

The VSC model

- » Vesicle Supply Centre (VSC) model proposed by Bartnicki-Garcia (1989)
- » Moving VSC uniformly radiates vesicles in random angles, which accumulate more material in the cell membrane



CHALLENGES



- » Finding the right resolution to incorporate phenomena of very **different scales**
- » From 2D to 3D representation - 3D simulations are computationally expensive, but assumptions can be more accurate
- » Validation data - access and comparison can be non-trivial, should start early with this

CHALLENGES



- » Finding the right resolution to incorporate phenomena of very **different scales**
- » **From 2D to 3D representation** - 3D simulations are computationally expensive, but assumptions can be more accurate
- » Validation data - access and comparison can be non-trivial, should start early with this

CHALLENGES




- » Finding the right resolution to incorporate phenomena of very **different scales**
- » **From 2D to 3D representation** - 3D simulations are computationally expensive, but assumptions can be more accurate
- » **Validation data** - access and comparison can be non-trivial, should start early with this

INPUT DATA

Experiments by Ijadpanahsaravi et al.

- » Germination data from *A. niger* based on fitted asymmetrical model (P_{max} , τ , d)[8, 9, 7]
- » Can be used to model the distributions from which germination times of individual spores are randomly drawn



VALIDATION DATA

»

- » Available data on 3D pellet morphologies of a hyper-branching *A. niger* strain
- » Can serve as an extreme case validation
- » Alternative sources might also be available




Figure: Process of identifying *A. niger* pellets in 3D microtomography imaging[14].

POSSIBLE EXTENSIONS



- » The asymmetric model for germination by Dantigny[4] yields a good fit with experimental data, but is not necessarily based on **first principles**.
- » Could the framework for our model be used to model explain the statistical results through **spatial interactions** between conidia?
- » Then the input data could become validation data.




Figure: Fitted asymmetrical model[4] on *A. niger* data measured in different media: alanine (A, F), proline (B, G), glutamate (C, H), histidine (D, I), phenylalanine (E, J)[8].

POSSIBLE EXTENSIONS



- » The asymmetric model for germination by Dantigny[4] yields a good fit with experimental data, but is not necessarily based on **first principles**.
- » Could the framework for our model be used to model explain the statistical results through **spatial interactions** between conidia?
- » Then the input data could become validation data.




Figure: Fitted asymmetrical model[4] on *A. niger* data measured in different media: alanine (A, F), proline (B, G), glutamate (C, H), histidine (D, I), phenylalanine (E, J)[8].

POSSIBLE EXTENSIONS



- » The asymmetric model for germination by Dantigny[4] yields a good fit with experimental data, but is not necessarily based on **first principles**.
- » Could the framework for our model be used to model explain the statistical results through **spatial interactions** between conidia?
- » Then the **input data** could become **validation data**.




Figure: Fitted asymmetrical model[4] on *A. niger* data measured in different media: alanine (A, F), proline (B, G), glutamate (C, H), histidine (D, I), phenylalanine (E, J)[8].

OUTLINE



1 Introduction

Aspergilli as model species

Relevance of mycelial morphology

Linking germination to hyphal outgrowth

2 Scope of Proposal

The *Aspergillus* life cycle

Unifying principles

3 Model

Ideas and challenges

Data

Possible extensions

4 Outlook

Tentative Schedule

Next steps

TENTATIVE SCHEDULE



1. Week 43, 2024 - Submission of Thesis Proposal;
2. Week 49, 2024 - Completed literature review (biological principles and existing models);
3. Week 3, 2025 - Minimal model completion and data access setup;
4. Week 9, 2025 - Validation, verification and potential correction of the model;
5. Week 13, 2025 - Analysis of model results;
6. Week 17, 2025 - Model extensions;
7. Week 22, 2025 - Documentation (writing of Thesis).

NEXT STEPS



- » In-depth literature review on relevant phenomena - agglomeration, apical extension, lateral branching.
- » In-depth literature review on related models.
- » Extraction, connection and reduction of relevant principles and their model representations.

THANK YOU!



BIBLIOGRAPHY I



- [1] Tim J. H. Baltussen et al. "Molecular Mechanisms of Conidial Germination in *Aspergillus* spp". In: *Microbiology and Molecular Biology Reviews* 84 (2019).
- [2] Timothy C. Cairns, Corrado Nai, and Vera Meyer. "How a fungus shapes biotechnology: 100 years of *Aspergillus niger* research". In: *Fungal Biology and Biotechnology* 5 (2018).
- [3] Katherine Celler et al. "Structured morphological modeling as a framework for rational strain design of *Streptomyces* species". In: *Antonie Van Leeuwenhoek* 102 (2012), pp. 409 -423.
- [4] Philippe Dantigny et al. "A new model for germination of fungi.". In: *International journal of food microbiology* 146 2 (2011), pp. 176-81.
- [5] Jarrod R. Fortwendel et al. "Aspergillus fumigatus rasA and rasB regulate the timing and morphology of asexual development.". In: *Fungal genetics and biology : FG & B* 41 2 (2004), pp. 129-39.

BIBLIOGRAPHY II



- [6] Maria Gougouli and Konstantinos P. Koutsoumanis. "Relation between germination and mycelium growth of individual fungal spores.". In: *International journal of food microbiology* 161 3 (2013), pp. 231–9.
- [7] Maryam Ijadpanahsaravi, Wieke R. Teertstra, and Han A. B. Wösten. "Inter- and intra-species heterogeneity in germination of Aspergillus conidia". In: *Antonie Van Leeuwenhoek* 115 (2022), pp. 1151 –1164.
- [8] Maryam Ijadpanahsaravi et al. "Minimal nutrient requirements for induction of germination of Aspergillus niger conidia.". In: *Fungal biology* 125 3 (2020), pp. 231–238.
- [9] Maryam Ijadpanahsaravi et al. "The impact of inter- and intra-species spore density on germination of the food spoilage fungus Aspergillus niger.". In: *International journal of food microbiology* 410 (2023), p. 110495.

BIBLIOGRAPHY III



- [10] Jeff Dale Jones. "Characteristics of Pattern Formation and Evolution in Approximations of Physarum Transport Networks". In: *Artificial Life* 16 (2010), pp. 127–153.
- [11] Anne Lafon et al. "The Heterotrimeric G-Protein GanB(α)-SfaD(β)-GpgA(γ) Is a Carbon Source Sensor Involved in Early cAMP-Dependent Germination in Aspergillus nidulans". In: *Genetics* 171 (2005), pp. 71–80.
- [12] Roeland M. H. Merks et al. "Diffusion-limited aggregation in laminar flows". In: *International Journal of Modern Physics C* 14 (2003), pp. 1171–1182.
- [13] Henri Müller et al. "From spores to fungal pellets: A new high-throughput image analysis highlights the structural development of Aspergillus niger". In: *Biotechnology and Bioengineering* 119 (2022), pp. 2182 –2195.

BIBLIOGRAPHY IV



- [14] Henri Müller et al. "Synchrotron radiation-based microcomputed tomography for three-dimensional growth analysis of *Aspergillus niger* pellets". In: *Biotechnology and Bioengineering* 120 (2023), pp. 3244 –3260.
- [15] Nir Osherov and Gregory S. May. "Conidial germination in *Aspergillus nidulans* requires RAS signaling and protein synthesis.". In: *Genetics* 155 2 (2000), pp. 647–56.
- [16] Maria Papagianni and Michael Mattey. "Morphological development of *Aspergillus niger* in submerged citric acid fermentation as a function of the spore inoculum level. Application of neural network and cluster analysis for characterization of mycelial morphology". In: *Microbial Cell Factories* 5 (2006), pp. 3 –3.
- [17] Stefan Schmidededer et al. "Universal law for diffusive mass transport through mycelial networks". In: *Biotechnology and Bioengineering* 118 (2020), pp. 930 –943.

BIBLIOGRAPHY V



- [18] Gero Steinberg et al. "Cell Biology of Hyphal Growth". In: *Microbiology Spectrum* 5 (2017).
- [19] Lukas Veiter, Vignesh Rajamanickam, and Christoph Herwig. "The filamentous fungal pellet—relationship between morphology and productivity". In: *Applied Microbiology and Biotechnology* 102 (2018), pp. 2997 –3006.
- [20] Thomas A. Witten and Leonard M. Sander. "Diffusion-limited aggregation, a kinetic critical phenomenon". In: *Physical Review Letters* 47 (1981), pp. 1400–1403.