

# Some notes on CKM element stuff

## 1 Basic Framework

- So we have the usual BR description in 2HDM:

$$\mathcal{B}r_{exp} = \mathcal{B}r_{SM} \times (1 + r_H)^2$$

- Here,  $(1 + r_H)^2$  describes the WC terms, and the CKM element is part of the SM equation, but if the 2HDM is correct, then what we get from experiment for the value of  $|V_{ij}|^2$  is really  $|\tilde{V}_{ij}|^2 \times (1 + r_H)^2$ 
  - ➡  $r_H = 0$  means that 2HDM is wrong, and we're back to the SM value being correct
  - ➡  $r_H = -2$  means we're in the fine-tuned solution of the 2HDM and what we measure as the SM CKM element is the correct value, although the 2HDM is still correct
  - ➡  $r_H \neq 0, -2$  means that the real CKM element is different from the measured one and we can't use the given SM values
- So if we assume 2HDM to be right, we're probably using the wrong CKM element
- Now consider having

$$\mathcal{B}r_{exp} = \mathcal{B}r_{SM} \times \frac{|\tilde{V}_{ij}|^2}{|V_{ij}|^2} (1 + r_H)^2$$







as our equation, with  $\tilde{V}_{ij}$  being the 'real' CKM element value and  $V_{ij}$  being the measured one in the SM







- This lets us cancel the SM CKM value and input the 'real' one, if there is a difference in their value
- For working in flavio, we need to write any modifications to SM formulae as NP WCs to be added to the SM WCs, so we need to rewrite this:

$$\begin{aligned} \frac{|\tilde{V}_{ij}|^2}{|V_{ij}|^2} (1 + r_H)^2 &= \left\{ \frac{|\tilde{V}_{ij}|}{|V_{ij}|} + \frac{|\tilde{V}_{ij}|}{|V_{ij}|} r_H \right\}^2 \\ &= \left\{ 1 + \left( \frac{|\tilde{V}_{ij}|}{|V_{ij}|} - 1 \right) + \frac{|\tilde{V}_{ij}|}{|V_{ij}|} r_H \right\}^2 \end{aligned}$$

- Writing like this means we can just add on  $\frac{|\tilde{V}_{ij}|}{|V_{ij}|} - 1$  to a WC (e.g. to **CVL\_bumunumu** in **flavio's WET basis** for  $B^+ \rightarrow \mu^+ \nu_\mu$ ) and just multiply the WCs involved with  $r_H$  by  $\frac{|\tilde{V}_{ij}|}{|V_{ij}|}$
- If the CKM element is not changed in the 2HDM, its contribution to the above will go away and we'll be back to our original consideration for the 2HDM; if it is changed, then we will be able to notice it
- This leaves us with 3 free parameters to fit, and I'm not sure if we can do that properly within flavio as it is, but it is simple enough to build up a picture of this modification's impact by doing a range of 2D contours as before for various values of  $\frac{|\tilde{V}_{ij}|}{|V_{ij}|}$
- The issue with this is that then you have to fit for each quark current so that you're only considering a single CKM element at once, and for some, i.e. kaons and pions, their errors in form factors are quite large so I'm not sure if we can really resolve much information from those fits
- I've done some quick contours for various values of  $\frac{|\tilde{V}_{ij}|}{|V_{ij}|}$  for the leptonic decays we've been using, which you can look through at [github/mbr-physics/higgs-proj/tree/master/ckm\\_tests](#) in individual folders for each CKM element consider
- It would probably be worthwhile adding more leptonic decays to these fits for better indication of our validity of choice in  $\frac{|\tilde{V}_{ij}|}{|V_{ij}|}$  but I haven't looked into any extras yet
- Using our leptonic decays so far, we can analyse how  $V_{ud}$ ,  $V_{us}$ ,  $V_{ub}$ ,  $V_{cd}$ , and  $V_{cs}$  can be modified in the 2HDM, but for the other elements, we would have to extract from unitarity
  - ➡ We have all the first row elements here which means we could use unitarity as an extra constraint and test the elements together

- ➡ It might be possible to do similar to above for  $\mathcal{R}(D^{(*)})$  and Vcb so we could then use unitarity constraints on the first two rows
- Since  $\gamma$  comes from the phase of the elements, it shouldn't be affected by the 2HDM, so while just considering 2HDM, we can use the usual method to construct the full matrix from knowing how to modify Vub, Vcb, Vus
- Here's how the leptonic change for Vub and Vus from some of the quick plots ( $\frac{|\tilde{V}_{ij}|}{|V_{ij}|}$  increasing as you go down the page):






## 2 Mapping Out the Modification

- For leptonic decays, using  $r_H$  as above, I've done some heatmaps showing values that  $\frac{|\tilde{V}_{ij}|}{|V_{ij}|}$  can take
- As James discussed in his notes, if the 2HDM is real then the value we measure and assume in the SM is


$$|V_{ij}|^2 = |\tilde{V}_{ij}|^2 (1 + r_H)^2 \implies \frac{1}{(1 + r_H)} = \frac{|\tilde{V}_{ij}|}{|V_{ij}|}$$

So I have used this above for the following plots


- On the left are the plots across the full parameter space we've been using before; on the right are the plots in a reduced range for clarity - this region also corresponds to the approximate region seen in our usual plots for the leptonic decays
- Again, all plots and code are on the github link if you want to look closer




(a)  $m_{B^+} \Rightarrow V_{ub}$ ; on the right, the range is  $\sim 1 \rightarrow 1.4$




(b)  $m_{D^+} \Rightarrow V_{cd}$ ; on the right, the range is  $\sim 0.99965185 \rightarrow 0.99999978$



(c)  $m_{D_s} \Rightarrow V_{cs}$ ; on the right, the range is  $\sim 0.99964 \rightarrow 1.00227$



(d)  $m_{K^+} \Rightarrow V_{us}$ ; on the right, the range is  $\sim 0.9999997 \rightarrow 1.002386$



(e)  $m_{\pi^+} \Rightarrow V_{ud}$ ; on the right, the range is  $\sim 0.9999994 \rightarrow 1.000132$

- For each point in parameter space, I have taken the modification factors for each CKM element from the heatmaps, modified the accepted CKM elements accordingly, and tested for unitarity in the first two rows – this is just using a scan, no statistics yet
- I haven't looked at any modification of  $V_{cb}$  yet either from building a similar sort of heatmap if possible from semileptonics




Figure 3: The black region represents areas where the first two rows of the CKM matrix break unitarity from 2HDM modification as discussed above; the white is where unitarity is still possible within  $2\sigma$ .

### 3 Conclusions