

# Wilson Coefficient for the effective weak Hamiltonian

Mattia Bruno

March 11, 2024

This library is a Mathematica implementation of the Wilson Coefficients for the  $\Delta F = 1$  Weak Hamiltonian computed in *Buras et al.* [arXiv:hep-ph/95123801](#). The goal for this library is to provide the user with a simple-to-use series of functions which automatically evaluate the Wilson Coefficients at a given mass scale  $\mu$  and order in  $\alpha_s$ .

This library contains also all the basic blocks used in the evolution of the Wilson Coefficients, from the strong coupling constant coded up to 4 loops to the evolution matrices  $U$ . Starting from this basic functions it is possible to evaluate the Wilson Coefficients for all transitions related to the effective hamiltonian written below.

The documentation is organized as follows. The colored text represents Mathematica code and each function is described according to this structure:

---

`function[variable1,variable2, ...]`

- `variable1`: explanation
- `variable2`: explanation

Additional comments and explanations go here together with equations to define and describe what `function` does.

# 1 A brief theoretical introduction

By integrating out the weak bosons from the standard model we obtain an effective theory, called weak effective hamiltonian. As an effective theory it loses the property of being renormalizable, meaning that at each order in perturbation theory a new (finite) set of couplings and operators is needed to properly cancel the remaining divergences. Nevertheless, if we work at a fixed order the theory is finite and in the following we will concentrate only on the leading order effective theory.

Composite operators in the effective theory are expected to mix under renormalization, unless there is a symmetry protecting them. This mixing can be generically expressed through a  $Z$ -factor matrix, which can be perturbatively expanded in terms of the so-called anomalous dimension matrices. Once they are known, the operators or alternatively the Wilson Coefficients can be run to low energies and the implementation provided in this library includes effects up to  $O(\alpha_s)$  and  $O(\alpha_e)$ .

The Wilson Coefficients are obtained by comparing (order by order in perturbation theory) amplitudes computed in the standard model against amplitudes computed in the effective theory. Obviously amplitudes are scheme-independent but the two parts of the calculation, the Wilson Coefficients and the low-energy matrix elements, individually depend on the scheme.

## 1.1 The $\Delta F = 1$ Hamiltonian

We are interested here in weak decays where either the strange or the bottom quantum numbers are violated by one unit. In general these processes can be described by a single effective hamiltonian

$$H_W = \frac{G_F}{\sqrt{2}} V_{us}^* V_{ud} \sum_{i=1}^{10} [z_i(\mu) + \tau y_i(\mu)] Q_i(\mu). \quad (1)$$

The 10-operator basis is given below. The scope of this library is to provide the user with a few easy-to-use Mathematica functions to immediately compute the (renormalized) coefficients  $z_i$  and  $y_i$  at a given renormalization scale  $\mu$ . The matrix elements  $Q_i$  must be computed in the same scheme to obtain consistent result at the end.

**Current-Current Operators**

$$Q_1 = (\bar{s}_i u_j)_{V-A} (\bar{u}_j d_i)_{V-A} \quad Q_2 = (\bar{s}u)_{V-A} (\bar{u}d)_{V-A} \quad (2)$$

**QCD-Penguins Operators**

$$Q_3 = (\bar{s}d)_{V-A} \sum_q (\bar{q}q)_{V-A} \quad Q_4 = (\bar{s}_i d_j)_{V-A} \sum_q (\bar{q}_j q_i)_{V-A} \quad (3)$$

$$Q_5 = (\bar{s}d)_{V-A} \sum_q (\bar{q}q)_{V+A} \quad Q_6 = (\bar{s}_i d_j)_{V-A} \sum_q (\bar{q}_j q_i)_{V+A} \quad (4)$$

## Electroweak-Penguins Operators

$$Q_7 = \frac{3}{2} (\bar{s}d)_{V-A} \sum_q e_q (\bar{q}q)_{V+A} \qquad Q_8 = \frac{3}{2} (\bar{s}_i d_j)_{V-A} \sum_q e_q (\bar{q}_j q_i)_{V+A} \quad (5)$$

$$Q_9 = \frac{3}{2} (\bar{s}d)_{V-A} \sum_q e_q (\bar{q}q)_{V-A} \qquad Q_{10} = \frac{3}{2} (\bar{s}_i d_j)_{V-A} \sum_q e_q (\bar{q}_j q_i)_{V-A} \quad (6)$$

## 2 Computation of the Wilson Coefficients

In this section we describe the most important functions of this library: those to compute the Wilson Coefficients at any given energy scale  $\mu$  between 1GeV and  $M_Z$ .

### 2.1 Definition of the Wilson Coefficients

C1[a]  
C2[a,ae]  
C3[a,mt,MW,ae]  
C4[a,mt,MW]  
C5[a,mt,MW]  
C6[a,mt,MW]  
C7[mt,MW,ae]  
C8  
C9[mt,MW,ae]  
C10

- **a**: value of  $\alpha_s$
- **mt**: value of the top quark mass
- **MW**: value of the W boson mass
- **ae**: value of  $\alpha_e$

The Wilson Coefficients at the weak scale are

$$C_1(M_W) = \frac{11}{2} \frac{\alpha_s(M_W)}{4\pi}, \quad (7)$$

$$C_2(M_W) = 1 - \frac{11}{6} \frac{\alpha_s(M_W)}{4\pi} - \frac{35}{18} \frac{\alpha_e}{4\pi}, \quad (8)$$

$$C_3(M_W) = -\frac{\alpha_s(M_W)}{24\pi} \tilde{E}_0(x_t) + \frac{\alpha_e}{6\pi \sin^2 \theta_W} [2B_0(x_t) + C_0(x_t)], \quad (9)$$

$$C_4(M_W) = \frac{\alpha_s(M_W)}{8\pi} \tilde{E}_0(x_t), \quad (10)$$

$$C_5(M_W) = -\frac{\alpha_s(M_W)}{24\pi} \tilde{E}_0(x_t), \quad (11)$$

$$C_6(M_W) = \frac{\alpha_s(M_W)}{8\pi} \tilde{E}_0(x_t), \quad (12)$$

$$C_7(M_W) = \frac{\alpha_e}{6\pi} [4C_0(x_t) + \tilde{D}_0(x_t)], \quad (13)$$

$$C_8(M_W) = 0 \quad (14)$$

$$C_9(M_W) = \frac{\alpha_e}{6\pi} \left[ 4C_0(x_t) + \tilde{D}_0(x_t) + \frac{1}{\sin^2 \theta_W} (10B_0(x_t) - 4C_0(x_t)) \right], \quad (15)$$

$$C_{10}(M_W) = 0, \quad (16)$$

$$x_t = \frac{m_t^2}{M_W^2}. \quad (17)$$

with the QED-related auxiliary functions

$$B_0(x) = \frac{1}{4} \left[ \frac{x}{1-x} + \frac{x \ln x}{(x-1)^2} \right], \quad (18)$$

$$C_0(x) = \frac{x}{8} \left[ \frac{x-6}{x-1} + \frac{3x+2}{(x-1)^2} \ln x \right], \quad (19)$$

$$D_0(x) = -\frac{4}{9} \ln x + \frac{-19x^3 + 25x^2}{36(x-1)^3} + \frac{x^2(5x^2 - 2x - 6)}{18(x-1)^4} \ln x, \quad (20)$$

$$\tilde{D}_0(x_t) = D_0(x_t) - \frac{4}{9}. \quad (21)$$

and the QCD-related ones

$$E_0(x) = -\frac{2}{3} \ln x + \frac{x(18 - 11x - x^2)}{12(1-x)^3} + \frac{x^2(15 - 16x + 4x^2)}{6(1-x)^4} \ln x, \quad (22)$$

$$\tilde{E}_0(x_t) = E_0(x_t) - \frac{2}{3} \quad (23)$$

## 2.2 Evolution of the Wilson Coefficients

`ComputeZ[mu,initAlphaMZ,loop,MZ,aem,init12]`

- **mu**: mass scale at which the coefficients  $v_i$  are computed

- **initAlphaMZ**: initial value of  $\alpha_s$  at the electroweak scale
- **loop**: number of loops of  $\alpha_s$ . Accepted values are 1,2,3,4.
- **MZ**: energy scale corresponding to **initAlphaMZ**. Default is **MZ**.
- **ae**: value of  $\alpha_e$ . Default is 1/129.
- **init12**: bi-dimensional vector containing the values of the Wilson Coefficients  $C_1$  and  $C_2$  at the weak scale. Default corresponds to the results presented in the previous subsection.

The function returns the vector  $\vec{z}(\mu)$  by taking into account possible quark thresholds. If  $\mu > m_b$

$$z_1(M_W) = C_1(M_W), \quad z_2(M_W) = C_2(M_W). \quad (24)$$

$$\begin{pmatrix} z_1(m_c) \\ z_2(m_c) \end{pmatrix} = U_5(\mu, M_W) \begin{pmatrix} z_1(M_W) \\ z_2(M_W) \end{pmatrix}, \quad (25)$$

whereas if  $\mu = m_b$

$$\begin{pmatrix} z_1(m_c) \\ z_2(m_c) \end{pmatrix} = M(m_b) U_5(m_b, M_W) \begin{pmatrix} z_1(M_W) \\ z_2(M_W) \end{pmatrix}, \quad (26)$$

If  $\mu < m_b$

$$\begin{pmatrix} z_1(m_c) \\ z_2(m_c) \end{pmatrix} = U_4(\mu, m_b) M(m_b) U_5(m_b, M_W) \begin{pmatrix} z_1(M_W) \\ z_2(M_W) \end{pmatrix}, \quad (27)$$

whereas if  $\mu = m_c$

$$\begin{pmatrix} z_1(m_c) \\ z_2(m_c) \end{pmatrix} = U_4(m_c, m_b) M(m_b) U_5(m_b, M_W) \begin{pmatrix} z_1(M_W) \\ z_2(M_W) \end{pmatrix}, \quad (28)$$

$$\vec{z}(m_c) = \begin{pmatrix} z_1(m_c) \\ z_2(m_c) \\ \alpha_s/(36\pi)z_2(m_c) \\ -\alpha_s/(12\pi)z_2(m_c) \\ \alpha_s/(36\pi)z_2(m_c) \\ -\alpha_s/(12\pi)z_2(m_c) \\ \alpha_e/(6\pi)F_e(m_c) \\ 0 \\ \alpha_e/(6\pi)F_e(m_c) \\ 0 \end{pmatrix}, \quad (29)$$

with

$$F_e(m_c) = -\frac{4}{9} (3z_1(m_c) + z_2(m_c)). \quad (30)$$

Finally if  $\mu < m_c$

$$\vec{z}(\mu) = U_3(\mu, m_c) \vec{z}(m_c). \quad (31)$$

---

`ComputeY[z,mu,initAlphaMZ,loop,MZ,aem,init]`

- **z**: 10 dimensional vector containing the values of the Wilson coefficients  $z_i(\mu)$
- **mu**: mass scale at which the coefficients  $v_i$  are computed
- **initAlphaMZ**: initial value of  $\alpha_s$  at the electroweak scale
- **loop**: number of loops of  $\alpha_s$ . Accepted values are 1,2,3,4.
- **MZ**: energy scale corresponding to **initAlphaMZ**. Default is **MZ**.
- **ae**: value of  $\alpha_e$ . Default is 1/129.
- **init**: ten dimensional vector containing the values of the Wilson Coefficients  $C_i$  at the weak scale. Default corresponds to the results presented in the previous subsection.

$$y_i(\mu) = v_i(\mu) - z_i(\mu). \quad (32)$$

---

`ComputeV[mu,initAlphaMZ,loop,MZ,aem,init]`

- **mu**: mass scale at which the coefficients  $v_i$  are computed
- **initAlphaMZ**: initial value of  $\alpha_s$  at the electroweak scale
- **loop**: number of loops of  $\alpha_s$ . Accepted values are 1,2,3,4.
- **MZ**: energy scale corresponding to **initAlphaMZ**. Default is **MZ**.
- **ae**: value of  $\alpha_e$ . Default is 1/129.
- **init**: ten dimensional vector containing the values of the Wilson Coefficients  $C_i$  at the weak scale. Default corresponds to the results presented in the previous subsection.

The functions return the vector  $\vec{v}(\mu)$  by taking into account possible quark thresholds.  
If  $\mu > m_b$

$$\vec{v}(\mu) = U_5(\mu, M_W) \vec{C}(M_W), \quad (33)$$

whereas if  $\mu = m_b$

$$\vec{v}(\mu) = M(m_b)U_5(m_b, M_W)\vec{C}(M_W), \quad (34)$$

If  $\mu < m_b$

$$\vec{v}(\mu) = U_4(\mu, m_b)M(m_b)U_5(m_b, M_W)\vec{C}(M_W), \quad (35)$$

whereas if  $\mu = m_c$

$$\vec{v}(\mu) = M(m_c)U_4(m_c, m_b)M(m_b)U_5(m_b, M_W)\vec{C}(M_W), \quad (36)$$

Finally if  $\mu < m_c$

$$\vec{v}(\mu) = U_3(\mu, m_c)M(m_c)U_4(m_c, m_b)M(m_b)U_5(m_b, M_W)\vec{C}(M_W), \quad (37)$$

## 2.3 LO and NLO prescriptions

`ReduceOrder[expr, order]`

- **expr**: expression of the Wilson Coefficients  $y_i$  and  $z_i$
- **order**: accepted values are the strings "LO" and "NLO".

This function takes as input the full expression of the Wilson Coefficients  $y_i$  and  $z_i$  and reduce it to the desired order.



### 3 Basic QCD Functions and input parameters

In this section we describe the functions related to the strong coupling constant and the physical constants which are automatically loaded with this library.

#### 3.1 Strong coupling constant

`beta0[Nc,Nf]`

`beta1[Nc,Nf]`

- `Nc`: number of colors
- `Nf`: number of flavors

First coefficients  $b_0$  and  $b_1$  of the QCD  $\beta$ -function.

---

`alphas[mu,L,Nc,Nf,loop]`

- `mu`: energy scale at which  $\alpha_s$  is computed
- `L`: value of the  $\Lambda$  parameter
- `Nc`: number of colors
- `Nf`: number of flavors
- `loop`: number of loops. Accepted values are 1,2,3,4. Default is 2.

---

`FindLambda[a,mu,Nc,Nf,loop]`

- `a`: input value of  $\alpha_s$
- `mu`: energy scale at which  $\alpha_s$  is computed
- `Nc`: number of colors
- `Nf`: number of flavors
- `loop`: number of loops. Accepted values are 1,2,3,4. Default is 2.

The function returns the value of the  $\Lambda$  parameter which solves the matching equation

$$\alpha_s^{\text{input}} = \alpha_s^{(\text{loop})}(\mu, \Lambda, N_c, N_f) \quad (38)$$

## 3.2 Constants

`MW,MZ,mtop,mbottom,mcharm,alphasMZ`

$$M_W = 80.2\text{GeV} \quad , \quad M_Z = M_W / \sqrt{1 - 0.23^2} , \quad (39)$$

$$m_t = 170\text{GeV} \quad , \quad m_b = 4.4\text{GeV} , m_c = 1.3\text{GeV} , \quad (40)$$

$$\alpha_s(M_Z) = 0.117 \quad (41)$$

## 4 Examples

---

**Example 1 :** computation of the LO Wilson Coefficients at  $\mu = 4\text{GeV}$  using  $\Lambda^4 = 0.325\text{GeV}$  and  $\alpha_s$  in LO. Note that to obtain the Wilson Coefficients in the LO approximation we must use the one-loop running of  $\alpha_s$ , while for the NLO approximation the two-loop running.

Since we want to start from the  $\Lambda$  parameter in the 4-flavor theory we have to first find the  $\Lambda$  parameter in the 5-flavor theory where we match the standard model with effective weak hamiltonian. Hence we start by computing  $\alpha_s$  at the bottom threshold

```
amb = alphas[mbottom,0.325,3,4,1];  
and then we compute  $\Lambda^5$  by matching the 4 and 5 flavors theories  
L = FindLambda[amb,mbottom,3,5,1];  
aMW = alphas[mMW,L,3,5,1];
```

Now we can evolve the Wilson Coefficients down to 4GeV

```
z = ComputeZ[4,aMW,1,MW];  
y = ComputeY[z,4,aMW,1,MW];
```

To obtain the LO results we simply use

```
ReduceOrder[z,"LO"]  
ReduceOrder[y,"LO"]
```

---

**Example 2:** computation of the NLO Wilson Coefficients at  $\mu = 1\text{GeV}$  using  $\alpha_s(M_Z)$

```
z = ComputeZ[4,alphasMZ,2];  
y = ComputeY[z,4,alphasMZ,2];
```

To obtain the NLO results we simply use

```
ReduceOrder[z,"NLO"]  
ReduceOrder[y,"NLO"]
```

## 5 Anomalous Dimension Matrices

In this section we describe the functions to compute the anomalous dimension matrices used in the evolution of the Wilson Coefficients.

### 5.1 QCD anomalous dimension matrices

`gammas0[Nc,Nf,size,Nu]`

- `Nc`: number of colors
- `Nf`: number of flavors
- `size`: size of returned matrix. Default is 10
- `Nu`: number of up quarks. Default is 2

$$\gamma_s^{(0)} \tag{42}$$

---

`gammas1[Nf,size,Nu]`

- `Nf`: number of flavors
- `size`: size of returned matrix. Default is 10
- `Nu`: number of up quarks. Default is 2

$$\gamma_s^{(1)} \tag{43}$$

### 5.2 QED anomalous dimension matrices

`gammae0[Nc,Nf,size,Nu]`

- `Nc`: number of colors
- `Nf`: number of flavors
- `size`: size of returned matrix. Default is 10
- `Nu`: number of up quarks. Default is 2

$$\gamma_e^{(0)} \tag{44}$$

---

`gammae1[Nf,size,Nu]`

- **Nf**: number of flavors
- **size**: size of returned matrix. Default is 10
- **Nu**: number of up quarks. Default is 2

$$\gamma_{se}^{(1)} \quad (45)$$

## 6 Renormalization Group Functions

In this section we describe the core functions used to implement the RG evolution of the Wilson Coefficients.

### 6.1 QCD RG functions

**U0[a1,a2,b0,g0]**

- **a1,a2**: values of  $\alpha_s$  at two different scales  $\mu_1 > \mu_2$
- **b0**: coefficient  $b_0$  of the QCD  $\beta$ -function
- **g0**: LO anomalous dimension matrix  $\gamma_s^{(0)}$

$$U^{(0)}(\mu, m) = V \left( \left[ \frac{\alpha_s(m)}{\alpha_s(\mu)} \right]^{\frac{\tilde{\gamma}^{(0)}}{2\beta_0}} \right)_D V^{-1} \quad (46)$$

with

$$\gamma_D^{(0)} = V^{-1} \gamma^{(0)T} V \quad (47)$$

**J[b0,b1,g0,g1]**

- **b0**: coefficient  $b_0$  of the QCD  $\beta$ -function
- **b1**: coefficient  $b_1$  of the QCD  $\beta$ -function
- **g0**: LO anomalous dimension matrix  $\gamma_s^{(0)}$
- **g1**: NLO anomalous dimension matrix  $\gamma_s^{(1)}$

$$J = V H V^{-1} \quad (48)$$

$$H_{ij} = \delta_{ij} \gamma_i^{(0)} \frac{\beta_1}{2\beta_0^2} - \frac{G_{ij}}{2\beta_0 + \gamma_i^{(0)} - \gamma_j^{(0)}} \quad (49)$$

$$G = V^{-1} \gamma^{(1)T} V \quad (50)$$

$$\gamma_D^{(0)} = V^{-1} \gamma^{(0)T} V \quad (51)$$

**U[a1,a2,b0,g0,J]**

- **a1, a2**: values of  $\alpha_s$  at two different scales  $\mu_1 > \mu_2$
- **b0**: coefficient  $b_0$  of the QCD  $\beta$ -function
- **g0**: LO anomalous dimension matrix  $\gamma_s^{(0)}$
- **J**:  $J$  matrix obtained from **J[b0, b1, g0, g1]**

$$U(\mu, m) = U^{(0)}(\mu, m) + \frac{1}{4\pi} \left[ \alpha_s(\mu) J U^{(0)}(\mu, m) - \alpha_s(m) U^{(0)}(\mu, m) J \right] \quad (52)$$

## 6.2 QED RG functions

**M1[b0, b1, ge0, gse1, J]**

- **b0**: coefficient  $b_0$  of the QCD  $\beta$ -function
- **b1**: coefficient  $b_1$  of the QCD  $\beta$ -function
- **ge0**: LO anomalous dimension matrix  $\geq$
- **gse1**: NLO anomalous dimension matrix  $\gamma_{se}^{(1)}$
- **J**:  $J$  matrix obtained from **J[b0, b1, g0, g1]**

$$M^{(1)} = V^{-1} \left( \gamma_{se}^{(1)T} - \frac{\beta_1}{\beta_0} \gamma_e^{(0)T} + \left[ \gamma_e^{(0)T}, J \right] \right) V. \quad (53)$$

**R[a1, a2, b0, g0, ge0, M1, J]**

- **a1, a2**: values of  $\alpha_s$  at two different scales  $\mu_1 > \mu_2$
- **b0**: coefficient  $b_0$  of the QCD  $\beta$ -function
- **g0**: LO anomalous dimension matrix  $\gamma_s^{(0)}$
- **ge0**: LO anomalous dimension matrix  $\geq$
- **M1**:  $M_1$  matrix obtained from **M1[b0, b1, ge0, gse1, J]**
- **J**:  $J$  matrix obtained from **J[b0, b1, g0, g1]**

$$R(m_1, m_2) \equiv -\frac{2\pi}{\beta_0} V \left( K^{(0)}(m_1, m_2) + \frac{1}{4\pi} \sum_{i=1}^3 K_i^{(1)}(m_1, m_2) \right) V^{-1} \quad (54)$$

$$(K^{(0)}(m_1, m_2))_{ij} = \frac{M_{ij}^{(0)}}{a_i - a_j - 1} \left[ \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_j} \frac{1}{\alpha_s(m_1)} - \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_i} \frac{1}{\alpha_s(m_2)} \right] \quad (55)$$

$$\left(K_1^{(1)}(m_1, m_2)\right)_{ij} = \begin{cases} \frac{M_{ij}^{(1)}}{a_i - a_j} \left[ \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_j} - \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_i} \right] & i \neq j \\ M_{ii}^{(1)} \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_i} \ln \frac{\alpha_s(m_1)}{\alpha_s(m_2)} & i = j \end{cases} \quad (56)$$

$$K_2^{(1)}(m_1, m_2) = -\alpha_s(m_2) K^{(0)}(m_1, m_2) H, \quad (57)$$

$$K_3^{(1)}(m_1, m_2) = \alpha_s(m_1) H K^{(0)}(m_1, m_2) \quad (58)$$

with

$$M^{(0)} = V^{-1} \gamma_e^{(0)T} V \quad (59)$$

### 6.3 Quark threshold matching functions

**Fs[a,z12]**

- **a**: value of  $\alpha_s$  at the mass threshold  $m_c$ .
- **z12**: vector with the two Wilson Coefficients  $z_1$  and  $z_2$  computed at the quark threshold  $m_c$

The function **Fs** returns the vector below

$$\vec{z}(m_c) = \begin{pmatrix} z_1(m_c) \\ z_2(m_c) \\ \alpha_s/(36\pi)z_2(m_c) \\ -\alpha_s/(12\pi)z_2(m_c) \\ \alpha_s/(36\pi)z_2(m_c) \\ -\alpha_s/(12\pi)z_2(m_c) \end{pmatrix}, \quad (60)$$

**Fse[a,z12,ae]**

- **a**: value of  $\alpha_s$  at the mass thresholds  $m_b$  and  $m_c$ .
- **z12**: vector with the two Wilson Coefficients  $z_1$  and  $z_2$  computed at the quark thresholds  $m_c$  and  $m_b$
- **ae**: value of  $\alpha_e$ . Default is 0.

The function **Fse** returns the vector below

$$\vec{z}(m_c) = \begin{pmatrix} z_1(m_c) \\ z_2(m_c) \\ \alpha_s/(36\pi)z_2(m_c) \\ -\alpha_s/(12\pi)z_2(m_c) \\ \alpha_s/(36\pi)z_2(m_c) \\ -\alpha_s/(12\pi)z_2(m_c) \\ \alpha_e/(6\pi)F_e(m_c) \\ 0 \\ \alpha_e/(6\pi)F_e(m_c) \\ 0 \end{pmatrix}, \quad (61)$$

with

$$F_e(m_c) = -\frac{4}{9} (3z_1(m_c) + z_2(m_c)). \quad (62)$$

`M[mu,a,ae,size]`

- **mu**: energy scale of **a**. Values accepted are  $m_c$  and  $m_b$ .
- **a**: value of  $\alpha_s$  at the mass thresholds  $m_b$  and  $m_c$ .
- **ae**: value of  $\alpha_e$ . Default is 0.
- **size**: size of the returned matrix. Default is 10.

$$M(m) = 1 + \frac{\alpha_s(m)}{4\pi} \delta r_s^T + \frac{\alpha_e}{4\pi} \delta r_e^T. \quad (63)$$

The routine automatically uses

$$\begin{aligned} \delta r_s^T &= \frac{5}{18} P(0, 0, 0, -2, 0, -2, 0, 1, 0, 1) \\ \delta r_e^T &= \frac{10}{81} \bar{P}(0, 0, 6, 2, 6, 2, -3, -1, -3, -1) \end{aligned} \quad (64)$$

if  $\mu = m_b$ , whereas if  $\mu = m_c$

$$\begin{aligned} \delta r_s^T &= -\frac{5}{9} P(0, 0, 0, 1, 0, 1, 0, 1, 0, 1) \\ \delta r_e^T &= -\frac{40}{81} \bar{P}(0, 0, 3, 1, 3, 1, 3, 1, 3, 1) \end{aligned} \quad (65)$$

with

$$P^T = (0, 0, -\frac{1}{3}, 1, -\frac{1}{3}, 1, 0, 0, 0, 0), \quad (66)$$

$$\bar{P}^T = (0, 0, 0, 0, 0, 0, 1, 0, 1, 0). \quad (67)$$



## 6.4 QCD+QED RG functions

FullU[a1,a2,ae,b0,g0,ge0,M1,J]

- **a1,a2**: values of  $\alpha_s$  at two different scales  $\mu_1 > \mu_2$
- **ae**: value of  $\alpha_e$
- **b0**: coefficient  $b_0$  of the QCD  $\beta$ -function
- **g0**: LO anomalous dimension matrix  $\gamma_s^{(0)}$
- **ge0**: LO anomalous dimension matrix  $\geq$
- **M1**:  $M_1$  matrix obtained from **M1[b0,b1,ge0,gse1,J]**
- **J**:  $J$  matrix obtained from **J[b0,b1,g0,g1]**

$$U(m_1, m_2, \alpha_e) = U(m_1, m_2) + \frac{\alpha_e}{4\pi} R(m_1, m_2), \quad (68)$$

## 7 Prescriptions

Here we give a detailed explanation of the various subtleties which one encounters in the implementation of the running of the Wilson Coefficients, which are not described nor mentioned in the original paper by Buras et al.

### 7.1 LO vs. NLO prescription

- $C_1, C_2$ : the  $O(1)$  terms correspond to the leading order (0 and 1 for  $C_1$  and  $C_2$  respectively), while the  $O(\alpha_s)$  terms are the NLO corrections.
- $C_3 \dots C_6$ : in this case the  $O(\alpha_s)$  terms appear already at LO, since at typical weak scales the top quark can be integrated out and generate a tree-level graph from a QCD-penguin diagram. Hence all  $m_t$  dependent functions, such as  $B_0$ ,  $C_0$ ,  $D_0$  and  $E_0$  have to be taken already at LO. At NLO the effects of the other quarks travelling in the penguin loop are summarized in the constant terms  $-2/3$  and  $-4/9$  which therefore give the  $O(\alpha_s)$  NLO corrections. Note that the effects of these two constants amount to the redefinitions of  $D_0$  and  $E_0$  into  $\tilde{D}_0$  and  $\tilde{E}_0$ .
- $C_7 \dots C_{10}$ : similarly to QCD-penguin diagrams also for QED penguins the top quark can be integrated out and generate LO contributions at  $O(\alpha_s)$  with  $m_t$ -dependent functions. The same constants given above account for the NLO effects.

Let us consider now the RG evolution matrices.

$$U(\mu, m) = \overbrace{U^{(0)}(\mu, m)}^{\text{LO}} + \overbrace{\frac{1}{4\pi} \left[ \alpha_s(\mu) J U^{(0)}(\mu, m) - \alpha_s(m) U^{(0)}(\mu, m) J \right]}^{\text{NLO}} \quad (69)$$

$$R(m_1, m_2) \equiv -\frac{2\pi}{\beta_0} V \left( \overbrace{K^{(0)}(m_1, m_2)}^{\text{LO}} + \overbrace{\frac{1}{4\pi} \sum_{i=1}^3 K_i^{(1)}(m_1, m_2)}^{\text{NLO}} \right) V^{-1} \quad (70)$$

The quark threshold matching matrices at leading order correspond to the identity matrix only, while at NLO to the full formula given in the previous sections. Note that the prescription for the  $z_i$  coefficients at the charm threshold (recall the functions **Fs** and **Fse**) is at NLO, or in other words  $z_i(m_c) = 0$  for  $i = 3 \dots 10$  at leading order.

## 7.2 Apparent divergence in the $10 \times 10$ $R$ matrix

The  $10 \times 10$  case, with full QCD+QED, leads to an apparent divergence in the matrix  $K_1^{(1)}$  (off-diagonal components)

$$\left( K_1^{(1)}(m_1, m_2) \right)_{ij} = \begin{cases} \frac{M_{ij}^{(1)}}{a_i - a_j} \left[ \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_j} - \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_i} \right] & i \neq j \\ M_{ii}^{(1)} \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_i} \ln \frac{\alpha_s(m_1)}{\alpha_s(m_2)} & i = j \end{cases} \quad (71)$$

due to the presence of two couples of identical eigenvalues of  $\gamma_s^{(0)}$ . However one can easily verify that the divergence cancels by substituting  $a^i = a^j + \varepsilon$  in both numerator and denominator, and by expanding around  $\varepsilon = 0$

$$\frac{x^{a_j} - x^{a_i}}{a_i - a_j} = x^{a_j} \frac{1 - x^\varepsilon}{\varepsilon} = x^{a_j} \frac{1 - (1 + \varepsilon \log x + O(\varepsilon^2))}{\varepsilon} = -x^{a_j} \log x \quad (72)$$

In fact, the same type of calculation leads to the diagonal element of  $K_1^{(1)}$ , which can be re-expressed as

$$\left( K_1^{(1)}(m_1, m_2) \right)_{ij} = \begin{cases} \frac{M_{ij}^{(1)}}{a_i - a_j} \left[ \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_j} - \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_i} \right] & \mathbf{a}_i \neq \mathbf{a}_j \\ M_{ii}^{(1)} \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_i} \ln \frac{\alpha_s(m_1)}{\alpha_s(m_2)} & \mathbf{a}_i = \mathbf{a}_j \end{cases} \quad (73)$$

## 7.3 Apparent divergence in the $N_f = 3$ case

The divergence appearing in the limit  $N_f = 3$  looks similar to the case above, since it arises from the fact that two eigenvalues differ exactly by 1 and therefore the combination  $a_i - a_j - 1$ , present in various denominators, vanishes.

By performing the same type of expansions described above one can check that for pure QCD such a divergence cancels in the combination

$$\alpha_s(\mu)JU^{(0)}(\mu, m) - \alpha_s(m)U^{(0)}(\mu, m)J$$

while in the QED part it cancels automatically inside the matrix  $K^{(0)}$

$$(K^{(0)}(m_1, m_2))_{ij} = \frac{M_{ij}^{(0)}}{a_i - a_j - 1} \left[ \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_j} \frac{1}{\alpha_s(m_1)} - \left( \frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{a_i} \frac{1}{\alpha_s(m_2)} \right] \quad (74)$$

and in the difference

$$K_1^{(1)} - \alpha_s(m_2) K^{(0)} H + \alpha_s(m_1) H K^{(0)} \quad (75)$$

In the code we implemented the substitution  $N_f \rightarrow N_f + \varepsilon$  both in the  $U_0$  and  $J$  matrices and the limit  $\varepsilon \rightarrow 0$  is taken by calling the functions **U** and **R**, or alternatively **FullU**. Only in the definition of  $K^{(0)}$  we hard-coded the solution to the entire expansion