



Angular efficiency dependance of semileptonic W pair decays at the ILC

Matthew Koster
University of Cambridge
FLC group, Summer Student Programme

September 4, 2019

Supervisor: Jakob Beyer

Abstract

An analysis was performed on the muon semi-leptonic decay channel of a pair of W bosons produced in 500 GeV center of mass energy $e_L^- e_R^+$ collisions at the ILC. The invisible decay products were reconstructed using conservation laws and the visible reconstructed data. Angular distributions defining the semileptonic decay were extracted from generator level particles and cuts were applied on the reconstructed particles to obtain the angular dependance of the reconstruction efficiency. The analysis reveals an angular dependance of the efficiency which is not currently modelled in the Electroweak polarisation fit for the ILC.

Contents

12	Contents	
13	1 Introduction	2
14	2 Final state 4-momenta analysis	2
15	2.1 Beam Background Removal	2
16	2.2 4-momenta Reconstruction	2
17	2.3 Neutrino and ISR Corrections	3
18	2.3.1 The invisible system	3
19	2.3.2 Reconstructing the leptonically decaying W boson	4
20	2.3.3 Center of Mass frame	4
21	2.3.4 ISR invariant mass	4
22	2.3.5 Beam Background Removal	5
23	2.3.6 Negligable ISR case	5
24	2.4 Angle extractions	6
25	3 Angular dependance of efficiency	7
26	3.1 Cut Flow	7
27	3.2 Angle Cut Efficiencies	9
28	4 Conclusion	10
29	5 References	12
30	6 Acknowledgments	13
31	7 Appendix	14
32	7.1 Log Figures	14
33	7.2 Derivation of the ISR energy E_γ with non-trivial m_γ and m_ν	14




Figure 1: Lowest order Feynman diagram of semileptonic W pair decay in the $\mu\bar{\nu}q\bar{q}'$ final state at e^+e^- colliders, with two ISR photons emitted. Alternately the W^+ could decay leptonically and the W^- hadronically, which would result in a $\mu^+\nu q\bar{q}'$ final state (not drawn).

1 Introduction

DO ME

2 Final state 4-momenta analysis

This analysis reconstructs the 4-momenta of the particles in the hard collision, using data from 500 GeV ILC simulations containing generator level particles and reconstructed particles. Then several variables used in the analysis in Section. 3 are evaluated and stored in a .root file. In order to do this it must use some kinematically derived formulae (Section. ??) to reconstruct the invisible 4-momenta contribution of the neutrino and any initial state radiation (ISR)¹. This section contains the details of each part of this analysis.

2.1 Beam Background Removal

In a particle collider, to increase the probability of a hard collision, a 'bunch' [Herr:941318] of particles are collided and due to the low cross-section one may expect only one hard interaction to occur. These other particles, however, do interact with each other which may produce particles that leave traces in the detector and so will generate so-called beam background. This is modelled in the collision simulation so these particles get reconstructed. These reconstructed particles from the background must be removed before analysis of the hard collision can be conducted.

The FastJetProcessor [1] was used to conduct such beam background removal from the reconstructed particle collection. Another method of background removal is by using generator level information to help remove the background from the reconstructed particles², this was done with the TrueJet processor [2]. This is a liberty that is only possible because the collisions are simulated, therefore it is not possible at an active detector and is referred to in this report as 'cheating overlay removal'. The later method was used to explore the sensitivity of the reconstruction to this beam background removal in Section. 2.3.5.

2.2 4-momenta Reconstruction

For the generator level particles, each of the particles in the hard collision were directly extracted from the collection using their element number. It was then a simple task of inputting their energies and momenta into a TLorentzVector [fons_rademakers_2019_3336325]. The W

¹Photons radiated by e^- or e^+ before collision

²A different steering file was used for this set up, further discussed in Section. ??

63 bosons 4-momenta were then calculated by addition of the 4-momenta of its decay products.

64
65 The IsolatedLeptonTaggingProcessor [3] was used to extract the hard collision lepton³ from
66 the reconstructed particles. The FastJetProcessor [1] extracts the two reconstructed quarks.
67 The 4-momenta of these final state particles and subsequently that of the hadronically decaying
68 W boson (W_{had}) could then be reconstructed⁴.

69
70 Reconstruction of the leptonically decaying W boson (W_{lep}) required a bit more thought, this
71 is because of the neutrino and the ISR photons which leave no signal in the detector. This
72 challenge is discussed in detail in the following section.

73 2.3 Neutrino and ISR Corrections

74 2.3.1 The invisible system

The leptonically decaying W boson was reconstructed by summing the 4-momenta of the extracted lepton and the neutrino, however, the neutrino does not leave a signal in the detector. In addition, there is some initial state radiation of photons (ISR) which are aligned enough to the beam pipe to not be detected. This results in an 'invisible' system that is not at all detected by the detector, and, to obtain complete information about the W bosons, this invisible system had to be reconstructed.

The chosen method for reconstruction of the invisible system arises purely from conservation laws⁵. The reconstructed visible system was considered as being the hadronically decaying W boson plus the isolated lepton ($p^\mu = (E, p_x, p_y, p_z)$), and the invisible system as the accompanying neutrino and an ISR photon, travelling parallel to the beam. The total center of mass energy was assumed to be 500 GeV and the hard interaction was assumed to take place in the center of mass frame. Conservation of momentum and energy then exactly defines the 4-momenta of the ISR photon and the neutrino. In the simulation there are two ISR photons emitted. This means that when combining the two photons into one 'photon', this 'photon' may have a non zero invariant mass. Furthermore, the ISR photon could be going in either direction with respect to the beam axis, resulting in two solutions.

$$E_\gamma = \frac{\lambda(500 - E) \pm p_z \sqrt{\lambda^2 - [(500 - E)^2 - p_z^2]m_\gamma^2}}{(500 - E)^2 - p_z^2} \quad (1)$$

75 where $\lambda = \frac{1}{2}[(500 - E)^2 - p^2 + m_\gamma^2 - m_\nu^2]$ has been defined for convenience and no absolute
76 values have been assumed.

77
78 The solution that reconstructs the invariant mass of the W boson closer to the currently ac-
79 cepted value of 80.3 GeV [5] is selected. This may shift more of the background into the peak
80 introducing a bias and so the appropriate cut was made to mitigate this [6].

81
82 The rest of the unknown variables of momentum and energy are exactly defined from this
83 and can be easily evaluated. As a result the 4-momenta of all of the particles in the hard
84 interaction are known, as required for the analysis.

³If it correctly reconstructs the lepton, sometimes it returned zero particles, sometimes more than one, this was cut on later

⁴It was noted that running the LcplusProcessor [4] did improve this reconstruction

⁵ a full mathematical derivation is included in the appendix for completeness (Appendix. 7.2)

86 A simplified form of this equation with $m_\gamma = m_\nu = 0$ is used in Ivan's thesis. Equation. (1)
 87 correctly simplifies to this solution as shown in the appendix.

$$E_\gamma = \frac{(500 - E)^2 - p^2}{1000 - 2E \mp 2p_z} \quad (2)$$

It is worth noting at this point that it is numerically possible for the particle reconstruction to result in a visible 4-momenta with a negative λ . It can be seen in the $m_\gamma = m_\nu = 0$ solution that this corresponds to the invisible part of the system having an imaginary invariant mass.

$$\lambda_{m_\gamma, m_\nu=0} \propto (500 - E)^2 - p^2 = E_{inv}^2 - p_{inv}^2 = m_{inv}^2 \quad (3)$$

88
 89 This is obviously not physical and will affect the solutions attained above. For example as
 90 $p_\gamma = \pm E_\gamma$, a negative energy will reverse the direction of the photon. This needs to be handled
 91 carefully in the code to avoid inconsistent solutions.

92
 93 It can be analytically shown that Equation. 1 leads to Equation. 2 when simplified, there
 94 is however one subtlety when this is done computationally. The $\sqrt{\lambda^2}$ will computationally
 95 evaluate as $|\lambda|$, so if λ is negative it will in effect reverse the effect of the \pm in Equation. 1
 96 and lead to the opposite solution of the simplified form. So it is clear that these negative λ 's
 97 may cause some issues if the contribution from the m_γ in non-negligible. This was not explored
 98 further, because in Section. 2.3.4 it is shown that including the non-zero m_γ contribution from
 99 the Monte Carlo has little effect on the solutions attained and so Equation. 2 is used with the
 100 \pm swapped accordingly.

101 2.3.2 Reconstructing the leptonically decaying W boson

102 Using the method in Section. 2.3.1 the 4-momentum of the leptonically decaying W boson was
 103 reconstructed. As a test of the performance of this, the accuracy of the reconstructed invariant
 104 mass of this boson was evaluated. By exploring different several variations of this reconstruction
 105 the sensitivity of the ISR energy solution was tested in the following analysis.

106 2.3.3 Center of Mass frame

107 The e^+e^- collision is incident at a non zero angle. Consequently the total system in detector
 108 frame has a non-zero 3-momentum $p^\mu = (500 \sin(\frac{0.014}{2}), 0, 0)$ GeV. In theory addition of a boost
 109 into the centre of mass frame should result in an improvement in the reconstruction, because the
 110 assumption that the total energy is equal to the center of mass energy is only satisfied in this
 111 frame. In fact we see that it makes a considerable improvement to the mass peak (Figure. 2a)
 112 and is kept in the following analysis.

113 2.3.4 ISR invariant mass

114 Another assumption that was tested was the non-zero invariant mass of the ISR photon. As
 115 two ISR photons are emitted, but only one photon is modelled in the reconstruction, this
 116 modelled photon need not satisfy this constraint. The true invariant mass was obtained from
 117 the generator level particles and inputted into Equation. 1, which is derived with this constraint
 118 relaxed. Again this is a 'cheat' as this information would not be available in a real detector.
 119 This addition had a negligible effect on the reconstructed W mass (see Figure. 2b), which could
 120 be due to the size of the photon mass indeed being negligible, even if non-zero.




Figure 2: The effect on the reconstructed invariant mass of the leptonically decaying W with different reconstruction methods. The simulated e^+e^- collision is not head on, but with a finite momentum in the x direction, (a) shows the effect of boosting into the ZMF before applying the reconstruction method. (b) shows the effect of inputting a non-zero invariant ISR photon mass, obtained from generator level information, into the reconstruction.

2.3.5 Beam Background Removal

The third check that was performed was the sensitivity of the reconstructed $m_{W_{lep}}$ to the beam background. This was done by comparing the two different background removal methods suggested in Section. 2.1. It was found that cheating the beam background removal improved the reconstruction of the two quarks, which can be seen in the reconstruction of the invariant mass of the hadronically decaying W boson (Figure. 3). In particular this cheating greatly suppresses an overestimation tail observed in the properly reconstructed particles. This in turn improves the reconstruction of the visible portion of the system. The $m_{W_{lep}}$ on the other hand was generally unsensitive to this change. Explaining this is non-trivial due to the fairly complicated nature of the E_γ formula.

2.3.6 Negligible ISR case

The ISR energy formula regularly reconstructs a wide range of energies (of order ± 20 GeV) when the true ISR energy is less than 0.5 GeV (see Figure. 4a). Addition of a third solution to the photon energy formula was implemented, where the photon energy, and hence momentum, was set exactly equal to zero. The conservation laws then directly defined the 4-momenta of the neutrino, as it was then the only contribution to the invisible part of the system. As before the estimate of $m_{W_{lep}}$ was calculated using all three solutions to E_γ and the solution that more accurately reconstructed this mass was chosen. Once again this selection may draw more of the background into the peak and introduce a bias. The extent of this is not explored in this report, but it is important to keep into consideration in the following, and any further, analysis. This seems to improve the estimation of $m_{W_{lep}}$ as seen in (Figure. 4b), although it is unclear if this is a true improvement in the method due to this bias.

As another check of the performance of this edit, A histogram of just $E_\gamma^{Reco} - E_\gamma^{MC}$ was created (Figure. 4c), to see if this improvement in $m_{W_{lep}}$ was corresponded to an improvement in the estimation of E_γ . Here it can be seen that the addition of the $E_\gamma = 0$ solution reduces the number of under and over estimates quite significantly. The highest peak at zero, however, claimed




Figure 3: Plots displaying the effect of cheating the beam background removal on the reconstructed invariant mass of the two W bosons, (a) the leptonically decaying W and (b) decaying hadronically.




Figure 4: (a) Reconstructed ISR photon energy against the generator level ISR energy. (b) Reconstruction of the leptonically decaying W mass for the three different E_γ calculation methods. (c) Log histogram showing the reconstruction of E_γ from the three calculation methods.

by the solution where there is no ISR considered at all (green). This means that sometimes an over/under-estimation of the photon energy returns a better reconstruction of the $m_{W_{lep}}$.

2.4 Angle extractions

From this analysis, complete information about the final state particles, and hence the intermediate W pair, is obtained. From this we can evaluate the angular distributions required for the next section of the analysis.

The semileptonic decay of a pair of W bosons is fully described by just 6 angles; the polar coordinates of the W— in the center of mass frame; the polar coordinated of the lepton in the center of mass frame of the W it decayed from; and the polar coordinated of one of the quarks in the center of mass frame of the boson it decayed from. The ϕ coordinate of the W— is expected to be uniformly distributed and so returns no information about the chiral structure and is not considered in the fit. Further, it is hard to assign a charge to the quarks in their reconstruction,

so it is not possible to consistently chose one the the quarks to measure its angles. Hence the quarks polar coordinates are also not considered in this analysis. This leaves just 3 angles as defined in Figure. 5, namely the theta coordinate of the W^- in the centre of mass frame (θ_{W^-}), and the theta and phi coordinates of the lepton in the rest frame of the W_{lep} it decayed from (θ_l^* and ϕ_l^*). The polar coordinate system is the same in both frames, with the z axis aligned to the beam-pipe. The same lorentz boost framework used in Section. 2.3.3 was implemented here and so this was a computationally very simple task, and the extracted angles were put in the root tree.

The resulting angular distributions that were obtained are visible in Figure. 7, where the θ angles have been displayed in terms of their cosine for comparison with previous results [7].

3 Angular dependance of efficiency

The following analysis is performed on data generated using the reconstruction techniques described above, choosing the best of all three E_γ solutions, with $m_\gamma = m_\nu = 0$. Unless otherwise explicitly stated, the beam background removal was not cheated but the lorentz boost into the centre of mass frame was performed. The analysis was conducted considering only the muon signal of the data set, as this is the focus of this analysis.

3.1 Cut Flow

All the same cuts as in Ivans thesis [6] were sequentially applied to obtain a cut flow histogram, which tests the efficiency of each of the cuts. The results of this analysis were then compared to Ivan's to see how it is performing. Further, the flow diagram was created again by cheating the beam background removal, and by applying the additional initial cut that the ISR energy is small ($E_\gamma^{MC} < 1$ GeV). The final plot was made to see if the ISR made a considerable contribution to the efficiency of the cuts.

There are a few subtle differences between the cuts that Ivan and the cuts made in this analysis, due to the reconstruction methods each implemented. The first was the cut on the jet reconstruction variables y_+ and y_- . In this analysis FastJets [1] was forced to reconstruct two jets, corresponding to the 2 quarks, and physically this is the minimum number of jets that can occur, as single quark cannot be created. This means the cut on the y_- variable that Ivan makes is unphysical in this analysis. Ivan applies this cut because he also considered final states with 4 quarks. The lepton cut that was made here make is also different to Ivans. The IsolatedLeptonTaggingProcessor was used, so the implemented cut was that a single isolated lepton is reconstructed. There are a considerable number of events where the processor reconstructed zero particles, and so this cut can be significant. Ivan on the other hand used a jet reconstruction processor with a specific energy cut to reconstruct isolated leptons. The final difference is that Ivan makes a 'charged lepton' cut, which is not described in his thesis. A similar cut is not made in this analysis and so the corresponding efficiency will not change across this cut.

For convenience the cuts and efficiencies are tabulated in Table. 1, and the resulting total flow diagram can be seen in (Figure. 6).




Figure 5: Angle definition for the 4-fermion final state from W^- pair production in the semileptonic channel in which the W decays leptonically. In the top part, the production angles θ_{W^-} and θ_{W^+} are defined by the axis of the initial colliding particles and the direction for the respective boson. The azimuthal angles ϕ_{W^-} and ϕ_{W^+} describe the rotation around the axis of the initial colliding particles.

The bottom picture shows the angular definition of the decay products of the W^{lep} boson in its rest frame. It shows the polar angles θ_l^* and $\theta_{\bar{\nu}}^*$ and the azimuth angles ϕ_l^* and $\phi_{\bar{\nu}}^*$ of the corresponding leptons. The $*$ denotes quantities in the rest frame of the W^{lep} boson. (Edited from [7])

Table 1: Selection efficiency of sequentially applied cuts. Where the post ISR correction m_W^{lep} was calculated using all 3 possible E_γ solutions. (*) Indicates cuts where the current and Ivan's cuts differ, as discussed in the text.

Order	Cut description	Efficiency [%]			Ivan's Results n = 107233	
		My Results		no cheat		cheat
		n = 2129	n = 99419			
0	muon signal	100.00	100.00	100.00	100.00	
1	track multiplicity ⁶ $n_{tracks} \geq 10$	97.13	97.01	96.23	99.996	
2	center of mass energy $\sqrt{s} > 100$ GeV	92.29	91.69	84.35	97.96	
3	total transverse momentum $P_T > 5$ GeV	91.16	90.47	83.28	96.69	
4	total energy $E_{SUM} < 500$ GeV	89.66	89.28	82.70	95.36	
5	$\ln(y_+) \in [-12, -3]$ (*)	88.69	88.08	82.47	95.01	
6	1 lepton found (*)	80.65	80.77	81.50	78.75	
7	pre ISR correction $m_W^{lep} \in [20, 250]$ GeV	78.23	77.94	77.84	76.61	
8	tau discrimination ⁷	76.05	75.60	75.73	74.07	
9	charged lepton (*)	76.05	75.60	75.73	73.51	
10	isolation variable ⁸ $\Delta\Omega_{iso} > 0.5$	76.01	75.58	75.72	73.42	
11	post ISR correction $m_W^{lep} \in [40, 120]$ GeV	72.90	72.77	72.33	70.13	
12	post ISR correction $m_W^{had} \in [40, 120]$ GeV	63.21	62.92	70.52	66.93	
13	$\cos\theta_W > -0.95$	63.02	62.65	70.21	66.78	

There is a considerable difference in the cut efficiencies before the 'lepton' cut. In all of the events where no isolated leptons are reconstructed, the leptonic W boson is reconstructed entirely from the invisible neutrino, will be completely wrong. The efficiency of the cheated and un-cheated background removal reconstructions converge on this lepton cut, suggesting that the differences observed before the cut are due to these incorrectly reconstructed events. When this lepton cut is applied to the current results before the rest of the cuts, the large discrepancy is not observed, providing support to this hypothesis (see Appendix. ??).

After the lepton cut, the efficiency of the reconstruction is higher than in previous studies, until the cut on the m_W^{had} where it is considerably worse. By looking at the cheated results though, which don't drop in efficiency as dramatically, it can be suggested that this is due to the performance of the beam background removal process. Further, after the lepton cut the $E_\gamma^{MC} < 1$ GeV signal is consistently performing better than the full signal, which suggests that large ISR energies reduces the efficiency of the reconstruction and so the detector.

3.2 Angle Cut Efficiencies

In this section the final result of the analysis, to evaluate the angular dependency of the cut efficiency, is discussed. The efficiency was obtained for both the full signal and the $E_\gamma^{MC} < 1$ GeV signal, the results of which are shown in Figure. 7.

⁶track multiplicity was taken as the number of reconstructed charged particles.

⁷ τ_{discr} defined by $\tau_{discr} = \left(\frac{2E_{lep}}{\sqrt{s}}\right)^2 + \left(\frac{m_W^{lep}}{m_W^{true}}\right)^2$

⁸ $\Delta\Omega_{iso}$ defined as,

$$(\phi_{lep} - \phi_{had}) < \pi \rightarrow \Delta\Omega_{iso} = \sqrt{(\theta_{lep} - \theta_{had})^2 + (\phi_{lep} - \phi_{had})^2} \quad (4)$$

$$(\phi_{lep} - \phi_{had}) \geq \pi \rightarrow \Delta\Omega_{iso} = \sqrt{(\theta_{lep} - \theta_{had})^2 + (2\pi - |\phi_{lep} - \phi_{had}|)^2}. \quad (5)$$




Figure 6: Cut flows comparing the results of this analysis (with and without cheating overlay removal) to previous studies. The results of this analysis considering a signal where $E_\gamma < 1$ are also displayed. The cuts on the x axis are as defined in Table. 1.

The $\cos(\theta_{W-})$ distribution is statistically limited towards -1, in the backwards direction, greatly reducing the efficiency. Due to the low mass of the neutrino, it becomes more virtual as more momentum is transferred to it, reducing the cross section. As a result most of the momentum transfer is to the W bosons, aligning it to the lepton it decayed from. In this case the W^- momentum is preferably aligned to the momentum of the e^- and so we see the angular distribution is highly biased towards the $\cos(\theta_{W-}) = 1$ side. It is, however, reasonably constant throughout the rest of the distribution. The efficiency has a clear angular dependance on $\cos(\theta_l^*)$, with lower efficiency in regions closely aligned to the beam-pipe. Due to the coordinate system used this is not trivially explained, however for W bosons highly aligned to the beam pipe, the lepton appears to decay in a preferably transverse direction in the center of mass frame of such a boson. The ϕ_l^* of efficiency has a uniform efficiency as well as angular distribution, somewhat expected due to the uniformity of the W bosons ϕ distribution.

The $E_\gamma^{MC} < 1$ GeV signal has a higher efficiency than the full signal, but the angular dependance as a similar form. The magnitude of the efficiency differs by a constant factor.

4 Conclusion

DO ME




Figure 7: (a), (b) and (c) are the extracted angles as defined in Figure. 5. (d), (e) and (f) are the associated cut efficiencies, after applying the cuts outlined in Table. 1.

5 References


- [1] Matteo Cacciari, Gavin P. Salam, and Gregory Soyez. “FastJet User Manual”. In: *Eur. Phys. J. C* 72 (2012), p. 1896. DOI: 10.1140/epjc/s10052-012-1896-2. arXiv: 1111.6097 [hep-ph].
- [2] Mikael Berggen. “Truth information : RecoMCTruthLinker and TrueJet”. In: *Software & Analysis Pre-Meeting to ILD Workshop* (2018). URL: <https://agenda.linearcollider.org/event/7839/contributions/40940/>.
- [3] C. Duerig and J. Tian. “isolated lepton finder”. In: *GitHub repository* (2019). URL: <https://agenda.linearcollider.org/event/6787/session/10/contribution/17/material/slides/0.pdf>.
- [4] J. Strube. “LCFIPlus”. In: (2015). URL: <https://confluence.slac.stanford.edu/display/ilc/LCFIPlus>.
- [5] M. Tanabashi et al. “Review of Particle Physics”. In: *Phys. Rev. D* 98 (3 Aug. 2018), p. 030001. DOI: 10.1103/PhysRevD.98.030001. URL: <https://link.aps.org/doi/10.1103/PhysRevD.98.030001>.
- [6] I. Marchesini. “Triple Gauge Couplings and Polarization at the *ILC* and Leakage in a Highly Granular Calorimeter”. Hamburg University, Diss., 2011. Dr. Hamburg: Hamburg University, 2011. URL: <https://bib-pubdb1.desy.de/record/94888>.
- [7] Robert Karl. “From the Machine-Detector Interface to Electroweak Precision Measurements at the ILC Beam-Gas Background, Beam Polarization and Triple Gauge Couplings”. Bitte POF/Facility überprüfen, vielen Dank, Maren Stein; Dissertation, Universität Hamburg, 2019. Dissertation. Hamburg: Universität Hamburg, 2019, p. 330. DOI: 10.3204/PUBDB-2019-03013. URL: <https://bib-pubdb1.desy.de/record/424633>.

267 6 Acknowledgments

268 I would like to thank my supervisor Jakob Beyer for giving me a cool project and always
269 providing instant support whenever I asked for help. Not only this but he provided help to the
270 other summer students in our group without complaint which was really above and beyond.
271 I'd also like to thank the rest of the FLC group for providing a welcoming environment for us,
272 inviting us on group excursions in and outside of work. Further, Daniel Heuchel and Vladimir
273 Borchanikov were great office mates and were a pleasure to work alongside.
274 It was great getting to know the other summer students in the FLC group - Sukee Dharani,
275 Andreas Loeschke Centeno, Peter McKeown and Lorenzo Cotrozzi. They are all wonderful
276 people and made the long hours writing reports and presentations a lot more bearable, ordering
277 pizza to the office was a terrific idea.
278 I'd like to thank all the summer students who organised awesome group events on the weekends,
279 like our excursions to Cuxhaven and Lübeck, and made any food for the community, the cultural
280 food picnic on the last Sunday was fantastic. In particular Jana Wittmann, who organised many
281 events and made many cakes, even delivering some at 5 am!
282 Finally I'd like to thank Olaf Behnke and the organisation team for all their hard work in
283 making this summer programme run smoothly

7 Appendix

7.1 Log Figures



7.2 Derivation of the ISR energy E_γ with non-trivial m_γ and m_ν

Our initial assumptions are that the system is in the center of mass frame with an invariant mass of 500 GeV. The system contains a visible 4-momentum $p^\mu = (E, p_x, p_y, p_z)$ and an invisible 4-momentum (a neutrino $p_\nu^\mu = (E_\nu, p_{\nu,x}, p_{\nu,y}, p_{\nu,z})$ and an ISR photon $p_\gamma^\mu = (E_\gamma, 0, 0, p_\gamma)$). Both the neutrino and the photon have a non trivial invariant mass leading to the following equations.

Conservation of 3-momentum

$$p_x + p_{\nu,x} = 0 \quad (6)$$

$$p_y + p_{\nu,y} = 0 \quad (7)$$

$$p_z + p_{\nu,z} + p_\gamma = 0 \quad (8)$$

$$(9)$$

Conservation of Energy

$$E + E_\nu + E_\gamma = 500 \quad (10)$$

Energy-Momentum equations

$$E_\nu^2 = p_\nu^2 + m_\nu^2 \quad (11)$$

$$E_\gamma^2 = p_\gamma^2 + m_\gamma^2 \quad (12)$$

$$(13)$$

For a unique solution of E_γ we require 2 more constraints on m_γ and m_ν which have yet to be imposed, but assuming these constraints are independent of E_γ we can arrive at a solution as follows.

From conservation of 3-momentum

$$p_\nu^2 = p_{\nu,x}^2 + p_{\nu,y}^2 + p_{\nu,z}^2 \quad (14)$$

$$= p_x^2 + p_y^2 + (p_\gamma + p_z)^2 \quad (15)$$

$$= p^2 + p_\gamma^2 + 2p_\gamma p_z \quad (16)$$

$$= p^2 + E_\gamma^2 - m_\gamma^2 + 2p_\gamma p_z. \quad (17)$$

Conservation of energy then gives us,

$$(500 - E)^2 - p^2 = (E_\nu + E_\gamma)^2 - p^2 \quad (18)$$

$$= E_\nu^2 + E_\gamma^2 - p^2 + 2E_\gamma E_\nu \quad (19)$$

$$= p_\nu^2 + m_\nu^2 + E_\gamma^2 - p^2 + 2E_\gamma E_\nu. \quad (20)$$

Substituting in the expression for p_ν ,

$$(500 - E)^2 - p^2 = \cancel{p^2} + E_\gamma^2 - m_\gamma^2 + 2p_\gamma p_z + m_\nu^2 + E_\gamma^2 - \cancel{p^2} + 2E_\gamma E_\nu \quad (21)$$

$$(500 - E)^2 - p^2 + m_\gamma^2 - m_\nu^2 = 2(E_\gamma^2 + p_\gamma p_z + E_\gamma E_\nu) \quad (22)$$

$$= 2(\cancel{E_\gamma^2} + p_\gamma p_z + E_\gamma[500 - \cancel{E_\gamma} - E]) \quad (23)$$

$$= 2(p_\gamma p_z + 500E_\gamma - EE_\gamma). \quad (24)$$

292 Where we have one again used conservation of energy.

293

294 For convenience lets define

$$\lambda = \frac{1}{2}[(500 - E)^2 - p^2 + m_\gamma^2 - m_\nu^2] \quad (25)$$

and use $E_\gamma^2 = p_\gamma^2 + m_\gamma^2$ to arrive at a solvable equation in E_γ .

$$\lambda = p_\gamma p_z + 500E_\gamma - EE_\gamma \quad (26)$$

$$[\lambda - (500 - E)E_\gamma] = p_\gamma p_z \quad (27)$$

$$[\lambda - (500 - E)E_\gamma]^2 = (E_\gamma^2 - m_\gamma^2)p_z^2 \quad (28)$$

$$\lambda^2 - 2\lambda(500 - E)E_\gamma + (500 - E)^2 E_\gamma^2 = p_z^2 E_\gamma^2 - p_z^2 m_\gamma^2 \quad (29)$$

$$[(500 - E)^2 - p_z^2]E_\gamma^2 - 2\lambda(500 - E)E_\gamma + (\lambda^2 + p_z^2 m_\gamma^2) = 0 \quad (30)$$

This can be solved with the quadratic formula to give,

$$E_\gamma = \frac{\lambda(500 - E) \pm \sqrt{\lambda^2(500 - E)^2 - [(500 - E)^2 - p_z^2][\lambda^2 + p_z^2 m_\gamma^2]}}{(500 - E)^2 - p_z^2} \quad (31)$$

$$= \frac{\lambda(500 - E) \pm p_z \sqrt{\lambda^2 - [(500 - E)^2 - p_z^2]m_\gamma^2}}{(500 - E)^2 - p_z^2}. \quad (32)$$

As expected, the solution with $m_\gamma = m_\nu = 0$ reduces to the previously calculated solution

$$\lambda = \frac{1}{2}[(500 - E)^2 - p^2 + \cancel{m_\gamma^2}^0 - \cancel{m_\nu^2}^0] \quad (33)$$

$$E_\gamma = \frac{\lambda(500 - E) \pm p_z \sqrt{\lambda^2 - [(500 - E)^2 - p_z^2]\cancel{m_\gamma^2}^0}}{(500 - E)^2 - p_z^2} \quad (34)$$

$$= \frac{\lambda[(500 - E) \pm p_z]}{(500 - E)^2 - p_z^2} \quad (35)$$

$$= \frac{\frac{1}{2}[(500 - E)^2 - p^2]}{(500 - E) \mp p_z} \quad (36)$$

$$= \frac{(500 - E)^2 - p^2}{1000 - 2E \mp 2p_z}. \quad (37)$$

295 c.f Ivan's result [**IvanMarchesini**]

296 It can also easily be shown for this case that the two solutions correspond to ISR photons
 297 travelling parallel or anti-parallel to the z axis. The \mp in the denominator hence corresponds
 298 to the sign of the photons z momentum,

$$E_\gamma = \frac{(500 - E)^2 - p^2}{1000 - 2E + 2\text{sgn}(p_\gamma)p_z} \quad (38)$$

299

$$E_\gamma = \frac{\lambda(500 - E) + \text{sgn}(p_\gamma)p_z\sqrt{\lambda^2 - [(500 - E)^2 - p_z^2]m_\gamma^2}}{(500 - E)^2 - p_z^2}. \quad (39)$$