

Hiding in the trees: Unmasking a cryptic genus of arboreal New Guinea frogs with phylogenetic taxonomy

Allison R Fisher, Ke Cao, Allen Allison, Bulisa Iova, and Marguerite A Butler

2026-06-26

Abstract

Although cryptic species are known to underestimate biodiversity, cryptic genera may present greater challenges for evolutionary studies. Surprisingly there remain long-standing groups that contain unrecognized polyphyly because they are morphologically undiagnosable. A good example is the large and well-studied frog genus Oreophryne that has stood for 130 years and whose monophyly was never questioned. Recently, our group has shown it to be composed of two non-sister monophyletic clades using dense taxonomic sampling across the subfamily Asterophryinae. Here we show that the two arboreal clades are morphologically undiagnosable and define the clades using phylogenetic taxonomy. We include the type species O. senckenbergiana, resolving which clade retains the name Oreophryne, and name the strikingly similar genus **Auparoparo gen. nov.** from the indigenous Papuan language Motu. Their geographic distributions implicate plate tectonic activity in diversification, with Oreophryne distributed from the Philippines to Indonesia, along central to south New Guinea and its satellite islands, whereas Auparoparo occurs in Sulawesi and southern New Guinea and satellite islands where there is range overlap. As evidenced by this case, groups with unrecognized polyphyly are still in use today, so we urge careful confirmation of monophyly for a nomenclature based on evolution that will promote sound evolutionary research.

Introduction

Oreophryne (Boettger, 1895) is the oldest and arguably the best studied genus of the hyperdiverse subfamily Asterophryinae. Well known for its arboreal morphology and lifestyle, the taxonomy of the genus Oreophryne as currently recognized has been stable over the past century. Recently it was discovered that the single large clade is, in fact, two independently evolved clades (reviewed in Hill *et al.*, 2022). A natural question is how did this happen and why did it remain undiscovered for so long? One obvious answer is that their morphology is remarkably similar (Figure 1). Oreophryne are conspicuously arboreal and account for nearly all arboreal species of Asterophryinae (Günther, Richards, & Iskandar, 2001; Menzies, 2006), inhabiting lowland to montane rain forest, although there is a small group of diminutive, high-elevation, grassland species, a few of which are terrestrial (Zweifel, Cogger, & Richards, 2005; Günther & Richards, 2012; Putri *et al.*, 2023). In short, Oreophryne are morphologically distinctive, consistent with their arboreal lifestyle, and appear in the field to be a natural group preserved by niche conservatism.

Although taxonomic instability is well known in the history of the subfamily Asterophryinae (reviewed in Rivera *et al.*, 2017; Hill *et al.*, 2022), it appeared that Oreophryne was united by a useful diagnostic character distinct from all other genera. Based on a pectoral anatomy with highly reduced and distinctly curved clavicles (Parker, 1934; Zweifel, 2003), hypothetical new species could be assigned to Oreophryne, and for modern studies, confirmed by phylogenetic analysis with other members of Oreophryne. Over the past 130 years the genus Oreophryne has grown to 75 currently recognized species, fully 1/5 of the total diversity of subfamily

Asterophryinae and the largest of the ~16 genera (Hill et al., 2022; Frost, 2024). Oreophryne as currently recognized has the largest geographic distribution in the subfamily (Hill et al., 2023a), with species centered on mainland New Guinea and its offshore islands (Werner, 1898; Zweifel, 1956; Zweifel, Menzies, & Price, 2003; Kraus & Allison, 2004; Kraus & Allison, 2009; Kraus, 2013; Frost, 2024), but ranging westward to distant oceanic islands (Müller, 1894; Boulenger, 1896; Ahl, 1933; Putri et al., 2023) and northward to Indonesia (Boettger, 1895; Setiadi et al., 2010; Günther, Iskandar, & Richards, 2023) and the Southern Philippines Islands (Alcala, 1986; Balinton & Seña, 2015).

The apparently reliable diagnostic character, however, has never been demonstrated to be a synapomorphy, and recent studies have not found evidence for monophyly (Zweifel et al., 2005; Rivera et al., 2017). Recently, Hill et al. (2022, 2023b) established that Oreophryne sensu lato represents two reciprocally monophyletic groups that are not sister taxa (Figure 2), temporarily labeled “Oreophryne A” and “Oreophryne B”. Furthermore, it is unknown whether it is possible to identify these groups with any morphological characters – there may in fact be no reliable diagnostic characters, rendering phylogenetic taxonomic definitions for each clade an imperative for evolutionary study. The aims of this study are to assess the phylogenetic position of the newly available type species of Oreophryne to name and describe the new genus, and to develop phylogenetic definitions for both clades in the context of the subfamily-level Asterophryinae phylogeny including representatives from all known genera. Furthermore, we analyze the morphological characteristics to assess whether morphological diagnosis is possible, and consider whether there are any additional features to assist collectors when identifying

putative Oreophryne in the field (e.g., call type, geography, other characters), and discuss the implications of having misleading diagnostic characters for evolutionary studies. Inclusion of samples in molecular phylogenetics is not enough, and indeed it is necessary to include explicit tests of monophyly when proposing a new taxonomy.

Taxonomic History:

Boettger (1895) described the genus Oreophryne and the species Oreophryne senckenbergiana on the basis of three specimens from Halmahera Island, in the Maluku Archipelago west of New Guinea, that lacked procoracoids and had cartilaginous sterna. He also noted the presence of expanded finger pads and transverse folds of skin in the palate. Parker (1934) noted that procoracoids were present, but were reduced and re-diagnosed the genus on the basis of this feature together with the presence of reduced clavicles, no omosternum, and a large cartilaginous sternum. Zweifel (2003) noted that “the clavicles are tiny, slightly curved bones lying on (ventral to) the procoracoid cartilage and apart from the midline and the scapula.” On the basis of these features there are now 75 recognized species in the genus (Frost, 2024).

We note that O. senckenbergiana was synonymized (Parker, 1934) with Microhyla achatina var. moluccensis described by Peters & Doria (1878) from Bacan and Ternate islands, based on morphology and the taxon is currently known as O. moluccensis. While Parker's (1934) methodology was standard at the time, genetic information has revealed that cryptic species are rampant in this group, with species readily separated by distance and oceanic barriers

(Hill *et al.*, 2022, 2023b). Because *O. senckenbergiana* was described from Halmahera island, and Bacan and Ternate are smaller offshore islands of Halmahera, it is likely that they are different species that should be confirmed with molecular evidence.

While *Oreophryne* sensu lato is morphologically similar (Figure 1), large scale molecular phylogenetic analysis with many ingroup samples were required to clarify that this group contains two reciprocally monophyletic clades that are not sister taxa (Hill *et al.*, 2022, 2023b). Early molecular phylogenetic studies suggested monophyly but only included a few taxa within a larger-scale study (Köhler & Günther, 2008 included 9 *Oreophryne* species; De Sa *et al.*, 2012 included 4). Rivera *et al.* (2017) provided the first molecular evidence of a polyphyletic *Oreophryne* with 36 *Oreophryne* species within a larger phylogeny Asterophryinae which included 155 samples, followed by Tu *et al.* (2018)’s sparse supermatrix approach with 18 species of *Oreophryne* falling out in various places within a much larger phylogeny, but both of these lacked strong nodal support connecting the hypothesized clades to the backbone of the phylogeny. The polyphyletic nature of *Oreophryne* sensu lato was not clearly demonstrated until Hill *et al.* (2022, 2023b) included 44 species of *Oreophryne* (with data 96% complete for sampled loci) within a 218 taxon phylogeny of Asterophryinae. This phylogeny is the most complete and robust phylogeny to date. The two clades, labeled “*Oreophryne* A” and “*Oreophryne* B” have deep histories of ~ 18MY and ~16MY, respectively, with “*Oreophryne* A” sister to the rest of the subfamily (2022, 2023b).




Figure 1: Photos of *Oreophryne* species on the left, *Auparoparo* species on the right. Both genera are aboreal with a few exceptions as noted in the text, and contain nearly all arboreal species within Asterophryinae. Comparison of finger and toe pad sizes in *Oreophryne inornata* BPBM16217 vs. *Auparoparo equus* BPBM15775, bottom row.

Materials and Methods

Taxon Sampling

Asterophryinae is an adaptive radiation with short branch lengths along the backbone separating currently recognized genera (Rivera et al., 2017; Hill et al., 2022, 2023b); we reconstruct the phylogeny of Oreophryne sensu lato within the larger context of the subfamily. We include 30 named species and 18 unnamed species of Oreophryne within a larger dataset of 234 samples of 203 species of Asterophryinae collected across New Guinea, the Philippines, and Indonesia with three outgroup species to root the phylogeny (Dyscophus antongilii, Scaphiophryne marmorata, and Platypelis grandis). Importantly, we include the type species Oreophryne senckenbergiana (Boettger, 1895) in a molecular phylogeny for the first time. The samples were collected in Halmahera, North Maluku Province, Indonesia by Iqbal Setiadi , Table 1, between Supu and Dodinga, two type localities for O. senckenbergiana (Boettger, 1895), which was later synonymized with O. moluccensis (Parker, 1934).

Table 1: Locality information for samples sequenced in this study. See Hill, et al., 2023, Table 1 for additional metadata.

			Latitude
ID	Species	Locality	Latitude, Longitude
BJE01143	<i>Oreophryne moluccensis</i>	Kabupaten Halmahera Utara, Halmahera Island, North Maluku	1.7899, 127.8910
BJE01144	<i>Oreophryne moluccensis</i>	Kabupaten Halmahera Utara, Halmahera Island, North Maluku	1.7899, 127.8910
BJE01120	<i>Oreophryne moluccensis</i>	Kabupaten Halmahera Utara, Halmahera Island, North Maluku	1.8228, 127.8300

Molecular Sequence

We obtained nearly complete DNA sequence data for five loci (three nuclear: Seventh in Absentia [SIA], Brain Derived Neurotrophic Factor [BDNF], Sodium Calcium Exchange subunit-1 [NXC-1], and two mitochondrial loci (Cytochrome oxidase b [CYTB], and NADH dehydrogenase subunit 4 [ND4] from four samples of two species not previously sequenced and combined these with previously published data (Rivera et al., 2017; Hill et al., 2022, 2023b) for a dataset of 240 samples with 207 species and 2475 base pairs that is 96% complete. All loci were

complete for the added samples. Sequence quality was evaluated using chromatogram quality scores, checks for stop codons indicating pseudogenes, BLAST searches (Ye *et al.*, 2012), and concordance of pairwise similarities with other Asterophryinae sequences. Primer sequences and detailed methods are provided in (Hill *et al.*, 2023b). We note that we excluded from the previous alignment (Hill *et al.*, 2022, 2023b) two samples of Aphantophryne pansa (BPBM 5299 and BPBM 8312) which were deemed unreliable as they came from formalin-fixed material, and three samples which fell out in odd places that should be confirmed with new material prior to recommending taxonomic revision (BPBM 22708 Copiula tyleri, BPBM 38939 Copiula sp. 7, BPBM 37753 Paedophryne dekot). We note that genetic material from 47 species of Oreophryne sensu lato was not available for this molecular analysis.

Morphology, Behavior, and Geographic Distribution

We collected new morphological characters and reexamined traits compiled from the literature, character states defined in Table 2 for all named species of Oreophryne in our phylogenetic dataset. Our sample included all species available in the collections of the Bishop Museum, Honolulu, Hawaii (BPBM): thirteen species of “Oreophryne A”, eight species “Oreophryne B” (Table 3). Finger and toe pads were photographed and length and width of the toe tips were measured using ImageJ software. Trait information for the remaining species were gathered from the literature, six species of “Oreophryne A” and one of “Oreophryne B”. In addition, we compiled information on call type from the literature and collection location from the respective collectors or the BPBM database (global positioning coordinates [GPS]; Table 3).

We examined the ability of morphological and behavioral characters to provide a diagnosis to the two Oreophryne clades consistent with molecular phylogenetics.

Table 2: Description of traits surveyed.

Column*	Trait	Definition
1	Procoracoids	Size and shape of the procoracoid cartilages, CR= curved, reduced
2	Clavicles	Size and shape of the clavicles, CR= curved, reduced
3	Omasternum	Presence (+) or absence (-) of the omasternum
4	Sternum	Cartilaginous sternum or sternal plate
5	Call type	Qualitative description of the call type
6	Finger:toe width	The ratio of finger pad width to toe pad width
7	Pectoral ligament	Cartilaginous or ligamentous connection between scapula and procoracoid
8	Rel toe length	Relative length of 5th toe compared to length of 3rd toe (5<3, 5=3, 5>3)
9	Toe webbing	Toe webbing assessed on 5th toe (absent (-), basally webbed, well webbed)
10	Tympanum:eye diameter	Tympanum diameter relative to eye diameter ($\frac{1}{4}$, $\square$, or $\frac{1}{2}$)
	Pupil horizontal	Whether the pupil is horizontal

Column*	Trait	Definition
	Teeth	Whether teeth are present
	Palatal folds	Whether two horizontal palatal folds in front of the pharynx are present

*Corresponding column in Table 3. The last three traits were scored and are invariable across species, thus not tabulated in Table 3

Nomenclature

We conducted a comprehensive taxonomic literature review of diagnoses for all species known to belong to the clades “Oreophryne A” and “Oreophryne B” (Hill et al., 2022). We applied the ICZN code (International Commission on Zoological Nomenclature, 1999) as well as the Phylocode (De Queiroz & Cantino, 2020). We support the nascent effort to assign indigenous names in taxonomy whenever possible (e.g., Gillman & Wright, 2020). One of us (B. Iova) provided guidance in proposing the novel genus name, Auparoparo, meaning “tree frog” in Motu, an indigenous language of Papua New Guinea, for Oreophryne “B”.

Phylogenetic Reconstruction

We used Partitionfinder in IQTREE2 to find the best-fit evolutionary model followed by Maximum Likelihood (ML) phylogenetic reconstruction (Lanfear et al., 2012; Minh et al., 2020). Following Hill et al. (2022), we tested models partitioned by both locus and codon versus by

loci only (15 vs. 5 partition scheme) with merging of partitions allowed. We found following merging that a 10-partition scheme involving locus and codon models with merged partitions was superior by the Akaike Information Criterion (models by partition included in Supplementary Materials). Nodal support values for the ML tree were inferred using 2000 bootstrap replicates.

Phylogenetic Definition

We use a maximum crown-clade definition under the PhyloCode (De Queiroz & Gauthier, 1990, 1992; De Queiroz & Cantino, 2020), specifically, the largest crown clade originating with the most recent common ancestor of the reference species for the clade (e.g., the type species of the genus), and all extant species that share a more recent common ancestor with that reference species than they do with reference species of other clades (i.e., in this study, type species of other genera or their proxies). This definition is robust to uncertainty in basal relationships such as in the case of adaptive radiations with short branch lengths along the backbone as in Asterophryinae, as well as being robust to identification of new species and new clades.

Analyses, Visualization, and Data Availability

All analyses were conducted and graphics produced in the R Statistical Computing Environment unless noted below (R Core Team, 2024). Initial DNA sequence alignment was conducted using MUSCLE (Edgar, 2004) with manual adjustments in Mesquite (Maddison & Maddison,

2023). Phylogenetic reconstruction was conducted using the IQTREE2 suite of software (Lanfear [et al.](#), 2012; Minh [et al.](#), 2020).

The known ranges of Oreophryne sensu lato were downloaded as shape files from the IUCN (2025) redlist database, and sites from collection locations for Oreophryne and Auparoparo were mapped using the `sf` and `ggplot2` packages (Wickham, 2016; Pebesma, 2018), overlaid on a basemap created using `rnaturalearth`, `grid`, and `ggspatial` packages (Dunnington, 2023; Massicotte & South, 2023). Tree graphics were produced with the `ggtree` package (Yu [et al.](#), 2017; Yu, 2020) supplemented by `ggplot2` (Wickham, 2016). All data, alignments, model and tree outputs, and code to reproduce analyses are available in the public repository ...

Results

Phylogeny and Phylogenetic Taxonomy

For our subfamily-level phylogeny, a partition scheme including codon as well as locus resulted in an improvement of fit by 3,127 AIC units (evolutionary model partitioned by locus and codon, 15-partitions: AIC of 155,492, versus partitioned by locus only, 5 partitions: AIC of 158,619). Unless otherwise noted, all phylogenetic reconstructions used the 15-partition scheme identified as best-fit by IQTREE2.

We recovered a maximum likelihood topology that confirms that Oreophryne sensu lato is polyphyletic, with species falling into two independently-evolved, reciprocally monophyletic, highly-supported clades that are not sister groups Figure 2. Nearly all nodes are moderately or

highly supported, with most of the weakly supported nodes falling along the backbone where there are very short branch lengths as characteristic of an adaptive radiation.

Furthermore, these findings demonstrate that the clade descended from node A should retain the name Oreophryne, as the type species, O. senckenbergiana, from North Maluku Island, Indonesia, is situated within Oreophryne with high support (Figure 2). We propose the new clade name Auparoparo for the clade descended from node B. Oreophryne is the earliest diverging genus-level clade in the Papuan subfamily, with genera Cophixalus and Paedophryne arising before Auparoparo, which split from the remaining genera of Asterophryinae.

Morphology, Behavior, and Ranked Taxonomy

We find no morphological synapomorphies that distinguish Auparoparo from Oreophryne. Oreophryne sensu lato is diagnosed by characters of the pectoral girdle: possessing highly reduced, curved procoracoid cartilages and clavicles, and lacking an omosternum (Parker, 1934; Zweifel, 2003). We found that this suite of characters is also present in Auparoparo, and therefore is not diagnostic for either Oreophryne or Auparoparo (Table 3).

Two traits imperfectly segregate between the clades, relative finger and toe pad widths and call type (Table 3), which may assist in tentative identification. Both Oreophryne and Auparoparo are arboreal clades with expanded digital pads. Species of Oreophryne have finger and toe pads generally of roughly equal width and most species of Auparoparo have visibly larger finger pads, with finger pads roughly 20-30% wider than toes. Both Oreophryne and Auparoparo use a variety of call types, with “honks” or “peeps” used only by Oreophryne, whereas “whinny” calls are used only by Auparoparo. A minority of species in both groups use the “rattle” call type. However, both toe pad width and call type are variable traits with some overlap and thus not unambiguously diagnostic. Furthermore, possession of expanded digital pads is clearly an adaptive trait for arboreality and prone to homoplasy.




Figure 2: Molecular phylogenetic hypothesis of Asterophryinae showing polyphyly of *Oreophryne* sensu lato, into two reciprocally monophyletic clades that are not sister taxa. Node A is *Oreophryne* (blue clade) and node B is *Auparoparo* (red clade) based on the position of the type species *Oreophryne senkenbergiana*=*O. moluccensis*. Nodal support: black dots indicate ML bootstrap support $\geq 95\%$, grey dots $\geq 75\%$, numeric values given for nodes below 75% support. Outgroup taxa excluded. Taxa in bold are the type species for each genus (or a proxy species used as an internal reference).

Table 3: Morphological character m

		(1)	(2)	(3)	(4)	(5)		(6)		(7)
Genus	species	proc	clav	omo	ster	call type		fing: toe width		pec lig
<i>Oreophryne</i>	<i>anamiatoi</i>	CR	CR	–	cart	chuckle or cackle		1.2		cart
	<i>anulata</i> ¹	CR	CR	–	cart			slightly larger		
	<i>aurora</i>	CR	CR	–	cart	harsh croaks				lig
	<i>biroi</i>	CR	CR	–	cart	rattle		1.1		lig
	<i>brachypus</i>	CR	CR	–	cart	long squeak		1.1		lig
	<i>cameroni</i>	CR	CR	–	cart	peep				cart
	<i>frontifasciata</i> ¹	CR	CR	–	cart			slightly larger		cart
	<i>geislerorum</i>	CR	CR	–	cart	rattle				lig
	<i>graminis</i> ¹	CR	CR	–	cart	peep		approx equal		cart
	<i>inornata</i>	CR	CR	–	cart	honk		1.0		lig
	<i>lemur</i>	CR	CR	–	cart	honk				lig

		(1)	(2)	(3)	(4)	(5)	(6)	(7)
Genus	species	proc	clav	omo	ster	call type	finger: toe width	pec lig
	<i>loriae</i>	CR	CR	—	cart	honk		lig
	<i>moluccensis</i> ¹	CR	CR	—	cart		slightly larger	
	<i>nana</i> ¹	CR	CR	—	cart		approx equal	cart
	<i>notata</i>	CR	CR	—	cart	peep		cart
	<i>parkeri</i>	CR	CR	—	cart	peep		lig
	<i>parkopanorum</i>	CR	CR	—	cart			cart
	<i>philosylleptoris</i>	CR	CR	—	cart	honk	1.0	lig
<i>Auparoparo</i> ²	<i>brunnea</i> ¹	CR	CR	—	cart	rattle	larger	lig
	<i>equus</i>	CR	CR	—	cart	whinny	1.3	lig
	<i>ezra</i>	CR	CR	—	cart	chatter		lig
	<i>insulana</i>	CR	CR	—	cart	whinny		lig
	<i>matawan</i>	CR	CR	—	cart	rattle		lig

		(1)	(2)	(3)	(4)	(5)	(6)	(7)
Genus	species	proc	clav	omo	ster	call type	finger: toe width	pec lig
	<i>meliades</i>	CR	CR	–	cart	whinny	1.2	lig
	<i>penelopeia</i>	CR	CR	–	cart	whinny	1.3	lig
	<i>phoebe</i>	CR	CR	–	cart	whinny	1.2	cart
	<i>picticus</i>	CR	CR	–	cart	whinny	1.3	lig
	<i>variabilis</i> ¹	CR	CR	–	cart		much larger	cart

¹Character states obtained from the literature, specimens not available for examination in this study.

²Hill et al. 2022 places *Aphantophryne pansa* within *Auparoparo*, but the status of *Aphantophryne* requires further study with samp

*External characters were examined from specimens, internal characters and pupil shape taken from the literature.

Geographic Distribution

Oreophryne has a wide distribution with species spanning from the southern Philippines to the Papuan Louisade Archipelago (Figure 3, localities for Oreophryne species included in our phylogeny are mapped with blue triangles). This range encompasses the additional species from the type locality in Indonesia. Auparoparo is mainly clustered in the southeastern tip

of mainland Papua New Guinea and its satellite islands with two species in Sulawesi, an Indonesian island (Figure 3, Auparoparo localities mapped in red circles). We also mapped Oreophryne sensu lato ranges obtained from the IUCN Red List (IUCN, 2025), which require phylogenetic diagnosis. Note the Indonesian region of New Guinea is difficult to attain samples from and it is unknown whether these groups inhabit this connective area.




Figure 3: The known distribution of Oreophryne as currently recognized is plotted in grey polygons (data source IUCN, 2025), with points indicating Auparoparo (red) and Oreophryne (blue) as assigned by our phylogenetic analysis Figure 2. Note that there is range overlap along the East Papuan Composite Terrane and on some offshore islands (see inset).

Taxonomy

We use a maximum crown-clade definitions for designating Oreophryne and Auparoparo because the monophyly of these taxa are strongly supported, and the relationships of these taxa to each other and to other genera of Asterophryinae are well-supported (this study, Hill et al., 2022, 2023b), as has been done for other anuran taxa (Brown et al., 2015) following the methods of phylogenetic nomenclature (De Queiroz & Gauthier, 1990, 1992, 1994). That is, we adopt phylogenetic definitions of these genera that explicitly identify them as clades. Specifically, we define each genus to include all descendants of the most recent common ancestor of the type species and all other extant species that are more closely related to it than to the type species of the other Asterophryine genera recognized here. We retain the name Oreophryne for species included in the phylogenetic redefinition of Oreophryne, and new clade name Auparoparo for newly designated species of Auparoparo. The previously defined polyphyletic Oreophryne is indicated by “sensu lato”, which contains many species which require molecular phylogenetic assignment.

Family Microhylidae Günther 1858

Subfamily Asterophryinae Günther 1858

Genus Oreophryne Boettger 1895 amended

Microhyla Peters 1878

Callula (Horst 1883)

Oreophryne Boettger 1895, by monotypy, Type species: Oreophryne senckenbergiana Boettger 1895. (= Microhyla achatina var. moluccensis Peters 1878) Placed on Official List of Generic

Names in Zoology by Opinion 1266, Anonymous, 1984.

Sphenophryne in part (Méhely 1897)

Phrynixalus (Stejneger 1908)

Mehelyia Wandolleck 1911, Type species: Meheylia lineata Wandolleck 1911 (= Sphenophryne biroi Mehely 1897, by subsequent designation of Parker 1934. Synonymy by Nieden (1926).

Cophixalus in part (Boettger 1892)

Chaperina in part (Taylor 1920)

Diagnosis: We note that there are no synapomorphies in adult Oreophryne that differ from the morphologically similar genus Auparoparo. Diagnoses should be made by molecular phylogenetic analysis, based on the phylogenetic definition below. Oreophryne differs from most other genera in subfamily Asterophryinae, but not from Auparoparo, by the combination of (1) presence of highly reduced clavicles (vs absence or elongate) which have a distinctly curved shape; (2) presence of well developed digital pads, often with finger and toe pads of similar size; (3) arboreal lifestyle (perch height >2m above ground); (4) most species range in body size from 16mm (O. graminis, Günther & Richards, 2012) to 30mm, with a few larger species ~40mm (O. inornata); (5) small to invisible tympanum; (6) large eyes with horizontal pupils; (7) occasional presence of toe webbing; (8) thin, elongated limbs; and (9) in many species males that call with “honk” or “peep” types or in some species “rattle” type. Some species of Oreophryne may be distinguished from Auparoparo by roughly equal finger to toe pad ratio and males calling with “honk” or “peep” but not “whinny” call types.

Phylogenetic definition: Oreophryne Boettger 1895 (Converted Clade Name) is defined as

the largest crown clade including Oreophryne senckenbergiana Boettger 1895 and excluding the type species of the other genus-ranked clades listed in Table Table 4.

Table 4: Genus-level clades of Asterophryinae and their internal specifiers (type species or proxies).

Genus-level clade	Internal Specifier (Type Species or proxy)
<i>Asterophrys</i>	<i>Asterophrys turpicola</i> (Schlegel 1837)
<i>Auparoparo</i>	<i>Auparoparo penelopeia</i> (Kraus 2016)
<i>Austrochaperina</i> *	<i>Austrochaperina palmipes</i> (Zweifel 1956)
<i>Austrochaperina</i> *	<i>Austrochaperina robusta</i> Fry 1912
<i>Barygenys</i>	<i>Barygenys cheesmanae</i> Parker 1936
<i>Callulops</i>	<i>Callulops doriae</i> Boulenger 1888
<i>Choerophryne</i>	<i>Choerophryne proboscidea</i> Van Kampen 1914
<i>Cophixalus</i>	<i>Cophixalus verrucosus</i> (Boulenger, 1898)
<i>Copiula</i>	<i>Copiula oxyrhina</i> (Boulenger 1898)
<i>Genyophryne</i> *	<i>Genyophryne thomsoni</i> Boulenger, 1890

Genus-level clade	Internal Specifier (Type Species or proxy)
<i>Hylophorbus</i>	<i>Hylophorbus rufescens</i> Macleay 1878
<i>Mantophryne</i>	<i>Mantophryne lateralis</i> Boulenger 1897
<i>Liophryne</i> *	<i>Liophryne rhododactyla</i> (Boulenger 1897)
<i>Oreophryne</i>	<i>Oreophryne senckenbergiana</i> Boettger 1895 (= <i>O. moluccensis</i> (Peters and Doria 1878))
<i>Oxydactyla</i>	<i>Oxydactyla brevicrus</i> (Van Kampen 1913)
<i>Paedophryne</i>	<i>Paedophryne kathismaphlox</i> Kraus 2010
<i>Sphenophryne</i> *	<i>Sphenophryne cornuta</i> Peters and Doria 1878
<i>Xenorhina</i>	<i>Xenorhina oxycephala</i> (Schlegel 1858)

*We note that the monophyly of *Austrochaperina*, *Liophryne*, *Oxydactyla*, and *Sphenophryne* (and reciprocal monophyly of *Genyophryne* [*Genyophryne thomsoni* Boulenger, 1890] with respect to *Liophryne* + *Oxydactyla* + *Sphenophryne*) has not yet been established. *Austrochaperina* is polyphyletic in recent phylogenies with possibly three monophyletic clades that are not sister taxa (Figure @fig-1).

Composition: Our results support the retention of 17 species in *Oreophryne* with molecular evidence (Table 5): *Oreophryne anamiatoi* Kraus 2009, *O. anulata* (Stejneger 1908), *O. aurora* Kraus 2016, *O. biroi* Mehely 1897, *O. brachypus* Werner 1898, *O. cameroni* Kraus

2013, O. geislerorum Boettger 1892, O. graminis Gunther 2012, O. inornata Zweifel 1956, O. lemur Kraus 2016, O. lorae Boulenger 1898, O. nana Brown 1967, O. notata Zweifel 2003, O. parkeri Loveridge 1955, O. parkoporum Kraus 2013, O. philosylleptoris Kraus 2016, and O. moluccensis (Peters 1878). A number of species were not available for our study but should molecular evidence demonstrate their relationship within Oreophryne, those species should also retain their generic identity.

Distribution: The range of confirmed Oreophryne species in this dataset remains nearly as broad as previously defined (Figure 3), from the Philippines and Indonesia to the southeastern most tip of mainland New Guinea and several near-shore islands (Fergusson, and Normanby Is.) and the distant Rossel Island of the Louisiade Archipelago. However, the transfer of species which occupy Woodlark Islands and Misima and Sudest Islands of the Louisiade Archipelago reduces the known range of Oreophryne. Woodlark Island is inhabited by O. phoebe [= Auparoparo phoebe], Misima Island is inhabited by O. picticus [= Auparoparo picticus], and Sudest Island is inhabited by O. ezra and O. meliades [= Auparoparo ezra and Auparoparo meliades].

Genus Auparoparo gen. nov.

Oreophryne in part (Boettger, 1895)

Sphenophryne in part (Boulenger, 1897)

Type species Oreophryne penelopeia (Kraus, 2016) [= Auparoparo penelopeia]

Diagnosis: Currently, there are no known synapomorphies in adult Auparoparo that differ from the morphologically similar genus Oreophryne. Diagnoses should be made by molecular

phylogenetic analysis, based on the phylogenetic definition below. Auparoparo differs from most other genera in subfamily Asterophryinae, but not from Oreophryne, by the combination of (1) presence of highly reduced clavicles (vs absence or elongate) which have a distinctly curved shape; (2) presence of well developed digital discs, with finger discs notably wider than toe discs (typically 20% or more); (3) arboreal lifestyle (perch height >2m above ground); (4) small to moderately large body size in known species, from 16mm (A. brunnea Kraus 2017) to 25 mm (A. picticus Kraus 2016); (5) small to invisible tympanum; (6) large eyes with horizontal pupils; (7) occasional presence of toe webbing; (8) and thin, elongated limbs; and (9) and in many species males that call with a “whinny” call type or in some species “rattle” type. Some species of Auparoparo may be distinguished from Oreophryne by larger finger pads than toe pads and males calling with “whinny” but not “honk” or “peep” call types.

Phylogenetic definition: Auparoparo (New Clade Name) applies to the largest crown clade including Auparoparo penelopeia (Kraus 2016) and excluding the type species of the other genus-ranked clades listed in Table 4.

Composition: We transfer eleven named species from Oreophryne into Auparoparo (Table 5): Auparoparo brunnea (Kraus, 2017), A. equus (Kraus, 2016), A. ezra (Kraus & Allison, 2009), A. insulana (Zweifel, 1956), A. matawan (Kraus, 2016), A. meliades (Kraus, 2016), A. penelopeia (Kraus, 2016), A. phoebe (Kraus, 2017), A. picticus (Kraus, 2016), A. penelopeia (Kraus, 2016), A. variabilis (Boulenger, 1896).

Distribution: Southeastern tip of mainland Papua New Guinea and satellite islands (Fergusson, Misima, Normanby, Rossel, Sudest, Woodlark Is.), Sulawesi.

Etymology: The generic name Auparoparo comes from the indigenous language, Motu, which is widely spoken in the region of Papua New Guinea in which this genus is found. It is a combination of the words “au” (tree) and “paro paro” (frog). Motu is not a gendered language, however, for nomenclatural purposes we designate the genus Auparoparo as feminine, thereby preserving the gender of existing species names.

Discussion

Our phylogenetic results (Hill *et al.*, 2022; 2023b, this study) provide clarity on the classification of the problematic genus Oreophryne sensu lato. We confirm that Oreophryne sensu lato is polyphyletic and consists of two reciprocally monophyletic clades that are not sister taxa (Figure 2). Here we designate “Oreophryne A” in Hill *et al.* (2022) as the genus Oreophryne because it contains the type species Oreophryne senckenbergiana Boettger 1895 with high support. Oreophryne is the earliest diverged genus in subfamily Asterophryinae originating approximately 18 MYA (Hill *et al.*, 2022), and is sister to a clade composed of all the other genera. The second clade, “Oreophryne B” in Hill *et al.* (2022), is highly supported as a separate, non-sister genus, which originated approximately 16 MYA and contains no known species that are the types of any previously proposed genera. Thus, we designate this genus as Auparoparo, a tribute to the Motu language of the indigenous peoples of the region.

The Difficulty with Cryptic Genera: Can we tell them apart?

Perhaps the greatest testament to the crypticism of Oreophryne vs. Auparoparo is that no taxonomist was able to distinguish that they are separate clades, independently evolved, in the 130 years between their discovery and the present classification [Boettger (1895); this study]. These species are indeed morphologically very similar (Figure 1) and our results provide some clues as to why Oreophryne taxonomy has been so difficult. We found morphological synapomorphies for neither Auparoparo nor Oreophryne, and moreover, no morphological features that can diagnose either clade.

The traditional basis for assigning species to Oreophryne has primarily relied on a signature pectoral anatomy (whether original or revised diagnosis, Parker, 1934; Zweifel, 2003). Our phylogenetic results clearly show that this trait appears more than once, in both Oreophryne and Auparoparo, is either a symplesiomorphy (shared ancestral character) or a result of convergent or parallel evolution, and perhaps adaptive for arboreality. Thus, the pectoral anatomy is not a reliable diagnostic character in differentiating between Oreophryne and Auparoparo. In studies across a wide diversity of anurans, Soliz et al. (2025) found that anatomical variation of the pectoral girdle corresponds with functional and ecological lifestyle demands, whereas Engelkes et al. (2020) found strong phylogenetic influence in the shape of the pectoral girdle. The pectoral girdle would be excellent for further study within Asterophryinae, as there are high elevation terrestrial species of Oreophryne sensu lato that possess the same pectoral trait (Zweifel et al., 2005), as well as a few arboreal species within other genera (Cophixalus, Asterophrys, Xenorhina) with unknown pectoral morphology.

We further reexamined additional characters previously used to diagnose Oreophryne as well as characters conventionally used to diagnose species of frogs. We conclude that pectoral girdle connector type (Parker, 1934) and relative lengths of the 3rd and 5th toe are uninformative and the degree of toe webbing is variable. It was previously suggested that species in Oreophryne sensu lato could be divided into two groups based on possessing a cartilaginous or ligamentous connection between the procoracoid cartilages and the scapula (Kraus, 2013), with roughly half possessing each type (Parker, 1934). However, we found that cartilaginous and ligamentous connectors are common across both genera (Table 3); therefore, these traits are not synapmorphies for neither Oreophryne nor Auparoparo and do not indicate monophyly for these clades. Of species with toe webbing, Oreophryne tends to be described as having “well webbed” toes compared to Auparoparo’s “basal” webbing (Table 3). However, the presence of intra-specific variability combined with subjectivity in judging relative landmarks (e.g., “reaching the penultimate tubercle”, “webbed 1/3 length of 5th toe”, etc.), makes this feature difficult to standardize for generic diagnostic use. Many of the morphological traits proposed by Parker (1934) and Zweifel (2003) may in fact be a result of homoplasy via convergent lifestyle or parallel evolution, and while they may prove useful for ecomorphological identification or species diagnoses in combination with other traits, they are not informative for generic diagnosis.

We found only two characters with potential for imperfectly distinguishing the genera: relative digital pad size (Parker, 1934; Zweifel, 1956), and call type (e.g., Kraus, 2016). The relative widths of the digit pads differs between genera: Auparoparo tends to have visibly larger finger

pads in comparison to toe pads, whereas in Oreophryne the finger and toe pads are roughly equal in size (Figure 1, bottom row), although there is some overlap in finger to toe pad ratio (Table 3). In some cases, Auparoparo's finger pads display T-shapes while the toes are often rounder and narrower. Whether there is any performance advantage in larger finger pads compared to toes is unknown. Furthermore, a handful of high-elevation Oreophryne species not included in our phylogeny and yet to be diagnosed by molecular phylogenetic analysis have evolved terrestrial or fossorial habits and possess smaller digital pads compared to their presumed arboreal relatives (Zweifel et al., 2005).

Call features are increasingly used for diagnosing species of Oreophryne, with advertisement call types commonly categorized as rattle, honk, whinny, peeping, or clicking (Kraus, 2016). We found honk or peep call types are only found in Oreophryne species in our sample, while whinny calls are only found in Auparoparo species, and rattle calls are found in both genera (Table 3). We note that several authors described species of both genera, thus the difference between genera cannot be attributed to the investigator. However, more complete sampling may add further complexity. As far as we are aware there has been no comprehensive acoustic analysis across Oreophryne sensu lato. More research is needed to explore call diversity both between species and across genera to understand call evolution and whether there any features that are distinctive of genera.

Geographic Distribution

Both Oreophryne and Auparoparo are geographically centered in the mainland New Guinea. Oreophryne extends southward to the Louisiade Archipelago and northward to the South Philippine Islands (Hill *et al.*, 2023b). Addition of O. moluccensis expands the known range eastward to North Maluku, Indonesia (Figure 3).

The known distribution of Auparoparo is mainly clustered in the southeastern tip of mainland Papua New Guinea (in the East Papuan Composite Terrane) and satellite islands with two species in Sulawesi, Indonesia. This suggests the possibility of interesting plate tectonics resulting in this biogeographic distribution. Exploration of intermediate localities may result in as yet undescribed or undiscovered species of Auparoparo. The geographic range is sure to grow with additional phylogenetic sampling. For example Oreophryne (sensu lato) chlorops (Günther *et al.*, 2023) is a very large (SVL > 40mm) species recently discovered from mid-elevation mountains of the bird s neck region of Western Papua with clearly enlarged finger pads relative to toe pads. If confirmed as Auparoparo, it would significantly expand the range of Auparoparo across New Guinea Island.

We found geographic range overlap between Oreophryne and Auparoparo at multiple localities: on the mainland along the Mt. Dayman Foothills, at the Mt. Simpson Massif, Mt. Trafalgar, and on the oceanic islands of Normanby and Fergusson (both nearshore islands), and Rossel island (hundreds of kilometers offshore), which presents interesting questions regarding biogeography as well as their ecological relationships. For example, Oreophryne inornata and

Auparoparo insulana have been collected at the same site (Goodenough Island) and even the same microhabitats (Zweifel, 1956). Do these species occupy the same niche, and what is the temporal history of each colonization? Are both genera also present in the nearby islands of Normanby and Fergusson (all part of the D'Entrecasteaux Archipelago)? Obtaining additional geographical samples for inclusion in phylogenetic analysis should be a high priority to gain a more complete understanding of the geographic distribution and evolutionary history of both Oreophryne and Auparoparo.

How to Avoid a Misleading Classification

We adopt an explicitly phylogenetic framework for the new taxonomy of the genera Oreophryne and Auparoparo, and to our knowledge, this is the first time that phylogenetic nomenclature has been applied within the subfamily Asterophryinae. A phylogenetic definition of genera is advantageous especially because Asterophryinae is an adaptive radiation. While we are increasingly confident that at least 16 genera are indeed monophyletic, nevertheless, very short branch lengths separate many of the genera along the backbone [Hill et al. (2022); this study]. A maximum-crown-clade definition with multiple external specifiers is robust to uncertainty (i.e., rearrangements of order) along the backbone. We define each genus as the largest crown clade including the internal specifier species for the clade (i.e., the type species of the genus) but not the internal specifier species of other asterophryine genera. For long-studied groups such as Asterophryinae, it may not be possible to obtain a genetic sample for type species that were collected in the 1800 s and early 1900 s, as these locations may be uncertain and/or

remote, it may be necessary to use a genetic sample from another species of the genus as a proxy for the type species. Nevertheless, it is increasingly clear that for Oreophryne and Auparoparo, a molecular phylogenetic approach is required for final confirmation of taxonomy.

Furthermore, a phylogenetic definition provides guidance for robust classification. It is imperative that when assigning new species to either Oreophryne or Auparoparo, molecular confirmation requires – at minimum – inclusion of members of the other clade to correctly identify the genus level clade. For example, a phylogenetic analysis with only Oreophryne sensu lato and outgroups to Asterophryinae will show incorrect grouping of Oreophryne and Auparoparo species because of the lack of intervening species assigned to other Asterophryinae genera. An additional difficulty here is that each clade, Oreophryne and Auparoparo, contains subclades with deep histories (Hill *et al.*, 2023a), so we recommend the inclusion of additional intervening species assigned to other Asterophryinae genera, particularly Cophixalus and Paedophryne, to reduce the potential for long-branch attraction and more clearly resolve the “boundaries” of each monophyletic clade.

This study furthermore highlights the importance of taxonomic sampling for robust phylogenetic results. Developing an orderly taxonomy in a clade with over 350 named species (with perhaps as much as 40% of species in museum collections that are yet to be described, A. Allison, unpublished) would be that much more challenging without the benefit of generic diagnoses that follow evolutionary history.

Why the Names Matter

Clarifying the generic structure also facilitates appreciation for the patterns of diversification within Asterophryinae. The evolutionary history of this subfamily is characterized by several interesting trends across genera: 1) a burst of ecological and species diversification leading to generic-level clades occurring early in the history of the subfamily (Hill et al., 2022), 2) a tendency toward niche conservatism within genera, and 3) simultaneous dispersal of multiple genera to offshore islands (Hill et al., 2022, 2023a) revealing the very strong influence of plate tectonics for explaining the historical dispersal patterns of this group. Finally, without distinct generic names for the two clades represented by Oreophyne and Auparoparo, it becomes much more difficult to appreciate independently evolutionary histories embodied by each clade, that aboreality has evolved at least twice in subfamily Asterophryinae, and would make it very difficult to appreciate the impressive parallelism embodied by these speciose and successful genera. An accurate and informative taxonomy serves to advance our understanding of Papuan microhylids and their very interesting evolutionary history leading to further discovery.

Lastly, we emphasize that future naming of species, genera, and higher taxa should seek to recognize the language of the indigenous stewards of the land and ecosystem from which they were discovered (Gillman & Wright, 2020). This is especially true of Asterophryinae frogs, whose awe-inspiring diversity mirrors the rich cultural beauty and historical complexity of Papuan people.

Acknowledgements

We thank Ben Evans and Iqbal Setiadi for generously providing samples critical for this study, and Ethan Hill for his guidance and training on the molecular and bioinformatics work. We are grateful to Molly Hagemann for her excellent assistance in accessing Bishop Museum Vertebrate Zoology Collections and data. We thank Neal Evenhuis for sharing expert knowledge on type species, and Brianna Correa and Alexis Shulga for providing comments which improved the manuscript. This is publication 234 from the School of Life Sciences, University of Hawai'i at Mānoa.

Funding

This study was made possible by a grant from the National Science Foundation awarded to MB (DEB-1145733). We gratefully acknowledge support from the School of Life Sciences, University of Hawai'i at Mānoa. University of Hawaii Information Technology Services – Cyberinfrastructure resources were funded in part by a National Science Foundation Major Research Instrumentation award (CNS-1920304).

Data Availability

The data underlying this article are available in the GenBank Nucleotide Database] at <https://www.ncbi.nlm.nih.gov/genbank/> , and can be accessed with [will be provided upon

publication]. The codes and datafiles needed to reproduce these results can be found at [GitHub repository] with DOI [supplied upon publication].

Supplementary Table

Table 5: Classification of Auparoparo and Oreophryne based on phylogenetic analysis of two mitochondrial gene fragments (CytB, ND4) and three nuclear gene fragments (RAG1, RAG2, NXC). Incertae sedis taxa require phylogenetic evidence for

Original designation	Citation	Previous generic placement	Current placement
<i>Oreophryne anamiatoi</i>	Kraus & Allison, 2009	<i>Oreophryne</i>	<i>Oreophryne</i>
<i>Phrynixalus anulatus</i>	Stejneger, 1908	<i>Oreophryne</i>	<i>Oreophryne</i>
<i>Oreophryne aurora</i>	Kraus, 2016	<i>Oreophryne</i>	<i>Oreophryne</i>
<i>Sphenophryne biroi</i>	Méhely, 1897	<i>Oreophryne</i>	<i>Oreophryne</i>
<i>Hylella brachypus</i>	Werner, 1898	<i>Oreophryne</i>	<i>Oreophryne</i>
<i>Oreophryne cameroni</i>	Kraus, 2013	<i>Oreophryne</i>	<i>Oreophryne</i>
<i>Cophixalus geislerorum</i>	Boettger, 1892	<i>Oreophryne</i>	<i>Oreophryne</i>

Original designation	Citation	Previous generic placement	Cu pla
<i>Oreophryne graminis</i>	Günther & Richards, 2012	<i>Oreophryne</i>	Or
<i>Oreophryne inornata</i>	Zweifel, 1956	<i>Oreophryne</i>	Or
<i>Oreophryne lemur</i>	Kraus, 2016	<i>Oreophryne</i>	Or
<i>Sphenophryne loriae</i>	Boulenger, 1898	<i>Oreophryne</i>	Or
<i>Microhyla achatina</i>	var. moluccensis Peters and Doria, 1878	<i>Oreophryne</i>	Or
<i>Oreophryne nana</i>	Brown & Alcala, 1967	<i>Oreophryne</i>	Or
<i>Oreophryne notata</i>	Zweifel, 2003	<i>Oreophryne</i>	Or
<i>Oreophryne parkeri</i>	Loveridge, 1955	<i>Oreophryne</i>	Or
<i>Oreophryne parkoporum</i>	Kraus, 2013	<i>Oreophryne</i>	Or
<i>Oreophryne philosylleptoris</i>	Kraus, 2016	<i>Oreophryne</i>	Or
<i>Oreophryne brunnea</i>	Kraus, 2017	<i>Oreophryne</i>	Au
<i>Oreophryne equus</i>	Kraus, 2016	<i>Oreophryne</i>	Au

Original designation	Citation	Previous generic placement	Cu pla
<i>Oreophryne ezra</i>	Kraus & Allison, 2009	<i>Oreophryne</i>	Au
<i>Oreophryne gagneorum</i>	Kraus, 2013	<i>Oreophryne</i>	Au
<i>Oreophryne insulana</i>	Zweifel, 1956	<i>Oreophryne</i>	Au
<i>Oreophryne matawan</i>	Kraus, 2016	<i>Oreophryne</i>	Au
<i>Oreophryne meliades</i>	Kraus, 2016	<i>Oreophryne</i>	Au
<i>Oreophryne penelopeia</i>	Kraus, 2016	<i>Oreophryne</i>	Au
<i>Oreophryne phoebe</i>	Kraus, 2017	<i>Oreophryne</i>	Au
<i>Oreophryne picticrus</i>	Kraus, 2016	<i>Oreophryne</i>	Au
<i>Sphenophryne variabilis</i>	Boulenger, 1896	<i>Oreophryne</i>	Au
<i>Oreophryne albitypanum</i>	Günther, Richards, and Tjaturadi, 2018	<i>Oreophryne</i>	inc
<i>Oreophryne albomaculata</i>	Günther, Richards, and Dahl, 2014	<i>Oreophryne</i>	inc
<i>Sphenophryne albopunctata</i>	Van Kampen, 1909	<i>Oreophryne</i>	inc

Original designation	Citation	Previous generic placement	Cu pla
<i>Oreophryne alticola</i>	Zweifel, Cogger, and Richards, 2005	<i>Oreophryne</i>	inc
<i>Oreophryne ampelos</i>	Kraus, 2011	<i>Oreophryne</i>	inc
<i>Oreophryne anser</i>	Kraus, 2016	<i>Oreophryne</i>	inc
<i>Sphenophryne anthonyi</i>	Boulenger, 1897	<i>Oreophryne</i>	inc
<i>Oreophryne asplenicola</i>	Günther, 2003	<i>Oreophryne</i>	inc
<i>Oreophryne atrigularis</i>	Günther, Richards, and Iskandar, 2001	<i>Oreophryne</i>	inc
<i>Oreophryne banshee</i>	Kraus, 2016	<i>Oreophryne</i>	inc
<i>Oreophryne brevicrus</i>	Zweifel, 1956	<i>Oreophryne</i>	inc
<i>Oreophryne brevirostris</i>	Zweifel, Cogger, and Richards, 2005	<i>Oreophryne</i>	inc
<i>Sphenophryne celebensis</i>	Müller, 1894	<i>Oreophryne</i>	inc
<i>Oreophryne choerophrynoides</i>	Günther, 2015	<i>Oreophryne</i>	inc
<i>Oreophryne chlorops</i>	Günther, Iskandar, and Richards, 2023	<i>Oreophryne</i>	inc

Original designation	Citation	Previous generic placement	Cu pla
<i>Oreophryne clamata</i>	Günther, 2003	<i>Oreophryne</i>	inc
<i>Cophixalus crucifer</i>	Van Kampen, 1913	<i>Oreophryne</i>	inc
<i>Oreophryne curator</i>	Günther, Richards, and Dahl, 2014	<i>Oreophryne</i>	inc
<i>Oreophryne flava</i>	Parker, 1934	<i>Oreophryne</i>	inc
<i>Oreophryne flavomaculata</i>	Günther and Richards, 2016	<i>Oreophryne</i>	inc
<i>Callula frontifasciata</i>	Horst, 1883	<i>Oreophryne</i>	inc
<i>Oreophryne furu</i>	Günther, Richards, Tjaturadi, and Iskandar, 2009	<i>Oreophryne</i>	inc
<i>Oreophryne geminus</i>	Zweifel, Cogger, and Richards, 2005	<i>Oreophryne</i>	inc
<i>Oreophryne habbemensis</i>	Zweifel, Cogger, and Richards, 2005	<i>Oreophryne</i>	inc
<i>Oreophryne hypsiops</i>	Zweifel, Menzies, and Price, 2003	<i>Oreophryne</i>	inc
<i>Oreophryne idenburgensis</i>	Zweifel, 1956	<i>Oreophryne</i>	inc
<i>Oreophryne jeffersoniana</i>	Dunn, 1928	<i>Oreophryne</i>	inc

Original designation	Citation	Previous generic placement	Cu pla
<i>Oreophryne kampeni</i>	Parker, 1934	<i>Oreophryne</i>	inc
<i>Oreophryne kapisa</i>	Günther, 2003	<i>Oreophryne</i>	inc
<i>Sphenophryne mertoni</i>	Roux, 1910	<i>Oreophryne</i>	inc
<i>Oreophryne minuta</i>	Richards and Iskandar, 2000	<i>Oreophryne</i>	inc
<i>Sphenophryne monticola</i>	Boulenger, 1897	<i>Oreophryne</i>	inc
<i>Oreophryne nicolasi</i>	Richards and Günther, 2019	<i>Oreophryne</i>	inc
<i>Oreophryne oviprotector</i>	Günther, Richards, Bickford, and Johnston, 2012	<i>Oreophryne</i>	inc
<i>Oreophryne pseudasplenicola</i>	Günther, 2003	<i>Oreophryne</i>	inc
<i>Oreophryne pseudunicolor</i>	Günther and Richards, 2016	<i>Oreophryne</i>	inc
<i>Oreophryne purari</i>	Günther, Nagombi, and Richards, 2025	<i>Oreophryne</i>	inc
<i>Oreophryne roedeli</i>	Günther, 2015	<i>Oreophryne</i>	inc
<i>Oreophryne rookmaakeri</i>	Mertens, 1927	<i>Oreophryne</i>	inc

Original designation	Citation	Previous generic placement	Cu pla
<i>Oreophryne riyantoi</i>	Putri, Trilaksono, Kurniati, Engilis, and Hamidy, 2023	<i>Oreophryne</i>	inc
<i>Oreophryne sibilans</i>	Günther, 2003	<i>Oreophryne</i>	inc
<i>Oreophryne streiffeleri</i>	Günther and Richards, 2012	<i>Oreophryne</i>	inc
<i>Oreophryne terrestris</i>	Zweifel, Cogger, and Richards, 2005	<i>Oreophryne</i>	inc
<i>Oreophryne unicolor</i>	Günther, 2003	<i>Oreophryne</i>	inc
<i>Oreophryne waira</i>	Günther, 2003	<i>Oreophryne</i>	inc
<i>Oreophryne wapoga</i>	Günther, Richards, and Iskandar, 2001	<i>Oreophryne</i>	inc
<i>Hylella wolterstorffi</i>	Werner, 1901	<i>Oreophryne</i>	inc
<i>Oreophryne zimmeri</i>	Ahl, 1933	<i>Oreophryne</i>	inc

Citations

Ahl E. 1933. Ergebnisse der celebes-und halmaheira-expedition heinrich 1930–32.

Mitteilungen aus dem Zoologischen Museum in Berlin **19**: 577–583.

Alcala AC. 1986. [Guide to philippine flora and fauna: Amphibians and reptiles](#) (Natural Resources Management Center (Philippines), Ed.). Natural Resources Management Center, Ministry of Natural Resources; University of the Philippines.

Balinton RB & Seña CA de la. 2015. [Karyotype of a rare minute frog, oreophryne cf. Anulata \(anura: Microhylidae\), in agusan marsh, mindanao, philippines.](#) The Herpetological Bulletin.

Boettger O. 1895. Liste der reptilien und batrachier der insel halmaheira nach den sammlungen prof. Dr. W. Kükenthal's. Zoologischer Anzeiger **18**: 129–138.

Boulenger GA. 1896. [Descriptions of new reptiles and batrachians obtained by mr. Alfred everett in celebes and jampea.](#) The Annals and magazine of natural history; zoology, botany, and geology, Series 6 **18**: 62–64.

Boulenger GA. 1897. Descriptions of new lizards and frogs from mount victoria, owen stanley range, new guinea, collected by mr. A. S. anthony. Annals and Magazine of Natural

History, Series 6 **19**: 6–13.

Brown RM, Siler CD, Richards SJ, Diesmos AC & Cannatella DC. 2015. [Multilocus phylogeny and a new classification for southeast asian and melanesian forest frogs \(family ceratobatrachidae\)](#). Zoological Journal of the Linnean Society **174**: 130–168.

De Queiroz K & Cantino P. 2020. [International code of phylogenetic nomenclature \(PhyloCode\)](#). CRC Press.

De Queiroz K & Gauthier J. 1990. [Phylogeny as a central principle in taxonomy - phylogenetic definitions of taxon names](#). Systematic Zoology **39**: 307–322.

De Queiroz K & Gauthier J. 1992. [Phylogenetic taxonomy](#). Annual Review of Ecology, Evolution, and Systematics **23**: 449–480.

De Queiroz K & Gauthier J. 1994. [Toward a phylogenetic system of biological nomenclature](#). Trends in Ecology & Evolution **9**: 27–31.

De Sa RO, Streicher JW, Sekonyela R, Forlani MC, Loader SP, Greenbaum E, Richards S & Haddad CFB. 2012. [Molecular phylogeny of microhylid frogs \(Anura: Microhylidae\) with emphasis on relationships among New World genera](#). BMC Evolutionary Biology **12**.

Dunnington D. 2023. [Ggspatial: Spatial data framework for ggplot2](#).

Edgar RC. 2004. [Muscle: A multiple sequence alignment method with reduced time and space complexity](#). BMC Bioinformatics **5**: 113.

Engelkes K, Kath L, Kleinteich T, Hammel JU, Beerlink A & Haas A. 2020. [Ecomorphology of the pectoral girdle in anurans \(amphibia, anura\): Shape diversity and biomechanical considerations](#). ECOLOGY AND EVOLUTION **10**: 11467–11487.

Frost DR. 2024. [Amphibian species of the world: An online reference. Version 6.2](#). American Museum of Natural History, New York, USA.

Gillman LN & Wright SD. 2020. [Restoring indigenous names in taxonomy](#). Communications Biology **3**: 609.

Günther R, Iskandar DT & Richards SJ. 2023. [A new large oreophryne species from the mountains of papua province, indonesian new guinea \(amphibia, anura, microhylidae\)](#). Vertebrate Zoology **73**: 153–159.

Günther R & Richards SJ. 2012. Five new microhylid frog species from enga province, papua new guinea, and remarks on albericus alpestris (anura, microhylidae). VERTEBRATE ZOOLOGY **61**: 343–372.

Günther R, Richards S & Iskandar DT. 2001. Two new species of the genus oreophryne from irian jaya, indonesia (amphibia, anura, microhylidae). [Spixiana](#) **24**: 257–274.

Hill EC, Fraser CJ, Gao DF, Jarman MJ, Henry ER, Iova B, Allison A & Butler MA. 2022. [Resolving the deep phylogeny: Implications for early adaptive radiation, cryptic, and present-day ecological diversity of papuan microhylid frogs.](#) [Molecular Phylogenetics and Evolution](#) **177**.

Hill EC, Jarman MJ, Fraser CJ, Gao DF, Henry ER, Fisher AR, Iova B, Allison A & Butler MA. 2023B. [Molecular and phylogenetic datasets for the asterophryinae frogs of new guinea with additional data on lifestyle, geography, and elevation.](#) [Data in Brief](#) **47**: 108987.

Hill EC, Gao DF, Polhemus DA, Fraser CJ, Iova B, Allison A & Butler MA. 2023A. Testing geology with biology: Plate tectonics and the diversification of microhylid frogs in the papuan region. [Integrative Organismal Biology](#).

International Commission on Zoological Nomenclature. 1999. [International Code of Zoological Nomenclature](#). London, International Trust for Zoological Nomenclature, c/o Natural History Museum, 1999.

IUCN. 2025. [Asterophryinae \(spatial data\).](#) [The IUCN Red List of Threatened Species. Version 2025-1.](#) [International Union for Conservation of Nature](#).

Köhler F & Günther R. 2008. The radiation of microhylid frogs (amphibia : Anura) on new guinea: A mitochondrial phylogeny reveals parallel evolution of morphological and life history traits and disproves the current morphology-based classification. Molecular Phylogenetics and Evolution **47**: 353–365.

Kraus F. 2013. Three new species of *oreophryne* (anura, microhylidae) from papua new guinea. ZOOKEYS: 93–121.

Kraus F. 2016. Ten new species of oreophryne (anura, microhylidae) from papua new guinea. Zootaxa **4195**: 1–68.

Kraus F. 2017. A new insular species of oreophryne (anura, microhylidae) from papua new guinea. Journal of Herpetology **51**: 552–558.

Kraus F & Allison A. 2004. Notes on poorly known new guinea frogs. I. Microhylids from the d'entrecasteaux islands. HERPETOFAUNA-SYDNEY- **34**: 26–39.

Kraus F & Allison A. 2009. New microhylid frogs from the muller range, papua new guinea. ZooKeys **26**: 53–76.

Lanfear R, Calcott B, Ho SYW & Guindon S. 2012. PartitionFinder: Combined Selection of Partitioning Schemes and Substitution Models for Phylogenetic Analyses. Molecular

Biology and Evolution **29**: 1695–1701.

Maddison WP & Maddison DR. 2023. [Mesquite: A modular system for evolutionary analysis](#).

Massicotte P & South A. 2023. [Rnaturalearth: World map data from natural earth](#).

Menzies J. 2006. The frogs of new guinea and the solomon islands. Pensoft Publishers.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Haeseler A von & Lanfear R. 2020. [IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era](#). Molecular Biology and Evolution **37**: 1530–1534.

Müller F. 1894. Reptilien und amphibien aus celebes. Verhandlungen der Naturforschenden Gesellschaft in Basel **10**: 825–843.

Parker HW. 1934. A monograph of the frogs of the family microhylidae. London: Trustees of the British Museum.

Pebesma E. 2018. [Simple Features for R: Standardized Support for Spatial Vector Data](#). The R Journal **10**: 439–446.

Peters WCH & Doria G. 1878. Catalogo dei rettili e dei batraci raccolti da o. Beccari, l.m. D'albertis a.a. Bruijn nella sotto-regione austro-malese. Annali del Museo civico di storia naturale di Genova **13**: 323–450.

Putri A, TRILAKSONO W, KURNIATI H, Hitch A, Jr A, Widayati K, Farajallah A & Hamidy A. 2023. [A new high elevation species of oreophryne boettger \(anura: Microhylidae\) from sulawesi, indonesia.](#) Zootaxa **5353**: 455–467.

R Core Team. 2024. [R: A language and environment for statistical computing.](#) Vienna, Austria: R Foundation for Statistical Computing.

Rivera JA, Kraus F, Allison A & Butler MA. 2017. [Molecular phylogenetics and dating of the problematic new guinea microhylid frogs \(amphibia: Anura\) reveals elevated speciation rates and need for taxonomic reclassification.](#) Molecular Phylogenetics and Evolution **112**: 1–11.

Setiadi MI, Hamidy A, Abidin Z, Susanto D, Brown RM, Peterson AT, Li X & Evans BJ. 2010. [Genetic structure of herpetofauna on halmahera island, indonesia: Implications for aketajawe-lolobata national park.](#) Conservation Biology **24**: 553–562.

Soliz MC, Abdala V & Tulli MJ. 2025. [The ecological drivers of variation in pectoral girdle anatomy in frogs.](#) Acta Zoologica **106**: 38–54.

Tu N, Yang M, Liang D & Zhang P. 2018. [A large-scale phylogeny of microhylidae inferred from a combined dataset of 121 genes and 427 taxa.](#) Mol Phylogenet Evol **126**: 85–91.

Werner Franz. 1898. [Vorläufige mitteilung über die von herrn prof. F. Dahl im bismarck-archipel gesammelten reptilien und batrachier.](#) Zoologischer Anzeiger **21**: 552–556.

Wickham H. 2016. [ggplot2: Elegant graphics for data analysis.](#) Springer-Verlag New York.

Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S & Madden TL. 2012. [Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction.](#) BMC Bioinformatics **13**: 134.

Yu G, Smith D, Zhu H, Guan Y & Lam TTY. 2017. [Ggtree: An r package for visualization and annotation of phylogenetic trees with their covariates and other associated data.](#) Methods in Ecology and Evolution **8**: 28–36.

Yu G. 2020. [Using ggtree to visualize data on tree-like structures.](#) Current Protocols in Bioinformatics **69**: e96.

Zweifel RG. 1956. [Microhylid frogs from new guinea, with descriptions of new species.](#) American Museum Novitates: 1–49.

Zweifel RG. 2003. A new species of microhylid frog, genus oreophryne, from papua new guinea. American Museum Novitates: 1–8.

Zweifel RG, Cogger H & Richards S. 2005. [Systematics of Microhylid Frogs, Genus Oreophryne, Living at High Elevations in New Guinea.](#) American Museum Novitates **3495**: 1–25.

Zweifel R, Menzies J & Price D. 2003. [Systematics of microhylid frogs, genus oreophryne, from the north coast region of new guinea.](#) AMERICAN MUSEUM NOVITATES: 1–31.