Britannica Volume 144850379 Row 1942

Encyclopaedia Britannica (1771)

Historical source

Entry ID: `entry-britannica-volume-144850379-row-1942-1942`
Volume ID: 144850379
Character Count: 6060

Extracted Text

PHAR MACY.

Prepara- tions.

brought from Italy, are prepared in this manner. The odorous parts may be again separated from the oil, and transferred into water or fpirit, by distillation with those liquors.

Sect: IV. _Conferration of recent Vegetables and_ _their Infusions, etc. by Sugar and Honey._

ğ 1. CONSERVES.

454. CONSERVES are compollions of recent vege table matters and fugar beaten together into an uniform mafs.

This management was introduced for preferving certain samples, undried, in an agreeable form, with as little attention to poblin in their native viruses; and to fome subjects it is very advantageously applied. Vegetables, whole virus use all or delivered in dry- ing, except in this form be kept unjusted for a length of time; For, by carefully focusing the mouth of the containing wells, the alteration, as well as diffusion, of their either principles, is generally prevented, and the fugar preserves them from the corruption which joley vegetable would otherwise undergo.

There are, however, fundry vegetables whole vi- tous are impaired by this treatment. Muchaginous fubilances, by long lying with fugar, become leis glu- tionus, and thatantly fully follow them the pa- late. Many of the fingers flowers are of to tender and delicate a texture, as almost entirely to lode their peculiar qualities on being been on brushed in general. It is obvious, that in this form, on ac- count of the large admixture of fugar, only fulbliances of considerable activity can be taken to advantage as medicines. And indeed, conferve are at perfect con- fidered chiefly as auxiliaries to medicines of greater efficacy, as interventions for joining them together. They are very convenient for echoing into boubles or pills the more pondrous powders, as mecurunu dul- cies, the colors of iron, and other mineral preparations; which will liquid or leis confident matters, as fyrroup, will not cohere.

The flops were formerly ecumulated with many conferences altogether insignificant; the few now retained have in general either an agreeable flavour to re- commend them, or are capable of answering some ofeful purpose as medicine. Their common dole is the bulk of a nutmey, or as much as can be taken up at once or twice upon the point of a knife. There is in general no great danger of exceeding in this parti- cular.

455. _General method of preferring conferences_.

Leaves are picked from the flalks, and flowers from their crops. They are beaten in a marble mortar, with a wooden peille, into a smooth mafs after which three there weight of double-refined fugar is added by degrees, and the beating continued till they are uniformly mixed.

The fugar should be pulverified by infilt, and patfed through a fleeve, before it is mixed with the vegetable mafs; otherwise it cannot easily be reduced to folfix- cleine friends, fo as to be daily incorporated. Some vegetables are cause reducible to the requisite finerals by beating in a mortar; fuch is orange-peel. This is molt conveniently raffed or grated off from the finis, then well mixed with the fugar, and the com-

Part II.

pound fet by in a clole veffel for fome weeks; after which it may be beaten Smooth with confidierably leis labour than at frift. This peel, and red role-buds, are commonly ground in a wooden mill made for that purpole.

456. _Conferve of the leaves of garden fuancy-grafs_. L.

This is the only form that feury-grafs in futultance can be kept in without the total line of its virtues. The conferve retains the full suite and virtue of the herb for a very conferable length of time, as a year two, provided the veffel be made perfectly cleo and ice in a cool place. It must be given in Ecurbotic habits three or four times a-day, so efficient.

437. _Conferve of the leaves of wood-forrel_. L.

This is a very elegant and grateful conferve y in talte it is lightly cushion, with a peculiar flavory which home refranchise totals of green-taa. It is taken because for quenching thirk, and cooling the mouth and fauces, in hot distempers. It may be uife- fully joined to the foregoing preparation, whole vic- ture to fomelan promotes, at the same time that it improves the taille.

458. _Conferve of the leaves of jewarmist_.

The conferve of mini retains the safe and virtues of the herb. It is given in weakouts of the stomach and perhaps to omit; and not unfrequently deea fervice in fome cafes of this kind, where the warmer and more active preparations of mini would be leis proper.

459. _Conferve of the leaves of rugs_. L.

This conferve is given from a dram to half an ounce in crudities of the prime via, for promoting digestion, and in by their disorders; I gently illuminate the fo- lids, attenuates vital juices, and excites the natural feecretions. Some have had a great opinion of it, ta- ken in a morning, as a preferative against the effects of contagious air or calitation.

460. _Conferve of the tops of jau-mesorwood_. L.

The conferve of wormwood has been celebrated in droplets. Mathihulus relates, that several persons were cured by it of that diltemper, without the diffiance of any other medicine. Where the diforder indeed proceeds from a simple laxity or flagidity of the fo- lids, the contained use of this medicine may be of bone ferives, as it appears to be a not malignant mild cor- robranant. It is directed to be given in the dole of half an ounce, about three hours before meals.

461. _Conferve of the buds of red refer_.

This is a very apparent and useful concre. A dram or two, diluted in warm milk, are frequently given as a light refrigerant in weakness of the blo- month, and likewise in couple and phithical com- plaints. In the German Ephemerides, examples are related of very dangerous phihifies cured by the con- tinned out of this medicine; in one of the cafes, zo pounds of the conferve were taken in the face of a month; and in another, upwards of go. Riverius mentions federal other influences of this kind.

461. _Conferve of refractory-flowers_. L.

Rofernary-flowers in great incature loft their pecu- liar

Source record

- **Dataset:** `NationalLibraryOfScotland/encyclopaedia_britannica_illustrated`
- **Dataset Revision:** `309f765b1ecc83e0a7f7762e44b4a927a7555443`
- **Page ID:** `britannica:309f765b1ecc83e0a7f7762e44b4a927a7555443:train:1942`
  Filename: `page-1942-99b5dc3b4c6c.jpg`
  Row Index: 1942
  SHA256: `99b5dc3b4c6cd19e...`

Preparations.

brought from Italy, are prepared in this manner. The odorous parts may be again ſeparated from the oil, and transferred into water or ſpirit, by diſtillation with thoſe liquors.

Sect. IV. *Conſervation of recent Vegetables and their Infuſions, &c. by Sugar and Honey.*

§ 1. Conserves.

454. Conserves are compoſitions of recent vegetable matters and ſugar beaten together into an uniform maſs.

This management was introduced for preſerving certain ſimples, undried, in an agreeable form, with as little alteration as poſſible in their native virtues; and to ſome ſubjects it is very advantageouſly applied. Vegetables, whoſe virtues are loſt or deſtroyed in drying, may in this form be kept uninjured for a length of time: for, by carefully ſecuring the mouth of the containing veſſel, the alteration, as well as diſſipation, of their active principles, is generally prevented; and the ſugar preſerves them from the corruption which juicy vegetables would otherwiſe undergo.

There are, however, ſundry vegetables whoſe virtues are impaired by this treatment. Mucilaginous ſubſtances, by long lying with ſugar, become leſs glutinous, and aſtringents ſenſibly ſofter upon the palate. Many of the fragrant flowers are of ſo tender and delicate a texture, as almoſt entirely to loſe their peculiar qualities on being beaten or bruiſed.

In general, it is obvious, that in this form, on account of the large admixture of ſugar, only ſubſtances of conſiderable activity can be taken to advantage as medicines. And indeed, conſerves are at preſent conſidered chiefly as auxiliaries to medicines of greater efficacy, or as intermediums for joining them together. They are very convenient for reducing into boluſes or pills the more ponderous powders, as mercurius dulcis, the calces of iron, and other mineral preparations; which with liquid or leſs conſiſtent matters, as ſyrups, will not cohere.

The ſhops were formerly encumbered with many conſerves altogether inſignificant; the few now retained have in general either an agreeable flavour to recommend them, or are capable of anſwering ſome uſeful purpoſes as medicines. Their common doſe is the bulk of a nutmeg, or as much as can be taken up at once or twice upon the point of a knife. There is in general no great danger of exceeding in this particular.

455. *General method of preſerving conſerves.*

Leaves are picked from the ſtalks, and flowers from their cups. They are beaten in a marble mortar, with a wooden peſtle, into a ſmooth maſs; after which thrice their weight of double-refined ſugar is added by degrees, and the beating continued till they are uniformly mixed.

The ſugar ſhould be pulveriſed by itſelf, and paſſed through a ſieve, before it is mixed with the vegetable maſs; otherwiſe it cannot eaſily be reduced to ſufficient fineneſs, ſo as to be duly incorporated. Some vegetables are ſcarce reducible to the requiſite fineneſs by beating in a mortar; ſuch is orange-peel. This is moſt conveniently raſped or grated off from the fruit, then well mixed with the ſugar, and the compound ſet by in a cloſe veſſel for ſome weeks; after which it may be beaten ſmooth with conſiderably leſs labour than at firſt. This peel, and red roſe-buds, are commonly ground in a wooden mill made for that purpoſe.

Preparations.

456. *Conſerve of the leaves of garden ſcurvy-graſs.* L.

This is the only form that ſcurvy-graſs in ſubſtance can be kept in without the total loſs of its virtues. The conſerve retains the full taſte and virtue of the herb for a very conſiderable length of time, as a year or two, provided the veſſel be made perfectly cloſe and ſet in a cool place. It may be given in ſcorbutic habits three or four times a-day, or oftener.

457. *Conſerve of the leaves of wood-ſorrel.* L.

This is a very elegant and grateful conſerve; in taſte it is lightly acidulous, with a peculiar flavour, which ſome reſemble to that of green-tea. It is taken occaſionally for quenching thirſt, and cooling the mouth and fauces, in hot diſtempers. It may be uſefully joined to the foregoing preparation, whoſe virtue it ſomewhat promotes, at the ſame time that it improves the taſte.

458. *Conſerve of the leaves of ſpearmint.* L.

The conſerve of mint retains the taſte and virtues of the herb. It is given in weakneſs of the ſtomach and retchings to vomit; and not unfrequently does ſervice in ſome caſes of this kind, where the warmer and more active preparations of mint would be leſs proper.

459. *Conſerve of the leaves of rue.* L.

This conſerve is given from a dram to half an ounce in crudities of the primæ viæ, for promoting digeſtion, and in hyſteric diſorders: it gently ſtimulates the ſolids, attenuates viſcid juices, and excites the natural ſecretions. Some have had a great opinion of it, taken in a morning, as a preſervative againſt the effects of contagious air or exhalation.

460. *Conſerve of the tops of ſea-wormwood.* L.

The conſerve of wormwood has been celebrated in dropſies: Matthiolus relates, that ſeveral perſons were cured by it of that diſtemper, without the aſſiſtance of any other medicine. Where the diſorder indeed proceeds from a ſimple laxity or flaccidity of the ſolids, the continued uſe of this medicine may be of ſome ſervice, as it appears to be a not inelegant mild corroborant. It is directed to be given in the doſe of half an ounce, about three hours before meals.

461. *Conſerve of the buds of red roſes.* L.

This is a very agreeable and uſeful conſerve. A dram or two, diſſolved in warm milk, are frequently given as a light reſtringent in weakneſs of the ſtomach, and likewiſe in coughs and phthiſical complaints. In the German Ephemerides, examples are related of very dangerous phthiſes cured by the continued uſe of this medicine: in one of theſe caſes, 20 pounds of the conſerve were taken in the ſpace of a month; and in another, upwards of 30. Riverius mentions ſeveral other inſtances of this kind.

462. *Conſerve of roſemary-flowers.* L.

Roſemary-flowers in great meaſure loſe their peculiar

Figure 1: *
Source scan: page-1942-99b5dc3b4c6c.jpg