Chap VI

Encyclopaedia Britannica (1771)

Historical source

Entry ID: `entry-chap-vi-2277`
Volume ID: 192545783
Character Count: 4113

Extracted Text

A N A T O M Y.

Of the muticles 12 12, The first cervical nerves; 13 13, The Brain and femoral cervical nerves both front upon in muticles that lie on the neck, and teguration of the neck and head. 14 14. The third cervical nerves, which after feeding off (12 15, Scc) the placenta nerve to the diaphragm, supply the mufles and tegurements that lie on the fide of the neck and top of the shoulder. 16 16, The brachial plexus, formed by the fourth, fifth, forth, towards cervical, and frit dorial nerves–which lupply the mufles, and tegurement of the effective extremity. 17 17, The twelve dorial, or proper intercedul nerves, which is peut upon the intercedul muscles and tone of the large mufles which are upon the thorax. 18 18, The five lumbar part of the nerves, which puppy the lumbar and abdominal muscles, and forms of the tegurism and mufles of the inferior extremity. 10 17, The fourscoles, as preferred cranial nerve, formed by the two inferior lumber, and three superior of the os Herman. This large nerve fulpthe the general part of the medicale and teguements of the inferior extremity. 20. The thoracic plexus, formed by the eighth pair. 21 21, Branches of the follor or callic plexus, formed by the eighth pair and inter-

CHAP. VI. OF THE SENSES, AND THEIR ORGANS.

140 IN treating of the feeling, we mean to confine ourselves to the external ones of touch, triple, fineing, hearing, and vision. The word [only], when applied to their lives, fears us imply not only the forlation occurred in the limited by certain impurities made on the body, but likewise the organ defined to receive and transmit basic ingredients in the forbidden. Each of these organs being at a peculiar threat, is if dependible only of particular impretions, which will be pointed out as we proceed to define each of them fe parately.

141 In this feature of touch may be defined to be the faculty of diltinguling certain properties of bodies by the feel. In a general accepted, this definition might perhaps not properly be extended to every part of the body political of identifying (), it is commonly combined to the nervous pupils, and consequently of exciting differences in the fenision communes. But the origin of touch, like all the other feels, it not equally distinct in every part of the body, or in every subjects, being in force much more equitable than it to others.

Sect. I. _Of Touch._

collals, which lupply the flomach and chylopotic vience. 22 19. Branch of the figure and anterior mesenteric plexus, formed by the eighth pair and fifth tercedula, which lupply the chyopotic vience. With part of the organs of urine and generation. 23 23, Nerve which accompany the fermatic cord. 24 24, The hypogotic plexus, while fusplies the organs of urine and generates within the pelvis.

Fig. 2, 3, 4. Show different views of the Inferior part of the Brain, cut perpendicularly through the Middle-with, the Origin and large portions of all the Nerve which puts through the Bones of the Cranium,and the three full Cervicals.

The anterior lobe, B, The lateral lobe of the cerebellum. C, One of the lobes of the cerebellum. To Tube aunture, K, Corpus pyramidal, in the middle of the medulla oblongata. G, The corpus olivare, in the fide of the medulla oblongata. G, The medulla oblongata. H, Hamilton mediala insula.

Naxxta - 2, 3,4, 6 5 and 3 Plate of nerves. 10 16, Nervus accoltionis, which comes from–11, 12, and 13, the three first cervical nerves.

Sect. II. _Of Teife._

This feeds of safe is faced chiefly in the tongue, the fixation and figure of which are sufficiently known.

On the upper further of this organ was may observe a great number of pupils, which, on occasion of their difference in free and hays, are commonly divided into three clades. The largest are trusted towards the final of the tongue. Their main community varies from fever to nine, and they form to be mucous folicles. Thief of the second clats and tunnels that matter, and of a cylindrical shape. They are most humorous about the middle of the tongue. Those of the third clats are very minute, and of a conical shape. They are

Source record

- **Dataset:** `NationalLibraryOfScotland/encyclopaedia_britannica_illustrated`
- **Dataset Revision:** `309f765b1ecc83e0a7f7762e44b4a927a7555443`
- **Page ID:** `britannica:309f765b1ecc83e0a7f7762e44b4a927a7555443:train:2277`
  Filename: `page-2277-cdde2fc730b9.jpg`
  Row Index: 2277
  SHA256: `cdde2fc730b9b43e...`

Of the Brain and Nerves.

muſcles. 12 12, The firſt cervical nerves; 13 13, The ſecond cervical nerves; both ſpent upon the muſcles that lie on the neck, and teguments of the neck and head. 14 14, The third cervical nerves, which, after ſending off (15 15, &c.) the phrenic nerves to the diaphragm, ſupply the muſcles and teguments that lie on the ſide of the neck and top of the ſhoulder. 16 16, The brachial plexus, formed by the fourth, fifth, ſixth, ſeventh cervicals, and firſt dorſal nerves,—which ſupply the muſcles, and teguments of the ſuperior extremity. 17 17, The twelve dorſal, or proper intercoſtal nerves, which are ſpent upon the intercoſtal muſcles and ſome of the large muſcles which lie upon the thorax. 18 18, The five lumbar pairs of nerves, which ſupply the lumbar and abdominal muſcles, and ſome of the teguments and muſcles of the inferior extremity. 19 19, The ſacro-ſciatic, or poſterior crural nerve, formed by the two inferior lumbar, and three ſuperior of the os ſacrum. This large nerve ſupplies the greateſt part of the muſcles and teguments of the inferior extremity. 20, The ſtomachic plexus, formed by the eighth pair. 21 21, Branches of the ſolar or cæliac plexus, formed by the eighth pair and intercoſtals, which ſupply the ſtomach and chylopoietic viſcera. 22 22, Branches of the ſuperior and inferior meſenteric plexuſes, formed by the eighth pair and intercoſtals, which ſupply the chylopoietic viſcera, with part of the organs of urine and generation. 23 23, Nerves which accompany the ſpermatic cord. 24 24, The hypogaſtric plexus, which ſupplies the organs of urine and generation within the pelvis.

Of the Brain and Nerves.

FIG. 2, 3, 4, 5. Show different views of the Inferior part of the Brain, cut perpendicularly through the Middle—with the Origin and large portions of all the Nerves which paſs out through the Bones of the Cranium,—and the three firſt Cervicals.

A, The anterior lobe. B, The lateral lobe of the cerebrum. C, One of the lobes of the cerebellum. D, Tuber annulare. E, Corpus pyramidale, in the middle of the medulla oblongata. F, The corpus olivare, in the ſide of the medulla oblongata. G, The medulla oblongata. H, The medulla ſpinalis.

NERVES.—1 2 3 4 5 6 7 8 and 9, Pairs of nerves. 10 10, Nervus acceſſorius, which comes from—11, 12, and 13, the three firſt cervical nerves.

CHAP. VI. OF THE SENSES, AND THEIR ORGANS.

140 IN treating of the ſenſes, we mean to confine ourſelves to the external ones of *touch*, *taſte*, *ſmelling*, *hearing*, and *viſion*. The word *ſenſe*, when applied to theſe five, ſeems to imply not only the ſenſation excited in the mind by certain impreſſions made on the body, but likewiſe the organ deſtined to receive and tranſmit theſe impreſſions to the ſenſorium. Each of theſe organs being of a peculiar ſtructure, is ſuſceptible only of particular impreſſions, which will be pointed out as we proceed to deſcribe each of them ſeparately.

SECT. I. *Of Touch.*

141 THE ſenſe of touch may be defined to be the faculty of diſtinguiſhing certain properties of bodies by the feel. In a general acceptation, this definition might perhaps not improperly be extended to every part of the body poſſeſſed of ſenſibility (r), but it is commonly confined to the nervous papillæ of the cutis, or true ſkin, which, with its appendages, and their ſeveral uſes, have been already deſcribed.

The exterior properties of bodies, ſuch as their ſolidity, moiſture, inequality, ſmoothneſs, dryneſs, or fluidity, and likewiſe their degree of heat, ſeem all to be capable of making different impreſſions on the papillæ, and conſequently of exciting different ideas in the ſenſorium commune. But the organ of touch, like all the other ſenſes, is not equally delicate in every part of the body, or in every ſubject; being in ſome much more exquiſite than it is in others.

SECT. II. *Of Taſte.*

142 THE ſenſe of taſte is ſeated chiefly in the tongue; the ſituation and figure of which are ſufficiently known.

On the upper ſurface of this organ we may obſerve a great number of papillæ, which, on account of their difference in ſize and ſhape, are commonly divided into three claſſes. The largeſt are ſituated towards the baſis of the tongue. Their number commonly varies from ſeven to nine, and they ſeem to be mucous follicles. Thoſe of the ſecond claſs are ſomewhat ſmaller, and of a cylindrical ſhape. They are moſt numerous about the middle of the tongue. Thoſe of the third claſs are very minute, and of a conical ſhape. They are

(r) In the courſe of this article, mention has often been made of the ſenſibility or inſenſibility of different parts of the body: it will therefore, perhaps, not be amiſs to obſerve in this place, that many parts which were formerly ſuppoſed to poſſeſs the moſt exquiſite ſenſe, are now known to have but little or no feeling, at leaſt in a ſound ſtate; for in an inflamed ſtate, even the bones, the moſt inſenſible parts of any, become ſuſceptible of the moſt painful ſenſations. This curious diſcovery is due to the late Baron Haller. His experiments prove, that the bones, cartilages, ligaments, tendons, epidermis, and membranes (as the pleura, pericardium, dura and pia mater, perioſteum, &c.), may in a healthy ſtate be conſidered as inſenſible. As ſenſibility depends on the brain and nerves, of courſe different parts will poſſeſs a greater or leſs degree of feeling, in proportion as they are ſupplied with a greater or ſmaller number of nerves. Upon this principle it is, that the ſkin, muſcles, ſtomach, inteſtines, urinary bladder, ureters, uterus, vagina, penis, tongue, and retina, are extremely ſenſible, while the lungs and glands have only an obſcure degree of feeling.

Figure 1: *
Source scan: `page-2277-cdde2fc730b9.jpg`