


# Air Quality - Agricultural Emissions

Max Callaghan

September 4, 2017

Outdoor air pollution contributes to millions of global premature deaths annually, with most health impacts air linked to fine particulate matter (PM<sub>2.5</sub>) (Lelieveld et al., 2015). Due to atmospheric reactions between alkaline ammonia (NH<sub>3</sub>) and acidic nitrogen oxides (NO<sub>x</sub>) and sulphur dioxide (SO<sub>2</sub>), that contribute to the formation of PM<sub>2.5</sub>, an optimal air pollution mitigation strategy should consider agricultural emissions (which primarily account for ammonia) as well as those from the transport and power sectors (Wang et al., 2015; Lee et al., 2015). Ammonia's contribution to premature mortality varies by region; in parts of Europe and North-Eastern America, reductions in Ammonia emissions may generate the largest reductions in mortality, although these results are sensitive to uncertainties about the relative toxicity of PM<sub>2.5</sub> particles by source (Lee et al., 2015; Lelieveld et al., 2015). Mitigation options for ammonia emissions are higher than other sectors, but behavioural changes towards reducing meat consumption would be highly effective, and would also generate health benefits for humans, a reduction in GHG emissions, a reduction in biodiversity loss, and the improvement of water quality (Backes et al., 2016; Leip et al., 2015).

# Agricultural Emissions



# Bibliography

- Backes, A. M., Aulinger, A., Bieser, J., Matthias, V., and Quante, M. (2016). Ammonia emissions in Europe, part II: How ammonia emission abatement strategies affect secondary aerosols. *Atmospheric Environment*, 126:153–161.
- Lee, C. J., Martin, R. V., Henze, D. K., Brauer, M., Cohen, A., and Donkelaar, A. V. (2015). Response of global particulate-matter-related mortality to changes in local precursor emissions. *Environmental Science and Technology*, 49(7):4335–4344.
- Leip, a., Billen, G., Garnier, J., Grizzetti, B., Lassaletta, L., Reis, S., Simpson, D., Sutton, M. a., de Vries, W., Weiss, F., and Westhoek, H. (2015). Impacts of European livestock production: nitrogen, sulphur, phosphorus and greenhouse gas emissions, land-use, water eutrophication and biodiversity. *Environmental Research Letters*, 10(11):115004.
- Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A. (2015). The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature*, 525(7569):367–71.
- Wang, S., Nan, J., Shi, C., Fu, Q., Gao, S., Wang, D., Cui, H., Saiz-Lopez, A., and Zhou, B. (2015). Atmospheric ammonia and its impacts on regional air quality over the megacity of Shanghai, China. *Scientific reports*, 5(October):15842.