

# TTIC 31230, Fundamentals of Deep Learning

David McAllester, Autumn 2023

## Continuous Time Models of SGD

Gradient Flow

The Diffusion SDE

The Langevin SDE

General SDEs

The SGD SDE

# Gradient Flow

Gradient flow is a non-stochastic (**deterministic**) model of **stochastic** gradient descent (SGD).

Gradient flow is defined by the **total gradient** differential equation

$$\frac{d\Phi}{dt} = -g(\Phi) \quad g(\Phi) = \nabla_{\Phi} E_{(x,y) \sim \text{Train}} \mathcal{L}(\Phi, x, y)$$

We let  $\Phi(t)$  be the solution to this differential equation satisfying  $\Phi(0) = \Phi_{\text{init}}$ .

# Gradient Flow

$$\frac{d\Phi}{dt} = -g(\Phi)$$

For small values of  $\Delta t$  this differential equation can be approximated by

$$\Delta\Phi = -g(\Phi)\Delta t$$

## Time as the Sum of the Learning Rates

Consider the update.

$$\Delta\Phi = -g\Delta t$$

Here  $\Delta t$  has both a natural interpretation as time in a numerical simulation of the flow differential equation.

But it also has a natural interpretation as a learning rate.

This leads to interpreting the sum of the learning rates as “time” in SGD.

## Gradient Flow and SGD

Consider a sequence of model parameters  $\Phi_1, \dots, \Phi_N$  produced by SGD with

$$\Phi_{i+1} = \Phi_i - \eta \hat{g}_i$$

and where  $\hat{g}_i$  is the gradient of the  $i$ th randomly selected training point.

Take  $\eta \rightarrow 0$  and  $N \rightarrow \infty$  using  $N = t/\eta$ . We will show that in this limit for SGD we have that  $\Phi_N$  converges to  $\Phi(t)$  as defined by gradient flow.

## Gradient Flow and SGD

For  $\Phi_{i+1} = \Phi_i - \eta \hat{g}_i$  we divide  $\Phi_1, \dots, \Phi_N$  into  $\sqrt{N}$  blocks.

$$(\Phi_1, \dots, \Phi_{\sqrt{N}}) (\Phi_{\sqrt{N}+1}, \dots, \Phi_{2\sqrt{N}}) \cdots (\Phi_{T-\sqrt{N}+1}, \dots, \Phi_N)$$

For  $\eta \rightarrow 0$  and  $N = t/\eta$  we have  $\eta\sqrt{N} \rightarrow 0$  which implies

$$\Phi_{\sqrt{N}} \sim \Phi_0 - \eta\sqrt{N}g$$

where  $g$  is the average (non-stochastic) gradient.

Since the gradients within each block become non-stochastic, we are back to gradient flow.

## Diffusion

Consider a discrete-time process  $z(0), z(1), z(2), z(3), \dots$  with  $z(n) \in \mathbb{R}^d$  defined by

$$\begin{aligned} z(0) &= y, \quad y \sim \text{pop}(y) \\ z(n) &= z(n) + \sigma\epsilon, \quad \epsilon \sim \mathcal{N}(0, I) \end{aligned}$$

We can sample from  $z(n)$  using

$$\begin{aligned} z(0) &= y, \quad y \sim \text{pop}(y) \\ z(n) &= z(0) + \sigma\epsilon\sqrt{n}, \quad \epsilon \sim \mathcal{N}(0, I) \end{aligned}$$

## Diffusion

Fix a numerical time step  $\Delta t$  and consider a discrete-time process  $z(0), z(\Delta t), z(2\Delta t), \dots$

$$z(0) = y, \quad y \sim \text{pop}(y)$$

$$z(t + \Delta t) = z(t) + \sigma \epsilon \sqrt{\Delta t}, \quad \epsilon \sim \mathcal{N}(0, I)$$

We now take the limit of this numerical simulation as  $\Delta t \rightarrow 0$ .

This limit defines a probability measure on the space of functions  $z(t)$ .



## The Diffusion SDE

$$z(t + \Delta t) = z(t) + \sigma \epsilon \sqrt{\Delta t}, \quad \epsilon \sim \mathcal{N}(0, I)$$

For simple diffusion (Brownian motion) this equation holds for any continuous  $t \geq 0$  and  $\Delta t \geq 0$ .

## The Langevin SDE

Consider gradient flow.

$$\frac{d\Phi(t)}{dt} = g(\Phi)$$

$$g(\Phi) = \nabla_{\Phi} \mathcal{L}(\Phi)$$

$$\mathcal{L}(\Phi) = E_{(x,y) \sim P_{\text{op}}} \mathcal{L}(\Phi, x, y)$$

## The Langevin SDE

In the Langevin SDE we add Gaussian noise to gradient flow.

$$\Phi(t + \Delta t) = \Phi(t) + g\Delta t + \sigma\epsilon\sqrt{\Delta t}, \quad \epsilon \sim \mathcal{N}(0, I)$$

We will show that the stationary distribution of Langevin Dynamics models a Bayesian posterior probability distribution on the model parameters where  $\sigma$  acts as a temperature parameter.

## The Langevin SDE

$$\Phi(t + \Delta t) = \Phi(t) + g(\Phi)\Delta t + \sigma\epsilon\sqrt{\Delta t}, \quad \epsilon \sim \mathcal{N}(0, I)$$

Let  $p(\Phi)$  be a probability density on the parameter space  $\Phi$ . The density  $p(\Phi)$  defines a gradient flow and a diffusion flow.

$$\text{gradient flow} = -p(\Phi)g(\Phi)$$

$$\text{diffusion flow} = -\frac{1}{2} \sigma^2 \nabla_{\Phi} p(\Phi)$$

The expression for the diffusion flow follows from the Fokker-Plank equation. A derivation of the diffusion flow expression from first principle is given in the appendix.

## The Langevin SDE

$$\text{gradient flow} = -p(\Phi)g(\Phi)$$

$$\text{diffusion flow} = -\frac{1}{2} \sigma^2 \nabla_{\Phi}(p(\Phi))$$

For the stationary distribution these two flows cancel each other out. In one dimension we have

$$\frac{1}{2} \sigma^2 \nabla_{\Phi} p = -p \nabla_{\Phi} \mathcal{L}$$

# The Langevin Stationary Distribution

$$\frac{1}{2}\sigma^2\nabla_{\Phi} p = -p\nabla_{\Phi}\mathcal{L}$$

$$\nabla_{\Phi} \left( \frac{1}{2}\sigma^2 \ln p \right) = \nabla_{\Phi}(-\mathcal{L})$$

$$\frac{1}{2}\sigma^2 \ln p = -\mathcal{L} + C$$

$$p(\Phi) = \frac{1}{Z} e^{\frac{-2\mathcal{L}(\Phi)}{\sigma^2}}$$

## A Bayesian Interpretation of Langevin Dynamics

If we incorporate a regularization into the loss function.

$$\mathcal{L}(\Phi) = E_{(x,y) \sim \text{Train}} [-\ln P_{\Phi}(y|x)] + \lambda ||\Phi||^2$$

Then the Gibbs distribution

$$p(\Phi) = \frac{1}{Z} e^{\frac{-2\mathcal{L}(\Phi)}{\sigma^2}}$$

can be interpreted as a Bayesian posterior density  $p(\Phi|\text{Train})$ .

## A General SDE

$$x(t + \Delta t) = f(t) + \mu(x, t)\Delta t + \sigma(x, t)\epsilon\sqrt{\Delta t}, \quad \epsilon \sim \mathcal{N}(0, I) \quad (1)$$

Here  $\sigma(x, t)$  is a matrix equal to  $\Sigma(x, t)^{1/2}$  where  $\Sigma = \sigma^2$  is the covariance matrix of the random variable  $\sigma(x, t)\epsilon$ .

This is conventionally written as

$$dx = \mu(x, t)dt + \sigma(x, t)dB \quad (2)$$

where  $B$  denotes a Wiener process (simple diffusion, aka Brownian motion)

I find (1) more intuitive than (2) but they are the same thing.



## The SGD SDE

We now consider SGD

$$\Phi_{i+1} = \Phi_i - \eta \hat{g}_i$$

Unlike gradient flow, we now hold  $\eta > 0$  fixed.

Holding  $\eta > 0$  fixed introduces gradient noise.

Gradient noise is shaped differently from the Brownian motion noise of Langevin dynamics.

## The SGD SDE

As in Gradient flow we take “time” to be the sum of the learning rates over the updates.

For  $N$  steps of SGD we define  $\Delta t = N\eta$

We consider  $\Delta t$  large enough so that  $\Delta t$  corresponds to many SGD updates.

We consider  $\Delta t$  small enough so that the gradient estimate distribution does not change over the interval  $\Delta t$ .

## Modeling Noise

The mean of  $\hat{g}$  is the true gradient  $g(\Phi)$ .

$$\begin{aligned}\Delta t &= \eta N \\ \Phi(t + \Delta t) &= \Phi(t) - \sum_{j=1}^N \eta \hat{g}_j \\ &= \Phi(t) - g(\Phi) \Delta t + \eta \sum_{j=1}^N (g(\Phi) - \hat{g}_j)\end{aligned}$$

## Modeling Noise

$$\Delta t = \Phi(t) - g(\Phi)\Delta t + \eta \sum_{j=1}^N (g(\Phi) - \hat{g}_i)$$

By the central limit theorem  $\sum_{j=1}^N (g(\Phi) - \hat{g}_i)$  is approximately Gaussian.

## Modeling Noise

$$\Delta t = \Phi(t) - g(\Phi)\Delta t + \eta \sum_{j=1}^N (g(\Phi) - \hat{g}_i)$$

Let  $\sigma(\Phi)$  be the square root of the covariance matrix of the random variable  $g - \hat{g}$  at parameter setting  $\Phi$ .

In one dimension we can approximately sample from  $\sum_{i=1}^N (g(\Phi) - \hat{g}_i)$  as

$$\sigma(\Phi)\epsilon\sqrt{N}, \quad \epsilon \sim \mathcal{N}(0, 1)$$

In high dimension this becomes

$$\sigma(\Phi)\epsilon\sqrt{N}, \quad \epsilon \sim \mathcal{N}(0, I)$$

## The SGD SDE

If the mean gradient  $g(\Phi)$  is approximately constant over the interval  $\Delta t = N\eta$  we have

$$\begin{aligned}\Phi(t + \Delta t) &\approx \Phi(t) - g(\Phi)\Delta t + \eta\sigma(\Phi)\epsilon\sqrt{N} \\ &= \Phi(t) - g(\Phi)\Delta t + \eta\sigma(\Phi)\epsilon\sqrt{\Delta t/\eta} \\ &= \Phi(t) - g(\Phi)\Delta t + \eta\sigma(\Phi)\epsilon\sqrt{\Delta t}\end{aligned}$$

This is the SGD SDE

## The SGD SDE in One Dimension

$$\Phi(t + \Delta t) = \Phi(t) - g(\Phi)\Delta t + \eta\sigma(\Phi)\epsilon\sqrt{\Delta t}$$

In one dimension, if the the gradient noise  $\sigma(\Phi)$  is constant, then the SGD SDE has the same form as Langevin dynamics and we get.

$$p(x) = \frac{1}{Z} \exp\left(\frac{-2\mathcal{L}(x)}{\eta\sigma^2}\right)$$

This is Gibbs and provides an interpretation of the learning rate as temperature.

## The SGD SDE in Higher Dimension

$$\Phi(t + \Delta t) = \Phi(t) - g(\Phi)\Delta t + \eta\sigma(\Phi)\epsilon\sqrt{\Delta t}$$

This is almost the general case of an SDE.

Here  $g(\Phi)$  is the gradient of a scalar function. This is not true for a general SDE.

But the matrix  $\sigma(\Phi)$  is arbitrary.

Here the learning rate  $\eta$  controls the level of noise but we do not in general have a Gibbs distribution.



## The SGD SDE, A Counter Example

If we have two dimensions  $x$  and  $y$  where the loss separates as  $\mathcal{L}(x, y) = \mathcal{L}(x) + \mathcal{L}(y)$ , and the matrix  $\sigma(\Phi)$  is constant and diagonal, each dimension behaves as an independent one dimensional SGD and we get.

$$p(x, y) = \frac{1}{Z} \exp \left( \frac{-2\mathcal{L}(x)}{\eta\sigma_x^2} + \frac{-2\mathcal{L}(y)}{\eta\sigma_y^2} \right)$$

This is not Gibbs.

**END**

## Appendix: Diffusion Flow

We consider the one dimensional case where we have a function  $x(t) \in \mathbb{R}$ . We consider a very small time step  $\Delta t$  and consider only the diffusion flow.

$$x(t + \Delta t) = x(t) + \sigma \epsilon \sqrt{\Delta t}, \quad \epsilon \sim \mathcal{N}(0, 1)$$

We assume a density  $p_x$  of values of  $x$  and let  $p_\epsilon(\epsilon)$  be the normal distribution  $\mathcal{N}(0, 1)$  on  $\epsilon$ .

The quantity of mass transfer in the time interval  $\Delta t$  from values above  $x$  to values below  $x$  is

$$\begin{aligned} & \int_{z=0}^{\infty} p_x(x + z) p_\epsilon(\sigma \epsilon \sqrt{\Delta t} \leq -z) dz \\ &= \int_{z=0}^{\infty} p_x(x + z) p_\epsilon \left( \epsilon \leq \frac{-z}{\sigma \sqrt{\Delta t}} \right) dz \\ &= \int_{z=0}^{\infty} p_x(x + z) \Phi \left( \frac{-z}{\sigma \sqrt{\Delta t}} \right) dz \end{aligned}$$

where  $\Phi$  is the cumulative function of the Gaussian.

## Appendix: Diffusion Flow

The quantity of mass transfer in the time interval  $\Delta t$  from values above  $x$  to values below  $x$  is

$$\int_{z=0}^{\infty} p_x(x+z) \Phi\left(\frac{-z}{\sigma\sqrt{\Delta t}}\right) dz$$

By a change of variables  $u = z/(\sigma\sqrt{\Delta t})$  we get

$$\int_{u=0}^{\infty} p_x(x + \sigma\sqrt{\Delta t} u) \Phi(-u) \sigma\sqrt{\Delta t} du$$

As  $\Delta t \rightarrow 0$  we can use the first order Taylor expansion of the density.

$$\sigma\sqrt{\Delta t} \int_{u=0}^{\infty} \left( p_x(x) + \sigma\sqrt{\Delta t} u \frac{dp_x(x)}{dx} \right) \Phi(-u) du$$

## Appendix: Diffusion Flow

$$\begin{aligned} & \sigma\sqrt{\Delta t} \int_{u=0}^{\infty} \left( p_x(x) + \sigma\sqrt{\Delta t} u \frac{dp_x(x)}{dx} \right) \Phi(-u) du \\ &= \sigma\sqrt{\Delta t} p_x(x) \left( \int_{u=0}^{\infty} \Phi(-u) du \right) + \sigma^2 \Delta t \frac{dp_x(x)}{dx} \left( \int_{u=0}^{\infty} u \Phi(-u) du \right) \end{aligned}$$

A similar analysis shows that the mass transfer from lower values to higher values is

$$= \sigma\sqrt{\Delta t} p_x(x) \left( \int_{u=0}^{\infty} \Phi(-u) du \right) - \sigma^2 \Delta t \frac{dp_x(x)}{dx} \left( \int_{u=0}^{\infty} u \Phi(-u) du \right)$$

The net mass transfer in the positive  $x$  direction is the second minus the first or

$$= -2\sigma^2 \Delta t \frac{dp_x(x)}{dx} \left( \int_{u=0}^{\infty} u \Phi(-u) du \right)$$

## Appendix: Diffusion Flow

The net mass transfer in the positive  $x$  direction is

$$-2\sigma^2\Delta t \frac{dp_x(x)}{dx} \left( \int_{u=0}^{\infty} u\Phi(-u)du \right)$$

Note that the mass transfer is proportional to  $\Delta t$ . Dividing by  $\Delta t$  gives the flow per unit time.

$$\text{Diffusion flow} = -\alpha\sigma^2 \frac{dp_x(x)}{dx} \quad \alpha = 2 \int_{u=0}^{\infty} u\Phi(-u)du$$

$\alpha$  can be calculated using integration by parts.

$$\begin{aligned} \alpha &= 2 \int_0^{\infty} u\Phi(-u)du \\ &= \int_0^{\infty} \Phi(-u)du^2 \\ &= u^2\Phi(-u)|_0^{\infty} + \int_0^{\infty} u^2\phi(-u)du \quad \text{where } \phi \text{ is the Gaussian density} \\ &= \int_0^{\infty} u^2\phi(-u)du \\ &= \frac{1}{2} \end{aligned}$$

# Appendix: Diffusion Flow

We now have that the diffusion flow is

$$\text{Diffusion flow} = -\frac{1}{2} \sigma^2 \frac{dp_x(x)}{dx}$$

For dimension larger than 1 we have

$$\text{Diffusion flow} = -\frac{1}{2} \Sigma \nabla_x p_x$$

Where  $\Sigma = E (\hat{g} - g)(\hat{g} - g)^\top$  is the covariance matrix of the gradient noise.

Here we have derived this from first principle but it also follows from the Fokker–Planck equation (see Wikipedia).

**END**