Massimiliano Canzi massimiliano.canzi@uni-konstanz.de

June 24 2021

Section 1

Four years ago...

Can event-related potential data inform information flow order in speech perception?

Can event-related potential data inform information flow order in speech perception? i.e. what the extent of top-down mediation is during speech perception.

Event-related potentials (ERP) are measured brain responses that are direct result of a **sensory**, **cognitive** or motor event (Luck 2005)

Event-related potentials (ERP) are measured brain responses that are direct result of a **sensory**, **cognitive** or motor event (Luck 2005)

Event-related potential components are measured with electroencephalography (EEG) equipment.

Top-down vs bottom-up in speech perception:

Methods 00000000 Experiment 1

Experiment 2

General Discussion 000000

Introduction

Top-down vs bottom-up in speech perception:

• Interactive models of speech perception e.g. **TRACE** (McClelland & Elman 1986)

Introduction

Top-down vs bottom-up in speech perception:

- Interactive models of speech perception e.g. **TRACE** (McClelland & Elman 1986)
- Feed-forward / modular models of speech perception e.g.
 Cohort (Marslen-Wilson 1984)

Experiment 1 000000000

Experiment 2 000000000

General Discussion 000000

 Elman, J. L., & McClelland, J. L. (1988). Cognitive penetration of the mechanisms of perception: Compensation for coarticulation of lexically restored phonemes. Journal of Memory and Language, 27(2), 143-165.

- Elman, J. L., & McClelland, J. L. (1988). Cognitive penetration of the mechanisms of perception: Compensation for coarticulation of lexically restored phonemes. Journal of Memory and Language, 27(2), 143-165.
- Pitt, M. A., & McQueen, J. M. (1998). Is compensation for coarticulation mediated by the lexicon? Journal of Memory and Language, 39, 347–370.

- Elman, J. L., & McClelland, J. L. (1988). Cognitive penetration of the mechanisms of perception: Compensation for coarticulation of lexically restored phonemes. Journal of Memory and Language, 27(2), 143-165.
- Pitt, M. A., & McQueen, J. M. (1998). Is compensation for coarticulation mediated by the lexicon? Journal of Memory and Language, 39, 347–370.
- Magnuson et al. (2003). Lexical effects on compensation for coarticulation: The ghost of Christmash past. Cognitive Science, 27(2), 285-298.

- Elman, J. L., & McClelland, J. L. (1988). Cognitive penetration of the mechanisms of perception: Compensation for coarticulation of lexically restored phonemes. Journal of Memory and Language, 27(2), 143-165.
- Pitt, M. A., & McQueen, J. M. (1998). Is compensation for coarticulation mediated by the lexicon? Journal of Memory and Language, 39, 347–370.
- Magnuson et al. (2003). Lexical effects on compensation for coarticulation: The ghost of Christmash past. Cognitive Science, 27(2), 285-298.
- McQueen, J. M. (2003). The ghost of Christmas future: didn't scrooge learn to be good?: Commentary on Magnuson et al. (2003). Cognitive Science, 27(5), 795-799.

- Elman, J. L., & McClelland, J. L. (1988). Cognitive penetration of the mechanisms of perception: Compensation for coarticulation of lexically restored phonemes. Journal of Memory and Language, 27(2), 143-165.
- Pitt, M. A., & McQueen, J. M. (1998). Is compensation for coarticulation mediated by the lexicon? Journal of Memory and Language, 39, 347–370.
- Magnuson et al. (2003). Lexical effects on compensation for coarticulation: The ghost of Christmash past. Cognitive Science, 27(2), 285-298.
- McQueen, J. M. (2003). The ghost of Christmas future: didn't scrooge learn to be good?: Commentary on Magnuson et al. (2003). Cognitive Science, 27(5), 795-799.
- McQueen, et al. (2009). No lexical-prelexical feedback during speech perception or: Is it time to stop playing those Christmas tapes?. Journal of Memory and Language, 61(1), 1-18.

Compensation for coarticulation: (Mann & Repp 1981)

Compensation for coarticulation: (Mann & Repp 1981)

 \bullet /t-k/ perceived more often as /k/ following /s/

Ganong effect (Ganong 1980)

Compensation for coarticulation: (Mann & Repp 1981)

- \bullet /t-k/ perceived more often as /k/ following /s/
- \circ /t-k/ perceived more often as /t/ following /J/

Ganong effect (Ganong 1980)

Compensation for coarticulation: (Mann & Repp 1981)

- /t-k/ perceived more often as /k/ following /s/
- \bullet /t-k/ perceived more often as /t/ following /ʃ/

Ganong effect (Ganong 1980)

 Ambiguous phonemes are solved more often with the choice that makes a word vs. a non-word

Compensation for coarticulation: (Mann & Repp 1981)

- /t-k/ perceived more often as /k/ following /s/
- \circ /t-k/ perceived more often as /t/ following /J/

Ganong effect (Ganong 1980)

- Ambiguous phonemes are solved more often with the choice that makes a word vs. a non-word
- e.g. Christma/s-f/ more often solved as Christma/s/.

111011 & 1110 010110110 (1000)

Compensation for coarticulation: (Mann & Repp 1981)

- \bullet /t-k/ perceived more often as /k/ following /s/
- \bullet /t-k/ perceived more often as /t/ following /ʃ/

Ganong effect (Ganong 1980)

- Ambiguous phonemes are solved more often with the choice that makes a word vs. a non-word
- e.g. Christma/s-\frac{1}{\sigma} more often solved as Christma/s/.
- Effect stronger at phoneme boundary.

 $Christma/s-\int//t-k/capes$

 $Christma/s-\int//t-k/capes$ Cool, huh?

Can event-related potential data inform information flow order in speech perception?

Can event-related potential data inform information flow order in speech perception? i.e. what the extent of top-down mediation is during speech perception.

 ERP (and $\operatorname{EEG})$ offer unparalleled temporal resolution,

ERP (and EEG) offer unparalleled temporal resolution, opening a direct window into cognitive processes of online language processing and speech perception.

ERP (and EEG) offer unparalleled temporal resolution, opening a direct window into cognitive processes of online language processing and speech perception.

The original goal of my thesis was that to **design** a handful of ERP experiments to investigate lexical feedback and top-down processes of speech perception.

ERP (and EEG) offer unparalleled temporal resolution, opening a direct window into cognitive processes of online language processing and speech perception.

The original goal of my thesis was that to **design** a handful of **ERP experiments to investigate lexical feedback** and top-down processes of speech perception. But how?

• Mismatch Negativity (MMN)

* Originally named Phonological Mismatch Negativity

- Mismatch Negativity (MMN)
- Phonological Mapping* Negativity (PMN)

 $\boldsymbol{*}$ Originanly named Phonological Mismatch Negativity

- Mismatch Negativity (MMN)
- Phonological Mapping* Negativity (PMN)
- N400
- $\boldsymbol{*}$ Originanly named Phonological Mismatch Negativity

- Mismatch Negativity (MMN)
- Phonological Mapping* Negativity (PMN)
- N400
- P600
- ullet Originally named Phonological Mismatch Negativity

MMN

The mismatch negativity (MMN) is a cross-sensorial ERP component often observed in frontocentral regions of the scalp between 150 and 250 ms post stimulus onset

MMN

The mismatch negativity (MMN) is a cross-sensorial ERP component often observed in frontocentral regions of the scalp between 150 and 250 ms post stimulus onset

The mismatch negativity reflects the perception of a deviant stimulus in a sequence of standard stimuli (e.g. Garrido et al., 2009)

MMN

The mismatch negativity (MMN) is a cross-sensorial ERP component often observed in frontocentral regions of the scalp between 150 and $250~\mathrm{ms}$ post stimulus onset

The mismatch negativity reflects the perception of a deviant stimulus in a sequence of standard stimuli (e.g. Garrido et al., 2009)

In the auditory domain, a deviant stimulus can be identified by differences in pitch, duration, stress and frequency range (Erlbeck et al., 2014)

Methods 000000000 Experiment 1 00000000

Experiment 2 000000000

General Discussion 000000

MMN

However, the MMN was also found to be sensitive to phonological mapping (Pulvermuller 2001) Methods 00000000 Experiment 1 00000000

Experiment 2

General Discussion

MMN

However, the MMN was also found to be sensitive to phonological mapping (Pulvermuller 2001)

• MMN to the presentation of mismatching Finnish words

MMN

However, the MMN was also found to be sensitive to phonological mapping (Pulvermuller 2001)

- MMN to the presentation of mismatching Finnish words
- No MMN in control group

The N400 (Kutas & Hillyard 1980) is part of the normal brain response to words and other meaningful stimuli.

The N400 (Kutas & Hillyard 1980) is part of the normal brain response to words and other meaningful stimuli.

nurse

The N400 (Kutas & Hillyard 1980) is part of the normal brain response to words and other meaningful stimuli.

nurse doctor

The N400 (Kutas & Hillyard 1980) is part of the normal brain response to words and other meaningful stimuli.

nurse doctor |

N400

The N400 (Kutas & Hillyard 1980) is part of the normal brain response to words and other meaningful stimuli.

nurse doctor | pizza

The N400 (Kutas & Hillyard 1980) is part of the normal brain response to words and other meaningful stimuli.

nurse doctor | pizza pineapple

The N400 (Kutas & Hillyard 1980) is part of the normal brain response to words and other meaningful stimuli.

nurse doctor | pizza pineapple

• Other paradigms include cloze-probability mismatch (e.g. Connolly and Phillips 1994)

The Phonological Mapping (or Mismatch) Negativity, \mathbf{PMN} is an event-related potential component hypothesized to index phonological mismatch and mapping

Experiment 1

Experiment 2

General Discussion

Phonological Mapping Negativity

The Phonological Mapping (or Mismatch) Negativity, **PMN** is an event-related potential component hypothesized to index phonological mismatch and mapping (e.g. Connolly and Phillips 1994; Connolly et al. 2001)

However, while some studies (e.g. Connolly and Phillips 1994) have linked the PMN to phonological mapping during the lexical selection stage of speech perception..

However, while some studies (e.g. Connolly and Phillips 1994) have linked the PMN to phonological mapping during the lexical selection stage of speech perception..

Others (e.g. Newman et al.) report that the PMN is a marker of acoustic and pre-lexical information.

Connolly and Phillips (1994)

Event-Related Potential Components Reflect Phonological and Semantic Processing of the Terminal Word of Spoken Sentences:

• The piano is out of

Connolly and Phillips (1994)

Event-Related Potential Components Reflect Phonological and Semantic Processing of the Terminal Word of Spoken Sentences:

• The piano is out of tune

Event-Related Potential Components Reflect Phonological and Semantic Processing of the Terminal Word of Spoken Sentences:

• The piano is out of tune (no mismatch)

- The piano is out of tune (no mismatch)
- The piano is out of

- The piano is out of tune (no mismatch)
- The piano is out of tuna

- The piano is out of tune (no mismatch)
- The piano is out of tuna (N400)

- The piano is out of tune (no mismatch)
- The piano is out of tuna (N400)
- The piano is out of

Connolly and Phillips (1994)

- The piano is out of tune (no mismatch)
- The piano is out of tuna (N400)
- The piano is out of pizza

- The piano is out of tune (no mismatch)
- The piano is out of tuna (N400)
- The piano is out of pizza (N400 and PMN)

Event-Related Potential Components Reflect Phonological and Semantic Processing of the Terminal Word of Spoken Sentences:

- The piano is out of tune (no mismatch)
- The piano is out of tuna (N400)
- The piano is out of pizza (N400 and PMN)
- o ...

Introduction

nods 000000 Experiment 1 000000000

Experiment 2

General Discussion 000000

Newman et al. (2003)

Phoneme deletion task to study the PMN:

Phoneme deletion task to study the PMN:

Delete /k/ from the word "clap"

Phoneme deletion task to study the PMN:

Delete /k/ from the word "clap"

lap

Phoneme deletion task to study the PMN:

Delete /k/ from the word "clap"

- lap
- aap

Introduction

Phoneme deletion task to study the PMN:

Delete /k/ from the word "clap"

- lap
- aap
- dog

Lewendon et. al (2020) suggest that the possibility exists that the PMN is an extension of either the Mismatch Negativity (MMN) or N400 components

Lewendon et. al (2020) also report that the majority of the literature on the PMN is characterized by contradictory findings and methodological limitations, e.g.

Contrasting theories of the PMN

Introduction

Lewendon et. al (2020) also report that the majority of the literature on the PMN is characterized by contradictory findings and methodological limitations, e.g.

- Contrasting theories of the PMN
- Mixed topographical locations:

Introduction

Lewendon et. al (2020) also report that the majority of the literature on the PMN is characterized by contradictory findings and methodological limitations, e.g.

- Contrasting theories of the PMN
- Mixed topographical locations:
 - Some studies report discovering the PMN in frontal and central sites, ohers in parietal / mid-line / evenly spread across the scalp.

Introduction

Lewendon et. al (2020) also report that the majority of the literature on the PMN is characterized by contradictory findings and methodological limitations, e.g.

- Contrasting theories of the PMN
- Mixed topographical locations:
 - Some studies report discovering the PMN in frontal and central sites, ohers in parietal / mid-line / evenly spread across the scalp.
- Methodological limitations:

Phonological Mapping Negativity

Introduction

Lewendon et. al (2020) also report that the majority of the literature on the PMN is characterized by contradictory findings and methodological limitations, e.g.

- Contrasting theories of the PMN
- Mixed topographical locations:
 - Some studies report discovering the PMN in frontal and central sites, ohers in parietal / mid-line / evenly spread across the scalp.
- Methodological limitations:
 - Few participants (usually < 10)

Phonological Mapping Negativity

Introduction

Lewendon et. al (2020) also report that the majority of the literature on the PMN is characterized by contradictory findings and methodological limitations, e.g.

- Contrasting theories of the PMN
- Mixed topographical locations:
 - Some studies report discovering the PMN in frontal and central sites, ohers in parietal / mid-line / evenly spread across the scalp.
- Methodological limitations:
 - Few participants (usually < 10)
 - Few trials (usually < 40)

Phonological Mapping Negativity

Introduction

Lewendon et. al (2020) also report that the majority of the literature on the PMN is characterized by contradictory findings and methodological limitations, e.g.

- Contrasting theories of the PMN
- Mixed topographical locations:
 - Some studies report discovering the PMN in frontal and central sites, ohers in parietal / mid-line / evenly spread across the scalp.
- Methodological limitations:
 - Few participants (usually < 10)
 - Few trials (usually < 40)
 - Confounding variables

• Is the PMN in response to acoustic, phonetic, phonological, lexical mapping and mismatch, none or a combination of all?

Introduction

Research questions

- Is the PMN in response to acoustic, phonetic, phonological, lexical mapping and mismatch, none or a combination of all?
- Is any other ERP component found in response to acoustic, phonetic and phonological mismatch in place of / together wih the PMN?

Why the PMN..

Why the PMN.. and why now?

Why the PMN.. and why now?

• The PMN might play an important role in future investigations of architectures of grammar (placed in between acoustic and lexical processing)

000000000000000000000

Introduction

Why the PMN.. and why now?

- The PMN might play an important role in future investigations of architectures of grammar (placed in between acoustic and lexical processing)
- Clinical studies have used the PMN as a <u>marker of phonological</u> <u>processing abilities</u> (Robson et al. 2017). However, it is not clear what processes the PMN reallys indexes.

Experiment 1

Experiment 2

General Discussion

Section 2

Methods

Experimental design

Experimental design

Three neuro-imaging experiments designed to introduce new contexts in which to probe the elicitation of the PMN ERP component.

Experimental design

Three neuro-imaging experiments designed to introduce new contexts in which to probe the elicitation of the PMN ERP component.

Experiments 1, 2 (and 3) were designed to simultaneously work independently while also being fully comparable.

Hardware:

Hardware:

• 64 active pin-type **BioSemi** electrodes

Hardware:

- 64 active pin-type **BioSemi** electrodes
- 6 (EX1 to EX6) face electrodes

Hardware:

- 64 active pin-type **BioSemi** electrodes
- 6 (EX1 to EX6) face electrodes
- BioSemi hardware (e.g. receiver)

Software:

• BioSemi Actiview


- BioSemi Actiview
- Neurobehavioral Systems' Presentation

- BioSemi Actiview
- Neurobehavioral Systems' Presentation
- MATLAB (2018b; 2019a; 2019b)

- BioSemi Actiview
- Neurobehavioral Systems' Presentation
- MATLAB (2018b; 2019a; 2019b)
- EEGLAB (Delorme & Makeig 2004)

- BioSemi Actiview
- Neurobehavioral Systems' Presentation
- MATLAB (2018b; 2019a; 2019b)
- EEGLAB (Delorme & Makeig 2004)
- ERPLAB (Lopez-Calderon & Luck, 2014)

- BioSemi Actiview
- Neurobehavioral Systems' Presentation
- MATLAB (2018b; 2019a; 2019b)
- EEGLAB (Delorme & Makeig 2004)
- ERPLAB (Lopez-Calderon & Luck, 2014)
- R (4.1) (R Core Team 2021)



Methods

000000000

Experiment 2

General Discussion

Equipment & Processing

EEG pre-processing:

• Offline average reference

- Offline average reference
- 512 Hz sampling frequency

Statistical analyses:

Statistical analyses:

• Exploratory channel-level multivariate testing with package ERP (Causeur et al. 2020) and the Adaptive Factor Adjustment (AFA) procedure (Sheu et al. 2016)

Statistical analyses:

- Exploratory channel-level multivariate testing with package ERP (Causeur et al. 2020) and the Adaptive Factor Adjustment (AFA) procedure (Sheu et al. 2016)
- Mean amplitude modelling with mixed-effect models & package lme4 (Bates et al. 2015)

Equipment

Data visualisation:

Equipment

Data visualisation:

 \bullet Grand-Average / difference ERP plots with ggplot2 (Wickham 2016)

Equipment

Data visualisation:

- Grand-Average / difference ERP plots with ggplot2 (Wickham 2016)
- Cubic spline interpolation scalp maps with package akima (Akima and Gebhardt 2020)

 $_{\rm 00000000}^{\rm Methods}$

Experiment 1 000000000

Experiment 2 000000000

General Discussion

Reproducibility



Reproducibility



Data, code and model summaries are freely available on GitHub at the repository mcanzi/phd_codedata

Reproducibility



Data, code and model summaries are freely available on GitHub at the repository mcanzi/phd codedata

PhD thesis has been submitted and will be available through open access following thesis defense (in August) and corrections.

Experiment 2

General Discussion

Section 3

Experiment 1

• Participants were trained to learn three pairs of tri-syllabic nonce words in a computerized training phase (e.g. pitabu dipida)

- Participants were trained to learn three pairs of tri-syllabic nonce words in a computerized training phase (e.g. pitabu dipida)
 - \bullet Transitional probabilities within the two items of each nonce-word pair was 1.0

- Participants were trained to learn three pairs of tri-syllabic nonce words in a computerized training phase (e.g. pitabu dipida)
 - \bullet Transitional probabilities within the two items of each nonce-word pair was 1.0
- Participants were tested on their knowledge of the experimental stimuli in a computerized task

- Participants were trained to learn three pairs of tri-syllabic nonce words in a computerized training phase (e.g. pitabu dipida)
 - Transitional probabilities within the two items of each nonce-word pair was 1.0
- Participants were tested on their knowledge of the experimental stimuli in a computerized task
- During EEG data collection, stimuli were played back to participants during a passive listening task, however...

- Participants were trained to learn three pairs of tri-syllabic nonce words in a computerized training phase (e.g. pitabu dipida)
 - Transitional probabilities within the two items of each nonce-word pair was 1.0
- Participants were tested on their knowledge of the experimental stimuli in a computerized task
- During EEG data collection, stimuli were played back to participants during a passive listening task, however...
 - In 33% of total trials (400 total trials), the first syllable of the second nonce-word of each pair would be manipulated to break expectations

pitabu

pitabu dipida

pitabu dipida pitabu

pitabu dipida pitabu **ba**pida

pitabu dipida pitabu **ba**pida pitabu

pitabu dipida pitabu **ba**pida pitabu **bu**pida

pitabu dipida pitabu **ba**pida pitabu **bu**pida

• Stimuli were synthesized using Mac OS Text-to-Speech

pitabu dipida pitabu **ba**pida pitabu **bu**pida

- Stimuli were synthesized using Mac OS Text-to-Speech
- Vowel, syllable and word length were controlled (each syllable was 200 ms long)

pitabu dipida pitabu **ba**pida pitabu **bu**pida

- Stimuli were synthesized using Mac OS Text-to-Speech
- Vowel, syllable and word length were controlled (each syllable was 200 ms long)
- Speaker and pitch contours were the same for all stimuli.

22 Participants (F = 13) took part to the experiment.

Experiment 2

Participants

22 Participants (F = 13) took part to the experiment.

• 22 right-handed adults

22 Participants (F = 13) took part to the experiment.

- ullet 22 right-handed adults
- 22 BrE speakers


- 22 Participants (F = 13) took part to the experiment.
 - 22 right-handed adults
 - ullet 22 BrE speakers
 - Age (M = 20, 18-25)

- 22 Participants (F = 13) took part to the experiment.
 - 22 right-handed adults
 - 22 BrE speakers
 - Age (M = 20, 18-25)
 - Normal or corrected to normal vision and hearing

- 22 Participants (F = 13) took part to the experiment.
 - 22 right-handed adults
 - 22 BrE speakers
 - Age (M = 20, 18-25)
 - Normal or corrected to normal vision and hearing
 - No reported use of psychoactive medications

Cubic-spline interpolation scalp maps. Mean amplitude betwee en 280 and $320~{\rm ms}$ post-stimulus onset.

Cubic-spline interpolation scalp maps. Mean amplitude betweeen 280 and 320 ms post-stimulus onset.



Results: PMN

We fitted a LMEM to mean amplitude measured between 280 and 320 ms PSO.

We fitted a LMEM to mean amplitude measured between 280 and 320 ms PSO. Condition, Region and Hemisphere were fitted as main effects as well as three-way interaction.

We fitted a LMEM to mean amplitude measured between 280 and 320 ms PSO. Condition, Region and Hemisphere were fitted as main effects as well as three-way interaction. Varying intercepts allowed for Subject

• No main effect of Condition $[F(_{1,1797}) = 0.01, \underline{p} = .89)]$

We fitted a LMEM to mean amplitude measured between 280 and 320 ms PSO. Condition, Region and Hemisphere were fitted as main effects as well as three-way interaction. Varying intercepts allowed for Subject

- No main effect of Condition $[F(_{1,1797})=0.01,\,\underline{p}=.89)]$
- No interaction of Condition & Region $[F(_{10,1797})=1.39,\,\underline{p}=.17)]$

Reesults: Other effects

• Small negative effect between 150-200 ms for mismatch condition (frontocentral) (MMN?)

Results: Other effects

- Small negative effect between 150-200 ms for mismatch condition (frontocentral) (MMN?)
- Bigger positive effect between 500-700 ms for mismatch condition (centroparietal) (**P600?**)

- Small negative effect between 150-200 ms for mismatch condition (frontocentral) (MMN?)
- Bigger positive effect between 500-700 ms for mismatch condition (centroparietal) (P600?)
- In case of a significant interaction between Condition and Region, pairwise contrasts were carried out with package emmeans (Lenth et al. 2018)

Discussion

No instance of PMN (in any of its expected forms) was found

Discussion

Possible explanations:

Experiment 2

General Discussion 000000

Discussion

Possible explanations:

• PMN is more "higher-level" than previously theorized

Experiment 2 000000000

Discussion

- PMN is more "higher-level" than previously theorized
- \bullet Methodological limitations of Exp. 1

- $\bullet\,$ PMN is more "higher-level" than previously theorized
- Methodological limitations of Exp. 1
 - Passive listening

- PMN is more "higher-level" than previously theorized
- Methodological limitations of Exp. 1
 - Passive listening
 - Possible P3a contamination?

Section 4

Experiment 2

Introduction Methods Experiment 1 Experiment 2 General Discussion 0.00000000

• Designed to be (fairly) comparable to experiment one

- \bullet Designed to be (fairly) comparable to experiment one
 - Same stimuli as Exp 1

- Designed to be (fairly) comparable to experiment one
 - Same stimuli as Exp 1
 - No lexical activation

- Designed to be (fairly) comparable to experiment one
 - Same stimuli as Exp 1
 - No lexical activation
- Includes active, behavioural tasks

- Designed to be (fairly) comparable to experiment one
 - Same stimuli as Exp 1
 - No lexical activation
- Includes active, behavioural tasks
- More streamlined

Stimuli

Same nonce words as Experiment 1

di +

di + (500 ms pause) +

di + (500 ms pause) + pi +

di + (500 ms pause) + pi + (500 ms pause) +

 $\mathrm{di}+(500\;\mathrm{ms}\;\mathrm{pause})+\mathrm{pi}+(500\;\mathrm{ms}\;\mathrm{pause})+\mathrm{da}$

$$\label{eq:constraint} \begin{array}{l} \mathrm{di} + (500 \; \mathrm{ms} \; \mathrm{pause}) + \mathrm{pi} + (500 \; \mathrm{ms} \; \mathrm{pause}) + \, \mathrm{da} \\ \\ & (4 \; \mathrm{s} \; \mathrm{pause}) \end{array}$$

$$\mathrm{di} + (500 \; \mathrm{ms} \; \mathrm{pause}) + \mathrm{pi} + (500 \; \mathrm{ms} \; \mathrm{pause}) + \mathrm{da}$$

$$(4 \; \mathrm{s} \; \mathrm{pause})$$

$$\mathrm{dipida}$$

However, in 33% of total trials

di

However, in 33% of total trials

di pi

However, in 33% of total trials

di pi da

Experiment 1

Experiment 2

General Discussion

Procedure

However, in 33% of total trials

di pi da

 \mathbf{ba} pida

20 Participants (F = 12) took part to the experiment.

20 Participants (F = 12) took part to the experiment.

 $\, \bullet \,$ 20 right-handed adults

20 Participants (F = 12) took part to the experiment.

- ullet 20 right-handed adults
- 20 BrE speakers

- 20 Participants (F = 12) took part to the experiment.
 - 20 right-handed adults
 - 20 BrE speakers
 - \bullet Age (M = 19, 18-24)

- 20 Participants (F = 12) took part to the experiment.
 - 20 right-handed adults
 - 20 BrE speakers
 - \bullet Age (M = 19, 18-24)
 - Normal or corrected to normal vision and hearing

20 Participants (F = 12) took part to the experiment.

- 20 right-handed adults
- 20 BrE speakers
- Age (M = 19, 18-24)
- Normal or corrected to normal vision and hearing
- No reported use of psychoactive medications

Results

Results

 Small negative eeffect between 75-125 ms for mismatch condition (frontal) (N1?)

Results

- Small negative eeffect between 75-125 ms for mismatch condition (frontal) (N1?)
- Small negative effect between 150-200 ms for mismatch condition (left hemisphere) (MMN? ELAN?)

General Discussion

Results

- Small negative eeffect between 75-125 ms for mismatch condition (frontal) (N1?)
- Small negative effect between 150-200 ms for mismatch condition (left hemisphere) (MMN? ELAN?)
- Bigger positive effect between 500-700 ms for mismatch condition (centroparietal) (**P600?**)

No instance of PMN (in any of its expected forms) was found

Possible explanations:

• PMN is more "higher-level" than previously theorized

Discussion

Possible explanations:

- PMN is more "higher-level" than previously theorized
- Methodological limitations of Exp. 1

Section 5

General Discussion

Contrasting findings in PMN literature cause:

• Difficulty in determining whether an observed response matches the PMN (in function and topographical distribution)

Contrasting findings in PMN literature cause:

- Difficulty in determining whether an observed response matches the PMN (in function and topographical distribution)
- Easy to mistake any component in a similar range as the PMN

The PMN appears to be linked to lexical processing more than to the processing of pre-lexical information (including acoustic and phonetic information).

The PMN appears to be linked to lexical processing more than to the processing of pre-lexical information (including acoustic and phonetic information).

• This supports earlier theories of the PMN (e.g. Connolly & Phillips 1994)

The PMN appears to be linked to lexical processing more than to the processing of pre-lexical information (including acoustic and phonetic information).

- This supports earlier theories of the PMN (e.g. Connolly & Phillips 1994)
- Goes against later interpretations (e.g. Newman et al. 2003)

The PMN appears to be linked to lexical processing more than to the processing of pre-lexical information (including acoustic and phonetic information).

- This supports earlier theories of the PMN (e.g. Connolly & Phillips 1994)
- Goes against later interpretations (e.g. Newman et al. 2003)
- How much of the PMN is in response to phonological information specifically?

The PMN appears to be linked to lexical processing more than to the processing of pre-lexical information (including acoustic and phonetic information).

- This supports earlier theories of the PMN (e.g. Connolly & Phillips 1994)
- Goes against later interpretations (e.g. Newman et al. 2003)
- How much of the PMN is in response to phonological information specifically?
 - Issues with uses of PMN in clinical settings

Earlier responses (150-200 ms) and later P600-like effects reinforce:

Earlier responses (150-200 ms) and later P600-like effects reinforce:

• Mismatch stimuli were recognised as such

Could the PMN be a later instance of the MMN?

Earlier responses (150-200 ms) and later P600-like effects reinforce:

- Mismatch stimuli were recognised as such
- Early, acoustic / phonetic mismatch

Could the PMN be a later instance of the MMN?

Earlier responses (150-200 ms) and later P600-like effects reinforce:

- Mismatch stimuli were recognised as such
- Early, acoustic / phonetic mismatch
- P600 as an index of sequence violation

Could the PMN be a later instance of the MMN?

Methodological limitations

If we consider the non-observation of the PMN as a cause of methodological limitations:

Methodological limitations

If we consider the non-observation of the PMN as a cause of methodological limitations:

 PMN most likely not reliable enough as a marker for clinical experiments

If we consider the non-observation of the PMN as a cause of methodological limitations:

- PMN most likely not reliable enough as a marker for clinical experiments
- Not a good candidate for experiments investigating information flow order in speech perception

Thank you!

Special thanks to my supervisors \mathbf{Dr} Wendell Kimper and \mathbf{Dr} Patrycja Strycharczuk