

APMA 1360 - Homework 7

Milan Capoor

21 March 2025

1 Brusselator model

Consider the system

$$\dot{x} = 1 - (b+1)x + ax^2y, \quad \dot{y} = bx - ax^2y,$$

which is a model for autocatalytic chemical reactions, where $x, y \geq 0$ and $a, b > 0$.

(i) Find all equilibria and use the Jacobian to classify them.

$$\begin{array}{rcl} 0 & = & 1 - (b+1)x + ax^2y \\ 0 & = & bx - ax^2y \\ \hline 0 & = & 1 - x \end{array}$$

Substituting $x = 1$,

$$0 = b - ay \implies y = \frac{b}{a}$$

so our only equilibrium is $\boxed{(1, b/a)}$. The Jacobian is

$$J(x, y) = \begin{pmatrix} -b-1+2axy & ax^2 \\ b-2axy & -ax^2 \end{pmatrix} \implies J(1, b/a) = \begin{pmatrix} -b-1+2b & a \\ b-2b & -a \end{pmatrix} = \begin{pmatrix} b-1 & a \\ -b & -a \end{pmatrix}$$

We know

$$\begin{aligned} \lambda_1 \lambda_2 &= -ab + a + ab = a \\ \lambda_1 + \lambda_2 &= b - 1 - a \end{aligned}$$

Since $a > 0$, $\det J > 0$ and we know there cannot be a saddle. $\text{tr } J = b - a - 1$ is

$$\begin{cases} > 0 & \text{if } b - a > 1 \\ < 0 & \text{if } b - a < 1 \end{cases} \implies \boxed{\begin{cases} \text{repeller if } b - a > 1 \\ \text{attractor if } b - a < 1 \end{cases}}$$

(ii) Show that a Hopf bifurcation occurs at some value of the parameter b (this value will depend on a).

Let

$$F(x, y) = \begin{pmatrix} 1 - (b+1)x + ax^2y \\ bx - ax^2y \end{pmatrix}$$

Clearly $F \in C^3$.

In part i, we showed that the Jacobian at the equilibrium is

$$J(1, b/a) = \begin{pmatrix} b-1 & a \\ -b & -a \end{pmatrix}$$

Notice that we could manually calculate the eigenvalues by

$$\begin{aligned}
 0 &= (b - 1 - \lambda)(-a - \lambda) + ab \\
 &= \lambda^2 + (a - b + 1)\lambda + a \\
 \lambda &= \frac{-(a - b + 1) \pm \sqrt{(a - b + 1)^2 - 4a}}{2} \\
 \lambda &= \alpha(a, b) \pm i\beta(a, b)
 \end{aligned}$$

In order for a Hopf Bifurcation to occur, we need

- (a) $\alpha(a, b) = 0$
- (b) $\beta(a, b) \neq 0$
- (c) $\frac{\partial \alpha}{\partial a} \neq 0$
- (d) $\frac{\partial \alpha}{\partial b} \neq 0$

This gives us a system of equations

$$-(a - b + 1) = 0 \tag{1}$$

$$(a - b + 1)^2 - 4a \neq 0 \tag{2}$$

$$\frac{\partial \alpha}{\partial a} = -\frac{1}{2} \neq 0 \tag{3}$$

$$\frac{\partial \alpha}{\partial b} = 1 \neq 0 \tag{4}$$

Equation (1) implies $b = a + 1$. Substituting this into (2) gives

$$(a - (a + 1) + 1)^2 - 4a = -4a < 0$$

and equations (3) and (4) are trivially satisfied.

Hence, for $\boxed{b = a + 1}$, a Hopf Bifurcation occurs around the equilibrium $(1, 1 + 1/a)$.

- (iii) Find the approximate period of the periodic orbit that bifurcates from the equilibrium at this Hopf bifurcation.

We know that the period of the periodic orbit is given by $\approx \frac{2\pi}{\beta(a, b)}$. Letting, $b = a + 1$,

$$\lambda^2 + (a - b + 1)\lambda + a = 0 \implies \lambda^2 + a = 0 \implies \lambda = \pm i\sqrt{a} \implies \beta(a, b) = \sqrt{a}$$

so the period is given by

$$\boxed{\approx \frac{2\pi}{\sqrt{a}}}$$

2 Bead on a rotating hoop revisited

Consider the second-order equation

$$\frac{d^2\phi}{dt^2} = -a\frac{d\phi}{dt} + (\gamma \cos \phi - 1) \sin \phi \quad (5)$$

that we discussed as a model for a bead that slides on a rotating hoop subject to gravitational and friction forces. The parameter a and γ are both strictly larger than zero: we keep $a > 0$ fixed and vary $\gamma > 0$.

- (i) Define the angular velocity $v := d\phi/dt$ and write (5) as a first-order system, which we refer to as equation (2), for the variables (ϕ, v) .

Let $v = d\phi/dt$. Then, we have

$$\begin{cases} \dot{\phi} = v \\ \dot{v} = -av + (\gamma \cos \phi - 1) \sin \phi \end{cases}$$

- (ii) Find the equilibria of the system (2) and determine their stability properties. Also, identify all bifurcations in this system.

We want $(\dot{\phi}, \dot{v}) = (0, 0)$ which immediately suggests $v = 0$, so

$$(\gamma \cos \phi - 1) \sin \phi = 0 \implies \begin{cases} \cos \phi = 1/\gamma \\ \sin \phi = 0 \end{cases} \implies \phi = \{\arccos \frac{1}{\gamma}, 0, \pi\}$$

where $\gamma \geq 1$ (else $\cos \phi = 1/\gamma$ has no solutions).

We have Jacobian

$$\begin{aligned} J(\phi, v) &= \begin{pmatrix} 0 & 1 \\ \gamma \cos 2\phi - \cos \phi & -a \end{pmatrix} \\ J(\arccos \frac{1}{\gamma}, 0) &= \begin{pmatrix} 0 & 1 \\ 2 - \frac{1}{\gamma} & -a \end{pmatrix} \\ J(\pi, 0) &= \begin{pmatrix} 0 & 1 \\ \gamma + 1 & -a \end{pmatrix} \\ J(0, 0) &= \begin{pmatrix} 0 & 1 \\ \gamma - 1 & -a \end{pmatrix} \end{aligned}$$

Further,

$$\det J(\arccos \frac{1}{\gamma}, 0) = 2 - \frac{1}{\gamma} > 0 \quad \forall \gamma$$

hence we have an attractor or repeller. We can check

$$\text{tr } J(\arccos \frac{1}{\gamma}, 0) = -a < 0 \quad \forall a > 0$$

which means that $\phi = \arccos \frac{1}{\gamma}$ is an attractor

Similarly,

$$\det J(\pi, 0) = -(\gamma + 1) < 0 \quad \forall \gamma$$

so $\phi = \pi$ is a saddle for all γ

Finally,

$$\det J(0, 0) = -(\gamma - 1)$$


is

$$\begin{cases} < 0 & \text{if } \gamma > 1 \\ > 0 & \text{if } \gamma < 1 \end{cases}$$

and

$$\text{tr } J(0,0) = -a < 0 \quad \forall \gamma$$

so $\phi = 0$ is a saddle for $\gamma > 1$ and an attractor $\gamma < 1$



Clearly, we have a **supercritical pitchfork bifurcation** at $\gamma = 1$.

- (iii) Compare your findings to those we obtained for the one-dimensional system where we omitted the second-order derivative $d^2\phi/dt^2$.

In the case where we omitted the second-order derivative, we had equilibria at $\phi = 0, \pi$, with

- $\phi = 0$ stable if $\gamma < 1$ and unstable if $\gamma > 1$
- $\phi = \pi$ is always unstable

and a pitchfork bifurcation at $\gamma = 1$.

In the case where we included the second-order derivative, we again have a pitchfork bifurcation at $\gamma = 1$, but now

- $\phi = 0$ is an attractor for $\gamma < 1$ and a saddle for $\gamma > 1$
- $\phi = \pi$ is a saddle
- $\phi = \arccos \frac{1}{\gamma}$ is an attractor

And this makes sense! In a sense, this is exactly the same result but in the higher dimension, we have saddles instead of unstable equilibria.